

Snow physics as relevant to snow photochemistry

**F. Domine¹, M. Albert², T. Huthwelker³, H.-W. Jacobi⁴, A. A. Kokhanovsky⁵,
M. Lehning⁶, G. Picard¹, and W. R. Simpson⁷**

¹Laboratoire de Glaciologie et Géophysique de l'Environnement, CNRS et Université Joseph Fourier, BP96, 54 rue Molière, 38402 Saint Martin d'Hères cedex, France

²Army Cold Regions Research and Engineering Lab, 72 Lyme Road, Hanover, N.H. 03755-1290, USA

³Paul Scherer Institute, Laboratory for Radiochemistry and Environmental Chemistry OFLB 106, 5232 Villigen, Switzerland

⁴Alfred Wegener Institute for Polar and Marine Research, Am Handelshafen 12, 27570 Bremerhaven, Germany

⁵Institute of Environmental Physics, Bremen University, O. Hahn Allee 1, 28334 Bremen, Germany

⁶WSL, Swiss Federal Institute for Snow and Avalanche Research, SLF Davos, Flüelastr. 11, C7260 Davos Dorf, Switzerland

⁷Department of Chemistry and Geophysical Institute, University of Alaska Fairbanks, Fairbanks, AK 99775-6160, USA

Received: 12 April 2007 – Accepted: 17 April 2007 – Published: 8 May 2007

Correspondence to: F. Domine (florent@lgge.obs.ujf-grenoble.fr)

Snow physics as
relevant to snow
photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Snow on the ground is a complex multiphase photochemical reactor that dramatically modifies the chemical composition of the overlying atmosphere. A quantitative description of the emissions of reactive gases by snow requires the knowledge of snow physical properties. This overview details our current understanding of how those physical properties relevant to snow photochemistry vary during snow metamorphism. Properties discussed are density, specific surface area, optical properties, thermal conductivity, permeability and gas diffusivity. Inasmuch as possible, equations to parameterize these properties as a function of climatic variables are proposed, based on field measurements, laboratory experiments and theory. The potential of remote sensing methods to obtain information on some snow physical variables such as grain size, liquid water content and snow depth are discussed. The possibilities for and difficulties of building a snow photochemistry model by adapting current snow physics models are explored. Elaborate snow physics models already exist, and including variables of particular interest to snow photochemistry such as light fluxes and specific surface area appears possible. On the other hand, understanding the nature and location of reactive molecules in snow seems to be the greatest difficulty modelers will have to face for lack of experimental data, and progress on this aspect will require the detailed study of natural snow samples.

1 Introduction

1.1 Context and general objectives

During the boreal winter, snow coverage includes most land masses north of 50° N, a significant fraction of land masses between 40 and 50° N, the whole Arctic ocean and Antarctica (e.g. Pielke et al., 2004), representing about 14% of the Earth's surface. Snow on the ground is a porous and permeable medium where air circulates easily

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(Albert et al., 2004). Because snow presents a large ice-air interfacial area (Legagneux et al., 2002), the potential for chemical interactions between the atmosphere and the surface is large when a snow cover is present. By analogy to the Leaf Area Index (LAI), Taillandier et al. (2006) defined the Snow Area Index (SAI) as the vertically integrated surface area of snow crystals (in m^2 of snow surface area per m^2 of land surface area). Taillandier et al. (2006) measured values of about 1000 for this dimensionless variable in the subarctic snowpack in Alaska, and Domine et al. (2002) measured values around 2000 for the Arctic snowpack. These values are much larger than for vegetation, as high LAI values are around 10 (e.g. Bremond et al., 2005). Therefore the physical properties of snow on the ground make it the surface type with the highest interaction potential with the atmosphere.

This suggestion is consistent with observations that atmospheric chemistry over snow surfaces cannot be explained without a significant contribution from snow emissions, that result for example in a 10-fold increase in OH concentrations (Domine and Shepson, 2002). Snow on the ground has been found to be the source of reactive gases such as NO_x (Jones et al., 2001; Honrath et al., 2002; Beine et al., 2002), HONO (Zhou et al., 2001; Beine et al., 2006), H_2O_2 (Hutterli et al., 1999; Jacobi et al., 2002), HCHO and other carbonyl compounds (Sumner and Shepson, 1999; Hutterli et al., 1999, 2002; Grannas et al., 2002; Jacobi et al., 2002), hydrocarbons and chloro-hydrocarbons (Swanson et al., 2002), organic acids (Dibb and Arsenault, 2002), and halogens (Spicer et al., 2002). Moreover, the complete destruction of ozone (Bottenheim et al., 1990) and the deposition of elemental mercury by the action of halogens (Steffen et al., 2002) observed in the lower polar troposphere have been traced back to processes that originate at least in part in the snow, so that it is fair to say that snow plays an important role in the composition of the planetary boundary layer over snow-covered areas.

However, elucidating the processes responsible for the quantitative understanding of these emissions remains a challenge. For example, HONO is thought to be formed from the photolysis of the NO_3^- ions contained in snow, but HONO emissions from some

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

snow-covered areas have been observed (Zhou et al., 2001), while other snow-covered regions where conditions for emissions were thought to have been met were found to produce no HONO flux (Beine et al., 2006). Part of the problem is that snow is a complex multiphase system whose chemistry cannot be treated as in a homogeneous gas phase. Thus, two reactants that are found to be present in snow by analytical techniques such as ion or liquid chromatography may be in two distinct phases and are thus not in contact. Another difficulty concerns the wide variation in physical properties such as grain size and air permeability that is found in deposited snow. If some stages of a reaction scheme requires solid phase diffusion of a species to the ice crystal surfaces, as suggested in the production of NO_x from NO_3^- photolysis (Beine et al., 2003 and 2006), areas with large-grained snow on the ground will be less productive than those with small-grains. Reactive gases produced in snow need to diffuse or be advected out of the snow. In areas with less permeable snow on the ground, the residence time may be sufficiently large to allow those gases to react with other reactants contained in snow, resulting in no observed emissions, even though production did take place.

These examples illustrate that emissions of reactive gases from snow on the ground depend on its chemical composition, on the distribution of chemical species between the different phases that form the snow, on the mobility of reactants, and on the physical properties of the snow. Figure 1 uses a very simplified scheme of NO_2 production from NO_3^- photolysis to illustrate the necessity to include snow physics in the description of its chemical emissions. One of the proposed channels for NO_3^- photolysis in snow is (Honrath et al., 1999; Jacobi and Hilker, 2007):



The rate of NO_2 production in snow interstitial air per unit volume $V_{\text{NO}_2, \text{snow}}$ is then proportional to the light actinic flux (i.e. the spherically-integrated light intensity) Φ_{light} and to the concentration of NO_3^- adsorbed onto the surface of snow crystals $[\text{NO}_3^-]_{\text{ads}}$, expressed in moles per cm^3 of snow. Φ_{light} is proportional to the light flux incident upon the snow surface Φ_{incident} and will depend on light attenuation by snow. Under certain

conditions (e.g. Simpson et al., 2002), Φ_{light} with decrease with depth z as $\exp(-z/\varepsilon)$ where ε is the snow e -folding depth. If we make the simplification that adsorbed NO_3^- comes from the adsorption of gas phase HNO_3 , then $[\text{NO}_3^-]_{\text{ads}}$ is proportional to the specific surface area of snow SSA (in cm^2/g), to the snow density ρ (in g/cm^3) and to the partial pressure of HNO_3 , P_{HNO_3} . We then have:

$$V_{\text{NO}_2, \text{snow}} \propto P_{\text{HNO}_3} \times \Phi_{\text{incident}} \times \text{SSA} \times \rho \times e^{-z/\varepsilon} \quad (1.1-2)$$

A subject of great importance to atmospheric chemistry is the snow-to-atmosphere flux of NO_2 , $F_{\text{NO}_2, \text{atm}}$. To a first approximation, this will be proportional to $V_{\text{NO}_2, \text{snow}}$, and will depend on the way NO_2 produced in the snow will be transported out of the snow. A priori, this will depend on the diffusivity of the snow if diffusive transport predominates, and on snow permeability if advective transport is more important. Overall, we have:

$$F_{\text{NO}_2, \text{atm}} \propto P_{\text{HNO}_3} \times \Phi_{\text{incident}} \times \text{SSA} \times \rho \times e^{-z/\varepsilon} \times f(K_p, D_{\text{snow}}) \quad (1.1-3)$$

Equation (1.1-3) shows that the oversimplified description of snow-to-atmosphere NO_2 fluxes from NO_3^- photolysis requires the knowledge of five physical properties of snow: specific surface area, density, e -folding depth, permeability and diffusivity. A more complete treatment would also mention that the absorption cross section and quantum yield of reaction (I-1) may be temperature-dependent, so that the snow temperature also comes into play. The snow temperature is determined by the surface energy balance and albedo, the density, the permeability, and the thermal conductivity of snow. The purpose of this paper is to present an overview of current knowledge on the physical properties of deposited snow, in order to assist modellers in their inclusion of snow physics in coupled air-snow chemical models.

Below we define the snow properties more precisely, we give a brief description of the dynamic character of snow, and we briefly list the physical properties relevant to this overview. The main section of the paper then details our knowledge of the selected physical properties, as studied in the field. The subsequent section shows how remote sensing may complement field studies in the determination of snow physical properties.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The final section discusses current snow physical models, and how these may be used and modified to contribute to a coupled air-snow chemical model.

1.2 Brief definition and description of surface snow layers

In the common vocabulary, the snow on the ground is usually called “snow”, a term that also applies to snow crystals in the atmosphere. This paper discusses seasonal snow on the ground where the terms “snowpack” and “snow cover” are commonly used, but also refers to snow on the surface of ice sheets and glaciers, where such terms are meaningless. We therefore refer to snow that has been deposited on the earth surface simply as “snow”. Our discussions do not specifically address snow falling through the atmosphere.

Because we are interested in snow-atmosphere interactions, our main focus will be surface layers, that most readily interact with the atmosphere. Defining the snow depth of interest is necessarily arbitrary. The photic zone is the depth down to which light penetrates, and is of the order of 20 cm (Simpson et al., 2002). Snowpack ventilation by wind takes place down to 50 to 100 cm (Albert and Schultz, 2002; Albert et al., 2002), so that in general our interest will be focused on the top 50 cm of the snow, hereafter called “surface snow”. When only the most superficial layers will be of interest, we will refer to “the photic zone”.

Deposited snow most often originates from the discontinuous precipitation of snow on the surface, resulting in a layered structure (Colbeck, 1991). Other processes contribute to the development of layering in snow, such as rain that subsequently freezes inside the snow, and the condensation of atmospheric water vapor on the snow surface, that form ice crystals known as surface hoar (Colbeck et al., 1990; Cabanes et al., 2002). Dry snow thus is a layered medium that is composed mostly of ice crystals and air. A property commonly used to describe the proportion of these two phases is the snow density, which varies between 0.01 g cm^{-3} or even lower for fresh dendritic snow to 0.61 g cm^{-3} for hard windpacked snow (e.g. Gray and Male, 1981; Sturm and Benson, 1997; Judson and Doesken, 2000; Domine et al., 2002 and unpublished results).

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The most common density values for snow are between 0.2 and 0.4 g cm⁻³ so that on a volume basis, snow is mostly formed of air. Snow also contains aerosol particles that have been either scavenged from the atmosphere by snow crystals during precipitation (Franz and Eisenreich, 1998; Lei and Wania, 2004), or deposited into snow on the ground by the circulation of air driven by surface wind (Cunningham and Waddington, 1993; Harder et al., 2000; Domine et al., 2004). Ice crystals within the snow also contain dissolved gases such as H₂O₂, HCHO, HCl and HNO₃ (Domine et al., 1995, Legrand et al., 1996, Hutterli et al., 2002 and 2004) that are incorporated within their crystalline lattice. Snow is thus a complex porous multiphase system formed of air, ice with dissolved gases, and any of the types of aerosols present in the lower troposphere. Snow on the ground is in contact with the atmosphere through the network of open pores. Even ice layers that are seen as thick and continuous to the naked eye have some degree of permeability (Albert et al., 2000).

On glaciers and ice sheets, where snow accumulates over many years, the open network of pore space continues down for many tens of meters as the snow evolves into firn (a term for snow older than one year) and finally becomes compressed into solid ice with only bubbles of air remaining. The depth at which there ceases to be connected pathways of interstitial air is called the depth of pore close-off, which is typically found in the range 60 to 105 m depth depending on the local climate (Barnola et al., 1991). In polar regions with low accumulation there are large areas (such as East Antarctica) where the top 50 cm is firn that is at least ten years old (Albert et al., 2004). In this paper we are concerned primarily with the top half-meter wherever it exists on the planet, and we refer to this medium as “snow”, whether it is seasonal snow or is in fact firn.

1.3 The dynamic character of snow: snow metamorphism

The transformation of snow to firn to ice illustrates the fact that snow is a dynamic medium whose physical properties need to be described as a function of time. The

crystal structure of near-surface snow evolves over time in response to changing weather. The processes responsible for physical evolution of snow are called “snow metamorphism”. When metamorphism takes place at 0°C, liquid water appears and the term “wet snow metamorphism” is used, while at temperatures below freezing, the term “dry snow metamorphism” applies (Colbeck, 1982).

In dry snow, a vertical temperature gradient almost always exists due to temperature changes at the surface. In temperate areas without permafrost, the base of the snowpack is often near 0°C, while its surface can be much colder, resulting in typical values of the temperature gradient of 10 to 30°C m⁻¹. These temperature gradients produce gradients in water vapor in the snowpack interstitial air. For example, a 1 m thick seasonal snowpack with a basal temperature of 0°C and a surface temperature of -15°C will have a water vapor pressure gradient of 444 Pa m⁻¹ that will generate upward water vapor fluxes, maintained by a net sublimation of lower layers and a net condensation in upper layers (Colbeck, 1983). Overall, it is observed that snow crystals grow during metamorphism, resulting in a decrease in their specific surface area and in the modification of other snow physical properties. These variations are complex and are also affected by other factors, such as wind that transports and compacts snow crystals and atmospheric relative humidity, that affects the snow mass balance.

When snow contains more than approximately 0.1% of water by volume, wet snow metamorphism occurs. Melting of the convex surfaces occurs first, with moisture movement to the concave (or less convex) surfaces, where it may refreeze. This process, towards an equilibrium rounded form, tends to minimize the specific surface area. When there is high liquid water content, the liquid can move easily, metamorphism proceeds rapidly, and intergranular bonds are weak. Under conditions of very low liquid water content, the liquid water is in disconnected menisci between crystals and may be immobile, slowing the metamorphism. If the bulk temperature of the snow cools and the water begins to freeze, the liquid water becomes part of the adjacent ice grains, causing rapid grain growth during melt-freeze cycling. Cold refrozen snow typically has strong intergranular bonds.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1.4 Snow physical properties relevant to photochemistry and to exchanges with the atmosphere

Many physical properties are needed to fully describe snow physics, such as mechanical resistance, which is required in avalanche research but is of little interest to this overview. The properties important for atmospheric chemistry discussed here are the following:

- Density ρ . Density is the mass of ice and water per unit volume of snow. The associated property, snow porosity ϕ , is the fraction of the snow occupied by air. With ρ_i the density of ice, porosity is given by:

$$\phi = 1 - \rho / \rho_i \quad (1.4-1)$$

- The specific surface area (SSA) is the surface area of snow crystals accessible to gases per unit mass. It is often expressed in $\text{cm}^2 \text{g}^{-1}$ and is in fact a measure of the area of the snow-air interface.
- The effective thermal conductivity of snow k_{eff} is the proportionality factor between the temperature gradient dT/dz and the heat flux q :

$$q = -k_{\text{eff}} \frac{dT}{dz} \quad (1.4-2)$$

- The air permeability of snow K_p is the proportionality factor between the pressure gradient dP/dz and the interstitial air flow velocity v_{air} :

$$v_{\text{air}} = -\frac{K_p}{\mu} \frac{dP}{dz} \quad (1.4-3)$$

where μ is the air viscosity. Equation (1.4-3) is known as Darcy's law and under this formulation K_p is called the "intrinsic permeability", but will be simply referred to as "permeability" hereafter.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- The gas diffusivity, D_g , is the proportionality factor between the gas concentration gradient and the interstitial gas flux F_{gas} :

$$F_{\text{gas}} = -\frac{D_g}{\phi} \frac{dC}{dz} \quad (1.4-4)$$

- The diffusion of gases in the snowpack can be complicated and considerably slowed down by the adsorption of gases onto snow crystal surfaces. This process will be referred to as “adsorptive diffusion” and will be briefly discussed below.
- Light intensity in snow is relevant to the calculation of photochemical reactions. It depends on snow albedo and on the ability of snow to transmit visible and UV light, this latter property being often described by its e -folding depth ε , defined above. The variable ε depends on snow scattering properties, which is affected by grain size and shape, and on snow absorption, which is determined mostly by the snow content in absorbing impurities such as carbonaceous and mineral aerosols.

2 Physical properties of snow, as determined from field studies

2.1 Preamble: variety of snow crystal shapes and of snowpacks

Snow crystals show a wide variety of sizes and shapes, caused by their conditions of formation in the atmosphere or by their metamorphic history (e.g. Colbeck et al., 1990; Wergin et al., 1996; Legagneux et al., 2002; Domine et al., 2003) and we present here a brief overview of crystal characteristics relevant to this paper. The reader is referred to Colbeck et al. (1990) for a detailed classification.

The definition of the size of a snow crystal has been ambiguous (Colbeck et al., 1990; Aoki et al., 2000), especially in the case of precipitating crystals. For example, six-branched dendritic crystals have a largest dimension of a few mm, but a thickness

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of only a few tens of microns. The problem is less complex for metamorphic crystals whose shapes are simpler. The size is often taken as either the longest dimension (adopted here), or an average characteristic dimension, and the definition varies from publication to publication. Since SSA can now be measured with a fairly good accuracy (Legagneux et al., 2002; Domine et al., 2007) it would be possible to define the average size of crystals in a snow sample, d_{ef} , by the diameter of ice spheres of the same SSA so that $d_{ef} = 6/\rho_l SSA$.

The most frequent crystals types observed on the ground are described below, with some illustrated in Fig. 2.

- Fresh snow crystals have varied shapes depending mostly on the temperature and supersaturation in the cloud where they formed (Nakaya, 1954). Common shapes include plates, dendritic crystals, graupel, columns and needles, as abundantly illustrated by Wergin et al. (1995 and 1996) and Domine et al. (2003).
- Decomposing crystals are fresh crystals in early stages of transformation, so that the initial shapes of the precipitating crystals are recognizable. Relevant pictures are reported in the above three references.
- Depth hoar crystals in seasonal snow (Fig. 2a) are the result of the metamorphism of snow of low to moderate density ($<0.35 \text{ g cm}^{-3}$) under high temperature gradient ($>20^\circ\text{C m}^{-1}$, Marbouty, 1980) that generate strong water vapor fluxes and rapid growth, resulting in hollow faceted crystals with sharp angles. Their sizes range from 2 to 20 mm. The connectivity between crystals is often very weak, resulting in very uncohesive snow.
- Faceted crystals (Fig. 2b) have flat faces with sharp angles, and are usually 0.5 to 2 mm in size. They are produced in the early stages of depth hoar formation (Sturm and Benson, 1997; Taillandier et al., 2006), or under conditions of moderate temperature gradient ($10 \text{ to } 20^\circ\text{C m}^{-1}$, Domine et al., 2002 and 2003). They are poorly interconnected and layers of such crystals are as uncohesive as depth hoar layers.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.

- 5
- 10
- 15
- 20
- Rounded grains (Fig. 2e) are produced by low temperature gradient ($<10^{\circ}\text{C m}^{-1}$) metamorphism of precipitating snow, and layers of such crystals have densities between 0.2 and 0.35 g cm^3 (Albert and Shultz, 2002; Domine et al., 2003). Crystal size range from 0.1 mm to almost 1 mm at the end of the season. They can also be produced by hard-packed wind-blown snow crystals, in which case their density can reach 0.5 (Domine et al., 2002), or exceptionally 0.61 g cm^3 , as observed near the Antarctic coast, in areas subjected to strong katabatic winds (Domine, unpublished results). Crystals in hard windpacks seldom exceed 0.4 mm in size. The interconnectivity between rounded grains is strong: these layers are cohesive and their cohesiveness increases with density.
 - Mixed forms refer to sets of crystals displaying both flat faces and rounded shapes (Fig. 2f). Their size is often in the range 0.3 to 1 mm . They can be formed under low to moderate temperature gradients, or by the rounding of faceted crystals following a decrease in the temperature gradient.
 - Melt-freeze crusts and layers are formed by the thermal cycling of snow layers, usually with diurnal melting and nocturnal freezing (Colbeck, 1982). Grains grow with each cycle, and are often observed to have a size in the range 0.5 to 2 mm (Raymond and Tusima, 1979). The melting of low density snow allows facile percolation. This can form melt-freeze layers of low density, down to 0.1 g cm^{-3} , and of moderate cohesiveness when frozen (Fig. 2d). Melting in denser layers results in dense melt-freeze crusts that have a strong cohesiveness when frozen (Fig. 2c).

25

Table 1 reports typical values of physical properties of snow layers formed of the above seven types of snow crystals. The snow on the ground is usually formed of layers of several of the above snow crystal types, and the formation of each metamorphic type is governed by surface climatic variables that include the amount of precipitation, air temperature, wind speed and insolation. There is therefore a strong correlation

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

between climate and the structure of the snow, which has motivated several classifications. Below is a brief summary of typical forms of seasonal snowpacks as described by Sturm et al. (1995). We focus on the snowpack types relating to the photochemistry investigations currently underway, Some of which are illustrated in Fig. 3.

5 The tundra snowpack (Fig. 3) is found in polar treeless areas with low precipitation (<250 mm water equivalent per year) that can be episodically swept by strong winds. It is typically 40 cm thick, and formed of a basal depth hoar layer 10 to 20 cm thick covered by a hard windpacked layer formed of small rounded grains. Variations include several windpacks, sometimes separated by layers of faceted crystals or by ice layers (Domine et al., 2002).

10 The taiga snowpack (Fig. 3) forms mostly in wind-sheltered subarctic regions with trees and low to moderate precipitation and is about 50 cm thick. This snowpack, extensively described by Sturm and Benson (1997) and Taillandier et al. (2006), is subjected to strong temperature gradients and is eventually almost fully transformed into depth hoar, except perhaps top layers deposited in late winter or spring. In the middle of winter, it is typically formed of a thick (20 cm) basal depth hoar layer surmounted by a layer of faceted crystals and episodically by a fresh snow layer.

15 The Alpine snowpack (Fig. 3) forms in fairly cold areas with abundant precipitation. The snowpack thickness prevents the establishment of the high temperature gradient necessary for depth hoar formation, except at the beginning of the season. Frequent precipitation of thick layers result in many layers of fine-grained snow not compacted by wind, separated by occasional windpacks and rare melt-freeze crusts and ice layers.

20 The maritime snowpack forms in temperate areas with abundant precipitation. The temperature gradients within this snowpack are often low. However strong gradients early in the season can induce kinetic crystal growth forms in shallow snow, and steep near-surface gradients at night can impose kinetic growth in surface snow in deep snowpacks. In maritime snow, melting events can be frequent, resulting in ice layers with grains that grew during melt-freeze cycles interspersed with layers of snow that have not experience melt (e.g. Albert and Shultz, 2002). Ice layers are common, and

Snow physics as relevant to snow photochemistryF. Domine et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

so are vertical percolation channels (e.g. Marsh and Woo, 1984; Albert et al., 1999).

The crystal structure of snow and firn on the polar ice sheets is significantly different in form than that found in temperate snowpacks described above. In the cold, high altitude regions, snow typically falls as small plates or prisms. Cold strong winds create thick layers of windpack, which is composed of small solid rounded crystals. At high-accumulation (greater than approximately 20 cm water equivalent per year) polar sites, near-surface temperature gradients and solar radiation during warmer summer conditions create surface hoar or near-surface depth hoar at the snow-air interface. If additional snowfall occurs before strong winds develop, these persist as low-density buried layers that can be used as markers of summer, even years and decades after deposition in regions where the snow does not melt (Alley et al., 1990). In addition to layering created by changing seasons, the properties of snow and firn change over time due to post-depositional metamorphism. This results in property changes that combine influences of original form with influences of grain growth, sintering, and pressure sintering over time. At high accumulation cold sites, this results in grain size that generally increases with depth, while permeability and thermal conductivity profiles take on more complicated forms (Albert and Shultz, 2002; Albert et al., 2000; Rick and Albert, 2004). In very low accumulation areas such as East Antarctica, layering exists but may reflect topographically-driven accumulation patterns rather than snowfall patterns, and large, well-sintered, faceted crystals are grown in the near surface from decades of vapor transport under very cold conditions (Albert et al., 2004).

Thus snow on the ground takes on a variety of layering and crystal properties that relate to the local climate, and the contrasting physical properties should be readily apparent to the non-specialist who is a careful observer. With regards to snow photochemistry, what is of interest includes the ability of the snowpack to store reactants, as influenced for example by its SSA, to transmit light, as dictated by its e -folding depth, and to release products to the atmosphere, as determined in part by its permeability. Typical values of relevant physical variables encountered in the photic zone of selected seasonal snow types in late winter are reported in Table 2 to illustrate the relationship

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

between snowpack type and physical properties. The actual ranges of values encountered are actually fairly wide. For example the SSA of surface snow often exceeds $800 \text{ cm}^2 \text{ g}^{-1}$ in the Alpine snowpack, due to the frequent presence of thick layers of fresh snow, whose density can remain below 0.1 g cm^{-3} for several days under windless conditions at sub-freezing temperatures. The heat conductivity of such layers can then be very low, down to $0.03 \text{ W m}^{-1} \text{ K}^{-1}$. Details on the range of values encountered in major snowpack types, and on the physical processes that determine them, are given in the following sections.

2.2 Density

Density is defined as the mass per unit volume, and because the mass of a unit volume of air is significantly less than the mass per unit volume of ice ($\rho_i = 0.917 \text{ g cm}^{-3}$ at 0°C), in practice the density of snow measures the amount of ice in a unit volume of snow. Because density is intuitive and does not require complex equipment to be measured, it has traditionally been the parameter used to describe the nature of snow, however it is now widely recognized that density alone cannot be used as a descriptor of other snow properties. Density is usually measured in the field or the laboratory by weighing a sample of known volume. The porosity is then defined from the snow density according to equation (1.4-1) above.

The density of natural snow can change enormously over time: from less than 10 kg m^{-3} for fresh cold dendritic snow to more than 600 kg m^{-3} for very hard windpacks and snow subjected to numerous melt freeze cycles. Physical processes involved in density changes include compaction, drifting, and dry and wet metamorphism. Deposited snow, with densities usually in the range 10 to 200 kg m^{-3} (Judson and Doesken, 2000) is subject to settling and compaction especially within the first week after deposition. Wind-drifted snow often forms thick layers of small rounded grains that sinter well and produce hard windpacks, with the windpacks density in the range 0.3 to 0.6 g m^{-3} increasing with the speed of the wind that formed them. High temperature gradient metamorphism can produce intense upward water vapor fluxes

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

that can fully compensate the effects of compaction, so that for example the density in the taiga snowpack is vertically fairly constant around 0.2 g cm^{-3} (Sturm and Benson, 1997; Taillandier et al., 2006). This intense dry metamorphism can lead to large mass losses in Arctic windpacks in the fall (Domine et al., 2002) and transform them into depth hoar, showing that significant density decreases can take place in snow layers. The appearance of liquid water usually leads to crystal rounding, compaction and density increases. Snow densities ranges for a variety of snow types are reported in Table 1.

Density values are required for almost all applications. Density parameterizations are therefore proposed in most snow physical models. Snow densification involves several processes and the rate of snow settling depends on the evolving snow microstructure, whose description remains a challenge and which is not closely related to density. In contrast to this complexity, snow densification is usually modelled by assuming that snow behaves like a viscous material. The material (snow) viscosity, η_s , then relates the stress σ to the strain rate $d\epsilon/dt$:

$$d\epsilon/dt = -(\sigma_s + \sigma_0)/\eta_s \quad (2.2-1)$$

Here σ_s is the applied snow stress i.e. in general the self-weight of the snow pack and the initial stress, σ_0 , is a possibility to include the effect of the rapid initial settling associated with destructive metamorphism of dendritic snow. The complex snow processes during settling are then expressed by mostly empirical relationships for viscosity and the initial stress. Simple models typically use $\sigma_0 = 0$ and parameterize the viscosity as a non-linear function of density and temperature only (e.g. Kojima, 1975). More advanced models have complicated parameterizations for both. We report here the formulations in SNTHERM (Jordan, 1991), CROCUS (Brun et al., 1989) and SNOWPACK (Lehning et al., 1999).

In SNTHERM, the initial settling is described independently of viscosity, which can be written as a function of temperature, T , (in $^{\circ}\text{C}$), ice volumetric content, θ_i , and water

volumetric content, θ_w :

$$\sigma_0/\eta_s = -2.778 \cdot 10^{-6} c_3 c_4 e^{0.04T},$$

$$c_3 = c_4 = 1, \quad \theta_i \leq 0.15 \text{ and } \theta_w = 0$$

$$c_3 = e^{-0.046(\theta_i - 0.15)}, \quad \theta_i > 0.15$$

$$c_4 = 2, \quad \theta_w > 0 \tag{2.2-2}$$

The snow viscosity is then parameterized as function of temperature and snow density, ρ :

$$\eta_s = \eta_0 e^{-c_5 T_c} e^{c_6 \rho},$$

$$\eta_0 = 3.6 \cdot 10^6 \text{ N s m}^{-2}$$

$$c_5 = 0.08 \text{ K}^{-1}$$

$$c_6 = 0.021 \text{ m}^3 \text{ kg}^{-1} \tag{2.2-3}$$

In CROCUS, a similar yet somewhat simpler formulation based on the settling law of Navarre (1975) is implemented. In the original implementation (Brun et al., 1989), snow viscosity is described as a function of temperature and snow density alone. That formulation suggested a factor for a microstructure correction, which has been implemented since (Coléou et al., 2005).

The original implementation of SNOWPACK used a visco-elastic, natural strain settling law (Bartelt and Lehning, 2002) with different viscosity laws for fresh and old snow (Lehning et al., 2002a). The viscosity formulation made use of a grain-bond description of snow microstructure and reduced snow settling to the deformation of ice bonds between the grains distinguishing between linear and non-linear yields of the ice matrix. Additional experiments and sensitivity studies have shown, however, that in natural settling elasticity and non-linear yield can be neglected and Eq. (2.2-1) is a good description of the settling behaviour. The current version of SNOWPACK (version 9.x) therefore retains the microstructure based development of a linear snow viscosity as described by Eqs. (33)–(35) in Lehning et al. (2002a). However, the empirical adjustment factor, F_{η} , is reformulated to include the liquid water content θ_w , the age of

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the snow layer, t_d (days), and variables representing crystal shapes and taking values between 0 and 1, namely dendricity, dd , and sphericity, sp , as defined in Brun et al. (1992):

$$F_{\eta} = \frac{0.4 + 9.5\theta_w}{\theta_i} + c_t \left(1 - \frac{t_d}{90}\right),$$

$$t_d \leq 90;$$

$$c_t = 2.1\sqrt{dd} + 0.7sp; \quad (2.2-4)$$

Here the time coefficient, c_t , is a function of the additional grain shape parameters dd and sp for all snow types except for melt-freeze crusts, where it is set constant to 1.7. The initial stress is formulated as a function of the ice surface tension, σ_i (0.11 N m^{-2}), the grain size, r_g and the dendricity, dd :

$$\sigma_0 = \frac{\sigma_i}{r_g} (1.5 + 2\sqrt{dd}). \quad (2.2-5)$$

The new SNOWPACK formulation calculates reliable settling curves for an Alpine seasonal snow cover. As shown in Fig. 4, the settling of individual snow layers tracked in the Alps at the Weissfluhjoch Versuchsfeld (Fierz and Lehning, 2001) can be described with good accuracy for the full seasonal development. It has also been investigated that errors in the settling curves such as in the 4th layer from the bottom arise from errors in the density estimation for fresh snow (Lehning et al., 2002b). Both a good density estimation and a good settling description are mandatory to allow a reliable snow mass balance at a point (Lehning et al., 1999) or for a full catchment (Lehning et al., 2006) based on snow depth measurements.

2.3 The specific surface area of snow

The specific surface area (SSA) of snow is a measure of the area of the air-snow interface. Its usual definition is the surface area of snow crystals accessible to gases

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

per unit mass (Legagneux et al., 2002), often expressed in $\text{cm}^2 \text{g}^{-1}$. Values measured in surface snow range from $19 \text{cm}^2 \text{g}^{-1}$ for melt-freeze crusts to $1558 \text{cm}^2/\text{g}$ for fresh dendritic snow (Domine et al., 2007).

Three methods are reported in the literature to measure reliably snow SSA: CH_4 adsorption at 77 K, stereology, and X-ray tomography. The first one, detailed in Legagneux et al. (2002) and Domine et al. (2007) has produced the largest number of measurements (345) for surface snow. Its principle is to measure the adsorption isotherm of CH_4 on snow using a volumetric method. A mathematical method (called BET) proposed by Brunauer et al. (1938) is then used to derive the SSA from the isotherm. The principle of gas adsorption to measure the SSA of porous solids dates back to the beginning of the century. However, obtaining reliable measurements for snow is not simple because this solid has a much lower SSA than other porous solids for which the method was intended, such as catalysts whose SSA can exceed $100 \text{m}^2 \text{g}^{-1}$. Using CH_4 rather than the more commonly used N_2 lowered the vapor pressure of the adsorbate at 77 K, and considerably improved measurement accuracy. The detection limit for this method is about $10 \text{cm}^2 \text{g}^{-1}$. One of its advantages is that it requires no sample preparation: it is sufficient to sample snow into a vacuum container and immerse it in liquid nitrogen, which has the added advantage of stopping metamorphism. However, it requires liquid nitrogen, a potential difficulty in polar campaigns. One SSA measurement with this method takes about 3 h.

Stereology is the study of 2-dimensional images to obtain information on 3-dimensional objects (Underwood, 1970) and is often used to study porous media such as snow (Narita, 1971; Perla et al., 1986). One method to obtain 2-dimensional images of snow is to harden a snow sample by filling it with a water-insoluble liquid that freezes at $T < 0^\circ\text{C}$, such as dimethyl-phthalate (Perla et al., 1986). The sample is then polished and macrographs are analyzed using appropriate methods (Davis et al., 1987) to derive SSA. The succession of all these preparation stages takes about 4 h for one SSA measurement.

The most extensive set of SSA measurements of surface snow using stereology is

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

that of Narita (1971). This paper is in Japanese with only an English abstract, so that details are not readily accessible. Narita studied snow and firn samples, and it appears that a few tens of samples were surface snow, the highest SSA value obtained being about $750 \text{ cm}^2 \text{ g}^{-1}$. A few other snow SSA values are available in other studies focused on firn and ice cores. From the data of Alley and Koci (1988) a value for an unspecified surface snow sample of $170 \text{ cm}^2 \text{ g}^{-1}$ can be inferred.

SSA values of surface snow obtained by both methods are in the same range, but no comparative studies have been made. The former method is probably more suited for surface snow, while the latter appears more suitable for firn and ice cores because of a lower limit of detection. From the data of Alley and Koci, SSA values for ice cores of about $2 \text{ cm}^2 \text{ g}^{-1}$ are calculated, and this is below the detection limit of CH_4 adsorption. On the other hand, fresh snow crystals have microstructure with radii of curvature of only a few μm , such as thin dendrites, that can be destroyed during all the stages of sample preparation for stereological studies: filling with a liquid induces extreme temperature gradients and warming of the sample that can be destructive for small structures. Polishing may also destroy these fragile structures. Moreover, it is not clear that features a few μm in size can be detected in automated processing of optical micrographs, so that this method may underestimate the SSA of high-SSA snow samples. Stereology, however, does not require liquid N_2 .

Recently, X-ray tomography has been used to obtain 3-D images of snow samples (Flin et al., 2003; Schneebeli and Sokratov, 2004), from which the SSA can be computed. This method requires a synchrotron (Flin et al., 2003) with the benefit of good spatial resolution and fast data acquisition, or a microtomograph (Schneebeli and Sokratov, 2004). Neither of these instruments can be realistically used in the field, and the data processing can be time-consuming. The interest of this method is to yield 3-D images of snow samples, but it is not worth the effort if only SSA is sought.

At present, it appears that CH_4 adsorption at 77 K is the method best suited to the measurement of the SSA of snow in the photic zone. However, this method is time consuming and its need for liquid nitrogen limits its use, so that a faster method readily

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

useable in the field is highly desirable.

Recently, Domine et al. (2006) have observed an excellent correlation between snow SSA measured by CH₄ adsorption and the snow reflectance in the near and short-wave infrared (NIR and SWIR). Wavelength tested were 1310, 1629, 1740 and 2260 nm.

The relationship was linear for the last 3 wavelengths. This is based on only 9 snow samples, however, and further work is needed before the potential of using NIR or SWIR reflectance to measure snow SSA can be established. In particular, it is possible that grain shape affect the reflectance (Neshyba et al., 2003; Kokhanovsky and Zege, 2004), so that tests using a wide range of snow crystal shapes are needed before the accuracy of this new method can be established. Matzl and Schneebeli (2006) also used NIR reflectance around 900 nm to measure snow SSA with a camera. The signal was calibrated with stereological measurements. This method is very rapid and quite promising. Its accuracy and in particular the dependence of the NIR signal on crystal shape needs to be studied further, however, especially since this is a unidirectional measurement. In any case, both these IR studies suggest that the accurate and rapid measurement of snow SSA in the field may soon be at hand.

Snow models do not currently treat SSA. As done for other variables such as heat conductivity (Sturm et al., 1997) the simplest method would be to correlate SSA with density. Figure 5, however, show that the correlation is poor. A significant improvement is obtained if the snow type is considered, and further improvement is observed if both the snow type and the snowpack type are considered. For example, Domine et al. (2007) show that for depth hoar in the taiga snowpack, the following relationship is found, with R²=0.66 :

$$SSA = -23.97 \ln(\rho) - 241.9 \quad (2.3-1)$$

with SSA in cm² g⁻¹ and ρ in g cm⁻³. Domine et al. (2007) also show that depth hoar in the tundra and Alpine snowpacks follow a similar relationship with different coefficients, yielding SSA values higher than in the taiga snowpack, for a given density.

Rather than attempt to link SSA to density, one can attempt to predict its rate of decrease after precipitation. Cabanes et al. (2002 and 2003) monitored the rate of

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

decrease of dry snow SSA in the Arctic and in the Alps and proposed an empirical equation of the form :

$$SSA = SSA_0 \exp(-\alpha t), \quad \text{with } \alpha \text{ (day}^{-1}\text{)} = 76.6 \exp(-1708/T(\text{K})) \quad (2.3-2)$$

This formulation is based on a limited data set, and predicts SSA only as a function of temperature, time, and initial SSA of the snow, SSA_0 (i.e. the SSA of the freshly fallen snow) Legagneux et al. (2003 and 2004) and Taillandier et al. (2007) have studied the rate of decrease of dry snow SSA, both in the field and during cold room experiments and observed that in all cases studied, a good empirical fit of SSA decay was possible with an equation of the form :

$$SSA(t) = B - A \ln(t + \Delta t) \quad (2.3-3)$$

Taillandier et al. (2007) found that two sets of values of B, A and Δt could be found for two different types of conditions and expressed as a function of SSA_0 and the time-averaged temperature of the snow T_m : one set was valid when the temperature gradient in the snowpack was low ($<9^\circ\text{C m}^{-1}$, i.e. ET conditions, as defined in Sommerfeld and LaChappelle, 1970) and the other set was valid under high temperature gradient ($>20^\circ\text{C m}^{-1}$, i.e. TG conditions, Sommerfeld and LaChappelle, 1970). Under ET conditions, the coefficients are:

$$A_{ET} = 0.0760SSA_0 - 1.76(T_m - 2.96) \quad ; \quad B_{ET} = 0.629SSA_0 - 15.0(T_m - 11.2);$$

$$\Delta t_{ET} = e^{\left[\frac{-0.371 SSA_0 - 15.0 (T_m - 11.2)}{0.0760 SSA_0 - 1.76 (T_m - 2.96)} \right]} \quad (2.3-4)$$

Under TG conditions, the coefficients are

$$A_{TG} = 0.0961SSA_0 - 3.44(T_m + 1.90) \quad ; \quad B_{TG} = 0.659SSA_0 - 27.2(T_m - 2.03);$$

$$\Delta t_{TG} = e^{\left[\frac{-0.341 SSA_0 - 27.2 (T_m - 2.03)}{0.0961 SSA_0 - 3.44 (T_m + 1.90)} \right]} \quad (2.3-5)$$

In the above equations, SSA is in $\text{cm}^2 \text{g}^{-1}$, T_m is in $^\circ\text{C}$ and Δt is in hours. These values reproduce well the experimental data of Taillandier et al. (2007), but the authors note

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

a number of limitations, such as the lack of data for dense snow ($\rho > 300 \text{ kg m}^{-3}$) so that predicting the SSA decay of for example Arctic windpacks is not possible. Another crucial lack of data is for wet snow, as Domine et al. (2007) report only a handful of measurements from which field estimations are uncertain. Data on snow SSA is therefore still limited, despite its importance in snow chemistry, and would definitely benefit from a new rapid measurement method based on IR reflectance (Domine et al., 2006; Matzl and Schneebeli, 2007).

2.4 Optical properties of snow

2.4.1 Light transmission

The transmission of light in snow plays important roles in atmospheric and snow chemistry as well as climate. The relationship to climate occurs largely through the albedo of snow, which is an important factor in the radiation balance (Grenfell et al., 1994). Recent work also shows that photochemistry in snowpacks plays an important role in the atmospheric chemistry in snow-covered regions (Honrath et al., 1999; Sumner and Shepson, 1999; Jones et al., 2000; Beine et al., 2002; Dibb et al., 2002). Light transmission in snow is also an important variable when considering biological activity in the subnivean environment, e.g. Kappen (1993) and Cockell et al. (2002).

When light interacts with snow grains, there are two important physical processes: scattering and absorption. We can consider the probability of either scattering or absorption per unit length traversed by a photon in the snowpack. We will refer to these scattering and absorption coefficients (rates per length) as K_{sca} and K_{abs} , and each has units of inverse length, typically cm^{-1} . The sum of scattering plus absorption is the extinction coefficient, $K_{\text{ext}} = K_{\text{sca}} + K_{\text{abs}}$. The ratio of scattering to total extinction, $\omega_0 = K_{\text{sca}} / K_{\text{ext}}$, is called the single scattering albedo. Typically, for visible wavelengths, the scattering coefficient is much larger than the absorption coefficient, leading to a single scattering albedo very close to unity. This dominance of scattering makes reflection the most probable occurrence for a photon that hits the snow and makes snow

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

white in appearance. The simplest optical models of snow use two-stream radiative transfer theory (Wiscombe and Warren, 1980; Warren, 1982; Beaglehole et al., 1998; Thomas and Stamnes, 2002) to solve the transmission of light in snow. If simple two-stream theory is used for snow, the scattering and absorption parameters have little physical meaning and are similar to fitting parameters. More advanced treatments take into account the angular asymmetry of light scattering by snow. Because snow grains are much larger than the wavelength of light, they typically forward scatter the light. This angular asymmetry can be included into advanced 2-stream theory calculations through the asymmetry factor, g , defined as the average of cosine of the scattering angle:

$$g = \frac{1}{2} \int_0^\pi x(\theta) \cos \theta \sin \theta d\theta \quad (2.4-1)$$

where $x(\theta)$ is the phase function describing the angular distribution of the singly scattered light, or more advanced theories, such as the delta-Eddington (Wiscombe, 1977) or the discrete ordinate transformation, DISORT (Stamnes et al., 1988), which is included in the Tropospheric Ultraviolet and Visible (TUV) model (Lee-Taylor and Madronich, 2002). When using these more advanced models, one can relate the scattering and absorption coefficients to snow physical properties. Specifically, the scattering coefficient is related to the snow specific surface area, and the absorption coefficient is related to the wavelength-dependent absorption of ice (Perovich and Giovoni, 1991) plus impurity absorption and grain shape. These correlations are discussed in detail below.

The relationship between light intensity and depth is complex in the near-surface of the snow. This complexity is caused by a combination of upwelling radiation escaping from the snowpack and direct radiation converting to diffuse radiation by scattering near the snow surface. This results in the light intensity being greater in the subsurface than at the surface (Fig. 6). Below this level in a deep uniform snowpack, the light intensity

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

decreases exponentially (e.g. Bohren and Barkstrom, 1974; Zege et al., 1991), as shown in Fig. 6:

$$I(z) \propto e^{-\alpha_0(\lambda)z} \quad (2.4-2)$$

In this equation, $I(z)$ is the light intensity at depth z , and $\alpha_0(\lambda)$ is the asymptotic flux extinction coefficient (AFEC). The inverse of α_0 is the depth over which the intensity of radiation decreases by a factor of e and is called the e -folding depth, $\varepsilon(\lambda)$,

$$\varepsilon(\lambda) = 1/\alpha_0(\lambda) \quad (2.4-3)$$

The transmission of radiation in snow has been measured in a number of studies (Liljquest, 1956; Grenfell and Maykut, 1977; Kuhn and Siogas, 1978; Beaglehole et al., 1998; King and Simpson, 2001). The general result is that the e -folding depth in the visible and near UV is between 5 and 25 cm for most snowpacks. The AFEC, α_0 , is directly observable by light transmission measurements and is affected by both light absorption and scattering.

In simple two-stream theory, one derives for AFEC:

$$\alpha_0(\lambda) = \sqrt{K^2(\lambda) + 2K(\lambda)S} \quad (2.4-4)$$

Here K and S are Kubelka-Munk absorption and scattering coefficients, respectively. They are related to K_{abs} and K_{sca} , respectively, but as mentioned above, they do not have direct physical meaning due to their disregard of the angular nature of scattering. K and S can be determined from experimental measurements of light and transmission by a snow layer using Kubelka-Munk theory (Kubelka and Munk, 1933; Kubelka, 1948). This formulation shows that both scattering and absorption are important to light transmission. In the limit of low absorption snow, K is small, and the K^2 term is small compared to the product of $K \cdot S$. Therefore, the dependence of light transmission is on $\sqrt{2KS}$. Many authors have generated radiation transfer theories for snow that can predict the AFEC (Bohren and Barkstrom, 1974; Wiscombe and Warren, 1980; Warren

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1982; Zege et al., 1991; Lee-Taylor and Madronich, 2002) including consideration of the angular asymmetry of scattering. In particular, Zege et al. (1991) show that :

$$\alpha_0 = \sqrt{3(1 - \omega_0)(1 - g)K_{\text{ext}}} \quad (2.4-5)$$

Because snow grains tend to forward scatter radiation ($g \sim 0.8$) (Warren 1982, Grenfell and Warren, 1999; Neshyba et al., 2003; Kokhanovsky and Zege, 2004) the AFEC is longer than predicted based by simple two-stream theory that assumes light scattering is equal into the forward and backward hemispheres.

The density scaling of snow optics is somewhat complex because the density affects both the amount of ice absorption and the number of ice-air surfaces encountered by light rays propagating in snow. Therefore, density affects both K and S absorption and scattering terms. It is common to create density-normalized “cross sections” to K and S ,

$$k_{\text{tot}}(\lambda) = K(\lambda) / \rho_{\text{snow}} \quad \text{and} \quad \sigma_{\text{scat}} = S / \rho_{\text{snow}} \quad (2.4-6)$$

Here, we have explicitly listed the wavelength dependence of the absorption, while the scattering cross section is relatively independent of wavelength. Substituting the cross sections for their analogs in Eq. (2.4-4) and in the weak absorption limit gives

$$\alpha_0(\lambda) = \rho_{\text{snow}} \sqrt{2k_{\text{tot}}(\lambda) \sigma_{\text{scat}}} \quad (2.4-7)$$

Most commonly, k_{tot} can be derived by measurements of impurities in snow and the inherent absorption of ice, which is particularly important in the near IR and thermal IR, where water vibrations of the ice lattice are strong absorbers.

Modeling studies have suggested that using measurements of the largest axis of a snow grain to determine the snow scattering results in scattering coefficient values that are too low (Grenfell et al., 1981). Grenfell and Warren (1999) and Neshyba et al. (2003) discuss different methods for representing ice crystals to calculate their scattering and absorption properties in clouds. They recommend the use of independent

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

spheres with the same volume-to-surface area ratio, V/S , as a nonspherical ice crystal for calculating optical parameters and conclude that the approximation is good for the range of sizes that were studied. The V/S ratio is related to the specific surface area (SSA) of snow for spherical approximate grains, thus the scattering coefficient of snow should be related to the SSA.

The SSA is the surface area per mass. Therefore, it follows:

$$\text{SSA} = \frac{S}{\rho_{\text{ice}}V} = \frac{3}{\rho_{\text{ice}}a_{\text{ef}}} \quad (2.4-8)$$

where $a_{\text{ef}}=3 V/S$ is the effective radius that coincides with the radius for monodisperse spheres.

The snow extinction coefficient is given by (Kokhanovsky and Zege, 2004):

$$K_{\text{ext}} = \frac{3c_g}{2a_{\text{ef}}} \quad (2.4-9)$$

where c_g is the volumetric concentration of grains in snow. This means that

$$K_{\text{ext}} = \frac{\rho}{2} \text{SSA} = \frac{3\rho}{2\rho_{\text{ice}}a_{\text{ef}}} \quad (2.4-10)$$

where $\rho=c_g\rho_{\text{ice}}$ is the snow density. This expression is also valid for the scattering coefficient because $K_{\text{sca}}\approx K_{\text{ext}}$ for snow in the visible.

Therefore, we can see that the scattering coefficient of snow is proportional to the density of the snow divided by the ice density (the volumetric concentration of ice in snow) and inversely related to the equivalent-sphere grain radius a_{ef} . For aged snow, we typically have $\rho\sim 0.3\text{ g cm}^{-3}$, and $a_{\text{ef}}\sim 0.5\text{ mm}$, so that $K_{\text{ext}}\sim K_{\text{sca}}\sim 1\text{ mm}^{-1}$. This result implies that light is scattered roughly every millimeter in the snow. The direct solar beam is scattered into diffuse radiation on this length scale, while the diffuse component decays much more slowly (5–25 cm length scale).

The key findings of this section are that radiative transfer in snow is relatively well understood. The asymptotic flux extinction coefficient (AFEC) scales as the square

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

root of the product of the snow scattering and absorption coefficients. The scattering coefficient of snow is well modeled by a collection of equivalent spheres that have the same SSA as the snow. The scattering coefficient is thus directly related to SSA and inversely related to the equivalent-spherical radius. The equivalent sphere radius of snow is significantly smaller than the “snow grain radius” because it takes into account scattering from small features within the snow grain.

2.4.2 Light reflection

The photochemistry of snow requires information on the total amount of light absorbed in the snow at each wavelength. Therefore, it is of advantage to study the absorbance defined as

$$a = 1 - r_d \quad (2.4-11)$$

where r_d is the diffuse reflectance under directional illumination (e.g. clear sky). For non-absorbing snow, by definition: $r_d=1$ and $a=0$. In reality snow absorbs some amount of solar energy at each wavelength and therefore, $a \neq 0$. The value of r_d can then be calculated using the following simple approximation (Zege et al., 1991; Kokhanovsky, 2006):

$$r_d(\lambda) = \exp(-\rho(\lambda)u(\vartheta_0)) \quad (2.4-12)$$

Here ϑ_0 is the solar zenith angle, λ is the wavelength, $\rho=4\sqrt{(1-\omega_0)/3(1-\omega_0g)}$ and the function $u(\vartheta_0)$ can be parameterized as discussed by Kokhanovsky and Zege (2004):

$$u(\vartheta_0) = 3(1 + 2 \cos \vartheta_0) / 7 \quad (2.4-13)$$

These last three equations indicate that absorption of light by snow is larger for a higher sun position ($\vartheta_0 \rightarrow 0$). The physical basis for this effect is that it is more difficult for photons to escape from the snow if they are injected along the normal to the layer because scattering occurs predominantly in the forward direction.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Under diffuse illumination such as complete cloud cover, snow reflectance r_s is calculated by integrating the diffuse reflectance r_d over all incidence angles (Kokhanovsky, 2006):

$$r_s = \int_0^{\pi/2} r_d(\vartheta_0) \sin 2\vartheta_0 d\vartheta_0 \quad (2.4-14)$$

5 The value of r_s is called the diffuse reflectance under diffuse illumination conditions or, in short, spherical albedo. It follows in the same approximation (Kokhanovsky and Zege, 2004):

$$r_s(\lambda) = \exp(-\rho(\lambda)) \quad (2.4-15)$$

10 Kokhanovsky (2006) has proposed to use the following relationship between the parameter $\rho(\lambda)$ and the grain diameter d for the case of pure snow:

$$\rho(\lambda) = 8\sqrt{\pi\chi(\lambda)d/\lambda} \quad (2.4-16)$$

Here $\chi(\lambda)$ is the imaginary part of the refractive index of ice, as reported for example in Wiscombe and Warren (1980). Then it follows:

$$r_s(\lambda) = \exp(-8\sqrt{\pi\chi(\lambda)d/\lambda}) \quad (2.4-17)$$

15 This simple equation has been used to calculate the spherical albedo of pure snow for different grain diameter, treating snow as ice spheres (Fig. 7). Similarly, we have for the diffuse reflectance, r_d , (see Eqs. 2.4-12, 2.4-13):

$$r_d(\lambda) = \exp(-24\sqrt{\pi\chi(\lambda)d/\lambda}(1 + 2\cos\vartheta_0)/7) \quad (2.4-18)$$

20 In the presence of absorbing impurities this equation must be modified. In particular, in the case of absorption of light by impurities that show much stronger absorption than ice, the value of $\chi(\lambda)$ in Eqs. (2.4-17) and (2.4-18) must be substituted by the

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

function proportional to the product of the imaginary part of the refractive index of an absorber $\chi_a(\lambda)$ and its volumetric concentration, c_s , (Kokhanovsky and Zege, 2004):

$$\chi(\lambda) = A\chi_a(\lambda)c_s \quad (2.4-19)$$

where the value of A depends on the type of impurities. A can be determined using Eqs. (2.4-17) and (2.4-18) and adequate measurements. It can also be calculated using various theoretical models (Wiscombe and Warren, 1980).

In particular, the following model for the calculation of the parameter ρ in Eq. (2.4-15) can be used. It is supposed in this model that the extinction is only due to scattering of light by snow grains and by absorption caused by impurities (e.g. soot) treated as small Rayleigh scatterers. Then the absorption coefficient is given by (Kokhanovsky, 2004):

$$K_{\text{abs}} = N_s f(n) \gamma(\lambda) V_s \quad (2.4-20)$$

where $N_s = c_s / V_s$ is the number concentration of absorbing particles, V_s is their average volume, c_s is their average volumetric concentration, $\gamma(\lambda) = 4\pi\chi_s(\lambda) / \lambda$, $\chi_s(\lambda)$ is the imaginary part of the refractive index of absorbing particles and $f(n) = 9n / (n^2 + 2)^2$, where n is the refractive index of absorbing particles. On the other hand, it follows for the extinction coefficient of large snow grains:

$$K_{\text{ext}} = \frac{3c_g}{d_{\text{ef}}} \quad (2.4-21)$$

where c_g is the volume fraction of snow occupied by snow grains and d_{ef} is the diameter of ice spheres with the same SSA as the snow. Therefore, one derives finally:

$$1 - \omega_0 \equiv \frac{K_{\text{abs}}}{K_{\text{ext}}} = \frac{f(n) \gamma(\lambda) c_s d_{\text{ef}}}{3c_g} \quad (2.4-22)$$

and, therefore,

$$\rho = \frac{4}{3} \sqrt{\frac{f(n) \gamma(\lambda) c_s d_{\text{ef}}}{c_g(1 - g)}} \quad (2.4-23)$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

where we accounted for the fact that $\omega_0 \rightarrow 1$ in the spectral region under study.

Using this result, one derives:

$$r_s = \exp \left\{ -\frac{4}{3} \sqrt{\frac{f(n) \gamma(\lambda) c_s d_{ef}}{c_g(1-g)}} \right\} \quad (2.4-24)$$

and

$$r_d = \exp \left\{ -\frac{4u(\vartheta_0)}{3} \sqrt{\frac{f(n) \gamma(\lambda) c_s d_{ef}}{c_g(1-g)}} \right\} \quad (2.4-25)$$

The dependence of r_s on the concentration of impurities c_s is presented in Fig. 8, showing that a few ppm by volume of soot can have a dramatic impact on snow albedo.

2.4.3 Photochemistry in snow

Radiation, particularly UV, drives photochemistry both above and inside the snow. The radiation above the snow is enhanced by the high albedo of the snow. Many studies have documented this effect (Madronich, 1987; Meier et al., 1997; Simpson et al., 2002b). Additionally, the radiation in the snow is enhanced. The snowpack enhancement comes from the conversion of direct to diffuse radiation (Madronich, 1987; Simpson et al., 2002a; Lee-Taylor and Madronich, 2002). Laboratory studies have demonstrated that photolysis rate coefficients modeled in the snow pack agree with those measured by chemical actinometry (Phillips and Simpson, 2005). A number of studies have used chemical actinometry in the snow pack to measure photolysis rates directly (Qiu et al., 2002). Laboratory studies have measured quantum yields for important photochemical reactions in the snow; e.g. Nitrate photolysis (Honrath et al., 1999, Jones et al., 2000; Beine et al., 2002; Simpson et al., 2002a), and hydrogen peroxide photolysis (Chu and Anastasio, 2005). More details on photochemistry in the snow pack can be found in Grannas et al. (2007).

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.5 Thermal conductivity

Because of snow metamorphism, the thermal properties of snow are highly variable and can change dramatically in time. Traditionally, estimates of snow thermal properties are based on readily-measurable characteristics such as density and grain size, though it is well known that the geometric forms of the microstructure are a first-order effect. There are still no quantitative analysis that can predict thermal properties based on microstructure (Arons and Colbeck, 1995).

The thermal conductivity of snow controls the net heat transfer in the snow when there is no interstitial air flow. In one dimension, the heat flux q at a point is given by the Fourier equation:

$$q = -k_{\text{eff}} \frac{dT}{dz} \quad (2.5-1)$$

where k_{eff} is the effective thermal conductivity, T is temperature, and z is the spatial coordinate along the direction of flow. Heat flow through snow results from the combined effect of several physical processes, as shown in Fig. 9.

Heat conduction takes place through the network of interconnected snow crystals and through air in the pore space. In addition, there almost always exist a temperature gradient in dry snow, which generates water vapor sublimation from warmer grains, diffusion through the pore space and condensation on colder grains (Colbeck, 1983) leading to heat transport through latent heat exchanges. The effects of these processes can be measured using a heated needle probe (e.g. Sturm and Johnson 1992; Sturm et al., 1997; Mellor, 1977; Lange, 1985) where the rate of dissipation of a given heat pulse is analyzed to derive the “effective” thermal conductivity k_{eff} . For this type of measurement, the snow sample is maintained enclosed in a fairly air-tight container to avoid heat transport due to air flow.

Air flow in snow can be caused by wind-driven variations in surface pressure over a rough snow surface (Albert et al., 2002) or by thermal convection caused by a high temperature gradient in very permeable snow such as depth hoar (Sturm and Johnson,

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1991), but this heat transport process is discussed in a subsequent section.

The variable k_{eff} has contributions from several physical processes, and the relative contributions of each one of these processes is expected to depend on the geometric aspects of the crystal structure and on the snow temperature. We therefore expect to observe a large scatter in k_{eff} values, and for dry snow, values between 0.021 and $0.65 \text{ W m}^{-1} \text{ K}^{-1}$ have indeed been measured (Sturm et al., 1997, and references therein). To attempt to predict k_{eff} values, correlations with snow density have been explored. Sturm et al. (1997) and Jordan et al. (2007) do report a density- k_{eff} correlation, although a significant scatter exists, that could be reduced if snow type were taken into account. Reported correlations show k_{eff} values ranging from approximately $0.07 \text{ W m}^{-1} \text{ K}^{-1}$ for a density of 0.1 g cm^{-3} to $0.7 \text{ W m}^{-1} \text{ K}^{-1}$ for a density of 0.5 g cm^{-3} (Jordan et al., 2007). Measured k_{eff} values of windpacked surface snow on polar ice sheets is between $0.15\text{--}0.20 \text{ W m}^{-1} \text{ K}^{-1}$ for a density near 0.35 g cm^{-3} , while aged near-surface firn that has not experienced melt on the East Antarctic ice sheet shows k_{eff} values ranging from 0.2 to $0.4 \text{ W m}^{-1} \text{ K}^{-1}$ for densities between 0.35 to 0.45 g cm^{-3} (Courville et al., 2007). Despite the large scatter in the density correlation, Sturm et al. (1997) proposed the following equations:

$$\begin{aligned} k_{\text{eff}} &= 0.138 + 1.01\rho + 3.2233\rho^2, \quad \{0.156 \leq \rho = 0.6\} \\ k_{\text{eff}} &= 0.023 + 0.234\rho, \quad \{\rho < 0.156\} \end{aligned} \quad (2.5-2)$$

that have been extensively used in various calculations and model parameterizations.

2.6 Permeability of snow and firn

The air permeability of snow is the property of snow that controls the ease with which a fluid, typically air or water, can move through the snow. Permeability K_p is the proportionality factor between the pressure gradient dP/dz and the interstitial air flow velocity

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

V_{air} :

$$V_{\text{air}} = -\frac{K_P dP}{\mu dz} \quad (2.6-1)$$

where μ is the air viscosity. Equation (2.6-1) is known as Darcy's law. The permeability is sensitive to the nature of the interconnected pore space, and it depends both upon the crystal structure and the snow layering. As in all physical properties of snow, permeability will change as the snow undergoes metamorphism. The permeability of snow is typically measured by drawing air through a snow sample and measuring the pressure drop and air flow rate through the sample, then calculating the permeability using Darcy's law (e.g. Albert et al., 2000). For seasonal snow, permeability ranges over two orders of magnitude depending on the snow type, density, grain size, and degree of metamorphism. The most permeable snow is depth hoar in seasonal snow, with permeability in the range from $100\text{--}600 \times 10^{-10} \text{ m}^2$. Freshly fallen snow typically has permeability in the range from 30 to $70 \times 10^{-10} \text{ m}^2$, while old coarse seasonal snow typically exists in the range from 8 to $50 \times 10^{-10} \text{ m}^2$ (Jordan et al., 1999). Ice layers and crusts formed by melt in seasonal snow, and subsequently buried by later snowfalls, have permeability ranging from 1 to $19 \times 10^{-10} \text{ m}^2$ depending on the degree of refreezing (Albert et al., 2000; Albert and Perron, 2000). Wind-packed snow on polar ice sheets typically falls in the range 5 to $30 \times 10^{-10} \text{ m}^2$ (Albert and Shultz, 2004). Coarse-grained, aged firn in very cold low accumulation sites like the East Antarctic plateau has high permeability in the range of $100\text{--}300 \times 10^{-10} \text{ m}^2$ (Albert et al., 2004). Natural stratification of snow can yield buried layers that are more permeable than surface layers, causing flow channelling through the buried high-permeability layers (Albert et al., 2000). Luciano and Albert (2002) reported measurements of anisotropy in snow permeability and concluded that differences in permeability between layers from different storms caused larger variation than directionality differences in a single layer.

Since snow density and grain size are the most commonly measured snow variables, several attempts have been made to relate snow permeability to density and grain size.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

One widely used equation is that of Shimizu (1970):

$$K_p = 0.077e^{(-0.0078\rho)}D^2 \quad (2.6-2)$$

with D the grain diameter in m and ρ the snow density in kg m^{-3} . As noted by Jordan et al. (1999), Shimizu's formula was obtained for small rounded grains, and can be in error by an order of magnitude for calculations based on grain size and density for other snow types. Jordan et al. (1999) also note that progress in predicting snow permeability may come from the use of the snow specific surface area rather than grain size, as SSA is a clearly defined variable, even though it is rarely measured in the field.

2.7 Gas diffusivity of snow

The gas diffusivity is a property of snow that indicates the ease with which gases can move through the interstitial pore space. The diffusivity, D_g , is the proportionality factor between the gas concentration gradient dC/dz and the interstitial gas flux F_{gas} :

$$F_{\text{gas}} = -\frac{D_g}{\phi} \frac{dC}{dz} \quad (2.7-1)$$

where ϕ is the snow porosity. Gas diffusivity will vary according to chemical species, as it depends on the size of the molecules relative to the size of an interstitial pathway. For reactive chemical species, sometimes the diffusivity is posed as an "effective" parameter that includes not only the obstruction aspects of diffusion in the pore space around snow grains, but also adsorption/desorption processes that occur on the grain surface. Thus, when using published values of gas diffusivity in snow, one must be careful to note whether or not the definition in the publication based the values on only the obstruction effect or whether they also include the adsorption/desorption reactive impacts. For this discussion, we define the diffusivity to be the behavior of the gas as though it were inert. Adsorption and desorption are then separate source-sink type processes.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Pioneering investigations of gas diffusivity of firn was done by Schwander et al. (1988) and more recently by Favre et al. (2000), whose experiments based on elution peaks of an advected tracer gas in a moving carrier gas led to the conjecture that snow microstructure and layering are inconsequential to gas diffusion. Recently, through experiments involving diffusion alone, in-situ measurements by Albert et al. (2007)¹ and on firn cores by Courville et al. (2007) have shown that the gas diffusivity of snow does depend on the structure of snow and firn. The measurement technique for in-situ measurements is described in Albert and Shultz (2002), which was adapted from a technique described by McIntyre and Philip (1964) for soils. Measurements of the diffusion of SF₆, an inert gas, through partially decomposed fresh dendritic seasonal snow has typical values close to 0.025 cm² s⁻¹ (Albert, unpublished data), while the SF₆ gas diffusivity of wind-packed snow typically ranges between 0.02 to 0.06 cm² s⁻¹ (Albert and Shultz, 2002). The SF₆ gas diffusivity of aged, coarse, near-surface polar snow in cold low accumulation sites in East Antarctica depends on the accumulation rate and ranges from 0.04 to 0.08 cm² s⁻¹ (Courville et al., 2007). For comparison, the diffusivity of SF₆ in air between 0 to -10°C is 0.08 cm² s⁻¹. Thus, depending on the nature of the snow, surface snow can either be little barrier to the transport of gas, or else slow down gas transport by as much as a factor of four compared to diffusion through air alone.

2.8 Effect of air advection in snow on the transport of heat and gases

Heat and chemical species can move through the snow by diffusion, a relatively slow, gradient-driven process, or by advection, a faster process by which heat and mass are transported by interstitial air flow (“ventilation”) in the pore space between the snow grains. In natural snow, diffusion is ubiquitous because the snow is almost never in

¹Albert, M. R., Neumann, T. A., Rick, U., and Steig, E.: Impact of post-depositional processes on interstitial transport with implications for diffusion modeling for ice core interpretation, J. Geophys. Res., submitted, 2007

perfect thermal or chemical equilibrium. Thus diffusion must always be considered. Advection, a faster process whereby chemical species in the firn air are transported by interstitial air flow, can occur both naturally and when interstitial air in the snow is sampled for chemical analysis (Albert et al., 2002). Wind-driven interstitial air flow within snow occurs mostly from pressure disturbances of the wind flowing over surface roughness. Thermal convection in extremely permeable snow subject to strong temperature gradients has been observed by Sturm and Johnson (1991) and resulted in easily detectable heat transport. Albert et al. (2004) found that extensively recrystallized firn formed under extremely cold conditions on the East Antarctic plateau has characteristics that would make it prone to thermal convection. However, buoyancy-induced convection is rarely found in other types of snow. Experimental studies performed under different conditions failed to detect convection (Brun and Touvier, 1987), and we therefore subsequently treat only wind-driven and sampling-induced advection.

Wind-driven ventilation can advect chemical species, and it has been shown to impact the concentrations of species contained in aerosols (Cunningham and Waddington, 1993; Harder et al., 2000, Domine et al., 2004), of species soluble in snow crystals (McConnell et al., 1998; Waddington et al., 1996), water vapor transport (Albert, 2002), and isotopic composition (Neumann and Waddington, 2004). In addition to advection caused naturally by the wind, advection also occurs when firn air is sampled for analysis. Because the technology of measuring trace gases currently uses through-flow chemical sampling devices, the sampling procedure itself induces ventilation and mixing of the air sampled. In the top meter or so of snow, this introduces ambient air dilution to the natural firn air composition (Albert et al., 2004). There have been no field experiments reported to date where the firn air gas sample concentrations reported are due purely to diffusion alone. Firn air sampling procedures for CO₂, methane, and other climate indicators that are sampled very deep in the firn typically call for withdrawal of discrete samples of gas in order to minimize the disturbance of the natural firn air concentration profile.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.9 Transport of species that interact with the snow surface

Diffusion and advection are not the only processes that affect trace gas transport through porous snow, because molecules hit ice surfaces while traveling through porous snow and those with sufficient affinity with these surfaces may become accommodated on the ice for various lengths of time. This “sticking” will therefore slow down gas transport through snow. The residence time of the molecule on snow crystal surfaces affects the partitioning of molecules between interstitial air and ice surfaces, and this can be characterized by an adsorption equilibrium, either a surface Henry’s law coefficient, H (in $\text{Pa m}^2 \text{ mol}^{-1}$) or an interfacial partitioning coefficient K_{ia} (in m) (Roth et al., 2004). Models that couple transport, adsorption and reaction in porous media have been derived from the study of reactions in porous catalysts. Such models are used for example to study transport processes in soil, and have also been applied to analyze reactions and uptake processes of trace gases on porous ice films in laboratory experiments, which lead to some controversy. (Hanson and Ravishankara, 1993; Keyser et al., 1993; Keyser et al., 1991; Leu et al., 1997).

What needs to be modeled is simply the fact that the diffusion of trace gases through the snow is hampered by resistance of the porosity and the finite residence time of the trace gases when adsorbed on the ice surface, and this requires the knowledge of the pore size and of SSA. Modeling can be achieved by numerical solution of the one-dimensional diffusion equation, with appropriate terms for the flux matching on the ice surface, as done by Hutterli et al. (1999) who modeled the seasonal transport of formaldehyde in snow by assuming one-dimensional diffusion for the gas phase transport.

A semi empirical approach was suggested by Herbert et al. (2006), which allows to assess the impact of the adsorption on the diffusion. In a first step, the theoretical diffusion coefficient D_{theo} in a porous medium was calculated from the tortuosity τ , the snow porosity ϕ and the diffusivity in the undisturbed gas phase D_{air} by $D_{\text{theo}} = D_{\text{air}} \phi \tau$. The additional resistance to the diffusive transport by the adsorption of the molecules

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

is described by an effective diffusion constant $D_{\text{eff}} = fD_{\text{theo}}$ with

$$f = \frac{1}{1 + r_{sp}K_{ia}SSA} \quad (2.9-1)$$

Parameters are the specific surface area (SSA), the solid ice to air volume ratio r_{sp} and the gas-solid partitioning coefficient K_{ia} (Roth et al., 2004). In Figure 10, we illustrate the impact of the value of K_{ia} on diffusion, by plotting f as function of K_{ia} . For K_{ia} smaller than 10^{-4} m (for example aliphatic hydrocarbons lighter than decane at $T = -6.8^\circ\text{C}$, Roth et al., 2004), there is almost no influence of the adsorption on the diffusion. For stronger adsorbing species, with $K_{ia} > 10^{-3}$ m (e.g. phenanthrene, Domine et al., 2007²), the impact of the adsorption can be critical.

3 Physical properties of snow, as determined from remote sensing

3.1 General remarks

Remote sensing methods (e.g. based on measurements performed by optical and microwave instruments onboard satellite platforms) cannot compete with in situ measurements of snow characteristics in terms of accuracy. For instance, a researcher on the field can use a microscope to determine the average grain size. This requires virtually no additional assumptions. On the other hand, a number a priori assumptions are involved in the interpretation of satellite signals, such as the shape of grains, the radiative transfer and atmospheric correction model, the surface roughness, the existence of sub-pixel bare soil or forest, clouds, etc. The assumptions used can be true or not. Therefore, in principle, highly biased results can be obtained. In particular, advanced cloud screening algorithms (e.g., using gaseous absorption features, Rozanov and Kokhanovsky, 2004) must be used for snow remote sensing applications.

²Dominé, F., Cincinelli, A., Bonnaud, E., Martellini, T., and Picaud, S.: Adsorption of phenanthrene on natural snow, Environ. Sci. Technol., submitted, 2007.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

However, while a researcher on the field can make reliable observations, their spatial representativity can be limited, while on the contrary satellite methods allow the monitoring of very large areas. Moreover, very remote areas such as polar oceans and ice caps are realistically accessible only to satellite investigations. Therefore, further developments of satellite techniques to monitor snow and ice from space are highly desirable.

3.2 Optical remote sensing

The reflection function of snow-covered areas in the visible, as observed from satellites, is highly sensitive to the fraction of the ground covered by snow and to snow pollution. The dependence on the grain size is less pronounced in the visible (Fig. 7). This is due to the fact that the grain sizes are quite large and their scattering characteristics are hardly influenced by the size of particles outside of the diffraction cone. However, the influence on the shape of particles remains (Grenfell and Warren, 1999; Neshyba et al., 2003; Zege and Kokhanovsky, 2004). A great simplification arises from the fact that snow can be considered as a semi-infinite scattering medium in the optical range. As a matter of fact, a snow layer of just 5 cm thick is virtually a semi-infinite medium in the visible and the near IR. Yet another simplification is due to the fact that the standard radiative transfer equation can be used for the interpretation of optical measurements for densely packed snow grains (Bohren and Beschta, 1979; Kokhanovsky and Zege, 2004). This is due to the fact that grains are quite large as compared to the wavelengths in the visible and near IR. Therefore, coherent scattering effects are of no importance. This assumption, however, does not hold for microwaves. Then grains are on the order of the wavelength and the dense media theories must be used instead of the standard radiative transfer equation. In particular, the spatial correlation of scatterers and its influences on scattering characteristics must be fully accounted for (Tsang et al., 2000).

Yet another complication is due to the fact that snow properties are not constant in the vertical direction. Therefore, the retrieval of the grain size using near IR measure-

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ments, where the snow reflection function is sensitive to the size of grains, and also the assumption of a vertically homogeneous layer will give different answers depending on the wavelength used. Light of smaller wavelength will penetrate to deeper layers, because generally ice absorption by ice grains is smaller for smaller wavelengths. This means that smaller wavelengths bring information not only from the upper snow layer but also from much deeper regions. This is generally not the case for larger wavelengths. Sizes retrieved for different wavelengths can be used to estimate the vertical inhomogeneity of a snow layer (Li et al., 2001). Future methods of snow grain size retrieval must rely on the measurements for a number of wavelengths (e.g., with a high spectral resolution). This will enable the determination of the grain size profile from correspondent fitting routines.

The retrieval of the grain size is based on the fact that the snow grain single scattering albedo, ω_w , which can be determined from the reflection function measurements and theoretical model based on the radiative transfer theory (see, e.g., Mishchenko et al., 1999), is highly dependent on the size of particles. Indeed larger sizes imply larger absorption of light by grains, which can be detected from measurements of the reflection function. For the retrieval of ω_w , one must assume the phase function of snow. In the past, Mie theory (Mie, 1908) has been used for this purpose (Nolin and Dozier, 1993; 2000). Nowadays, various nonspherical models are used (Mishchenko et al., 1999; Kokhanovsky and Zege, 2004; Xie et al., 2006). We underline that equivalent spherical grain models can perform well in the case of snow albedo measurements. However, satellite instruments measure the reflection function for a given incidence and observation directions. Therefore, they are based on the measurements of the directional properties of a surface, which differ for nonspherical and spherical scattering models (Mishchenko et al., 1999). The application of spherical models leads to the unphysical dependence of the grain size on the illumination or observation geometry.

The retrieved grain size can be used to estimate the snow reflection function in the visible. The difference of the measured reflection function from the correspondent theoretical model can be attributed to the pollution. This enables the determination of

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the concentration of pollutants (Tanikawa et al., 2002) from space-borne instrumentation assuming a type of a pollutant (soot, dust, etc.). This is of importance because dirty snow reduces the albedo of the planet and contributes to warming of the Earth-atmosphere system (Hansen and Nazarenko, 2004).

5 It must be noted that the retrieval of snow grain size and concentration of pollutants must rely on the atmospherically corrected radiances. In particular, correspondent MODerate Resolution Spectrometer data (MODIS) atmospherically corrected satellite products can be used (Vermote and Vermeulen, 1999; Hall et al., 2001; Tedesco and Kokhanovsky, 2007). The snow cover can be assessed directly from the snow mask algorithm, e.g., as described by Hall et al. (2001) for MODIS. MODIS enables the determination of snow cover with the resolution of 250 m×250 m. Remotely sensed snow parameters using the visible and near IR can be used in the retrieval of other snow parameters (e.g., the snow depth) from microwave measurements. Future algorithms must address the retrieval of snow properties using broad spectral ranges from the UV to microwave region.

15 The current capabilities of snow retrieval algorithms are demonstrated by works of Bourdelles and Fily (1993), Fily et al. (1997), Aoki et al. (2000), Nolin and Dozier (1993, 2000), and Painter et al. (2003). In particular, Painter et al. (2003) has found a good correlation of the retrieved and measured on the ground grain size. Although we would like to underline that the grain size measured by an orbiting instrument corresponds to the ratio of the volume of a grain to its surface area. This information is not readily available from 2-D image processing in field studies, and SSA measurements would be preferable.

3.3 Microwave remote sensing

25 Microwave sensors usually operate at a single or at several frequencies in the range 1 GHz–100 GHz. This range corresponds to wavelengths from a few millimeters (a bit larger than snow grains diameter) up to 30 cm (the depth of an Arctic snowpack or of an Alpine snow layer). Passive microwave sensors (called radiometers) measure the

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

radiation naturally emitted by the surface. Their spatial resolution is coarse, typically a few tens of kilometers. Active sensors illuminate the surface and measure the echo. Active instruments include scatterometers (imagers with resolution of few kilometers), synthetic aperture radars (SAR, imagers with decametric resolution) and altimeters (nadir looking radar).

Microwave remote sensing has the advantages to observe the surface even in cloudy conditions, to be weakly sensitive to the water vapor or aerosol thickness (except at some frequency bands) and not to rely on solar illumination. These advantages are crucial for applications requiring high temporal resolution such as the daily monitoring of the snow cover, or studies during the polar winter. Furthermore, many snow physical variables can be retrieved using microwave remote sensing, because these methods cover a large range of frequencies and use a wide diversity of measurement methods. Initially, methods were developed to derive a single variable from the data. This was possible for the following variables:

- The presence of snow on the ground.
- The snow water equivalent (SWE) on the ground.
- The net accumulation. This is inherently the same variable as the SWE, but applied to ice caps and integrated over an arbitrary period, not necessarily a season.
- The presence and amount of liquid water in the snowpack.

At present, more sophisticated algorithms are being written to retrieve simultaneously several interdependent variables, that include the former ones as well as:

- The snowpack temperature profile.
- The grain size.
- The layering of the snowpack.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- The surface roughness.

Before addressing the specific retrieval methods of each variable, we briefly recall in the next section the bases of the interactions between microwaves and snow. We also review the forward models developed to predict microwave observations from snow characteristics.

3.3.1 Background: interactions between snow and microwaves

Interactions between the snow and microwaves differ from those with visible/infrared radiation by two main features:

- Microwaves penetrate through dry snow. The e-folding depth depends on the absorption within the ice and scattering by the snow grains, and also on reflection at the boundaries between snow layers. Note that unlike in the optical range, the microwave e-folding depth is related to the direct rather than the diffused beam. Absorption mainly increases with density and, contrary to the optical range, scattering increases with grain size and frequency. Attenuation depends significantly on these variables: typical one-way e-folding depth is as large as hundreds of meters at 1 GHz (e.g. frequency of the SMOS radiometer and ALOS radar), about 10 m at 5 GHz (e.g. ENVISAT ASAR radar and ERS altimeter), 2 or 3 m at 20 GHz (e.g. SSM/I radiometer), 0.5 m at 37 GHz (SSM/I). In contrast to optical measurements, these large e-folding depths allow the retrieval of information on large snow thicknesses.
- The presence of liquid water in the snow radically increases the absorption of microwaves and as a consequence largely reduces the e-folding depth (to 1 cm at 20 GHz). Even a few percent of liquid water content efficiently absorbs microwaves because snow grains coated with a thin layer of water are seen, by microwaves, almost as if they were composed of water only. In practice, this difference between wet and dry snow implies that 1) detecting wet from dry snow is

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

easy, but 2) variable-retrieval algorithms work for dry snow only.

3.3.2 Passive measurements

Radiometers measure the thermal emission of the Earth expressed by the “brightness temperature”. For an opaque medium such as soil or rock, the brightness temperature is given, to a first approximation and neglecting the (weak) perturbing atmospheric term, by:

$$T_b = \varepsilon_{mw} \times T_s \quad (3.3-1)$$

where ε_{mw} is the emissivity and T_s the surface temperature.

For a transparent medium such as the snowpack, the relationship is more complex, as the signal received originates from a thickness related to the e-folding depth. The soil can therefore affect the measurements. Usually, radiometers acquire T_b at one or several of various possible frequencies (e.g. 19, 22, 37 and 85 GHz for SSM/I) and two polarisations: vertical and horizontal. Using several frequencies has the potential of yielding the vertical temperature profile in the snowpack.

For a rigorous calculation of T_b , radiative transfer equations need to be solved in the snowpack to account for the thermal emission (depending on the temperature vertical profile), scattering (depending on grain size and frequency), absorption (depending mostly on density and frequency), the reflective layers (depending on the density and grain size variations) and the underlying soil. Thermal emission and scattering by the atmosphere also need to be accounted for at some frequencies (e.g. 22 GHz is highly sensitive to water vapor, 37 GHz and higher are sensitive to intense liquid precipitation). Earlier models were based on the strong fluctuation theory (Stogryn 1986, Surdyk et Fily, 1993 and 1994). The HUT (Helsinki University of Technology) snow emission model is a semi-empirical model (Pulliainen et al., 1999) that treats homogeneous snowpack and includes vegetation effects, that are especially important in boreal regions. MEMLS (Microwave Emission Model of Layered Snowpacks, Wiesmann and Mätzler, 1999) or the model based on Dense Media Radiative Transfer (Macelloni et al.,

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2001) are more complex. They solve the radiative transfer for a multi-layered snowpack and account for reflection, interference effects, etc.

3.3.3 Active measurements

Active sensors measure the energy emitted by an antenna and scattered in the backward direction, expressed by the backscattering coefficient σ_0 . The coefficient σ_0 is sensitive to the scattering properties of the snowpack while thermal emission is negligible. Modeling the backscattering coefficient knowing the snow properties is also based on the radiative transfer theory and share common issues with modeling in passive mode. Most studies deal with radiative transfer (Bingham and Drinkwater, 2000) and the dense media radiative transfer theory (Zurk et al., 1996; Wen et al., 1990; Bingham and Drinkwater, 2000).

3.3.4 Application 1: continental snow cover extent

The most advanced application of microwave remote sensing on continental surfaces is probably the large scale determination of the areal extent of the snow cover (Grody and Basist, 1996; Mialon et al., 2005). The detection is based on the fact that snow attenuates the waves coming from the soil. In passive mode, the soil has a higher T_b than dry snow because it is both warmer and has a higher emissivity. As snow depth increases, the soil upwelling waves are attenuated and the brightness temperature decreases. The decreased amplitude depends on the snow scattering properties and this is usually enhanced at higher frequencies (except in special cases: Renseffl and Grody, 2000). This spectral signature is specific to the snow and can be used to discriminate snow from soil. In active mode, the soil acts as an opaque rough surface that backscatters efficiently the incoming wave. Soil backscattering coefficients are about one order of magnitude higher than those of dry snow. The snow layer attenuates the backscattered energy, thus decreasing the backscattering coefficient of bare soil.

The monitoring of snow areal extent using microwaves does not require sun illu-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

mination, an enormous advantage in the polar winter (Groody et al., 1991), and is furthermore insensitive to cloud cover. However, thin snow covers are undetected by microwave because of their larger penetration depth than in the visible/infrared, especially at the lowest frequencies (Armstrong and Brodzik, 2001). Working at 85 GHz improves detection but the sensitivity to atmospheric effects increases. Working in the visible or infrared further improves detection, but requires cloud-free conditions and day light. When possible, the combination of both microwave and visible/infrared observations offers the best potential (Romanov et al., 2000).

3.3.5 Application 2: snow water equivalent

A lot of effort has been devoted to the retrieval of Snow Water Equivalent (SWE: the height of water that would be obtained by melting a column of snow), for instance for hydrological applications (Grippa et al., 2005). SWE retrieval is based on the same principle as the snow areal extent, but the idea is here to quantify the absorption due to the snow covering the soil. Most studies use passive microwave as in the early work of Chang et al. (1987). The uncertainties in the method are mainly caused by the variations of attenuation due to grain size and density that change both spatially and through the season due to metamorphism. Other effects such as the presence of forest or wet snow further complicate the interpretation. Many methods have been developed to deal directly or indirectly with metamorphism including empirical methods (Josberger and Mognard, 2002; Grippa et al., 2004), statistical methods (Davis et al., 1993; Tedesco et al., 2003; Chang and Rango, 2000) and physical model-based methods (Pulliainen et al., 1999; Pulliainen, 2006; Wilson et al., 1999; Guo et al., 2003). Even under favorable conditions, such as over grassland, these methods are still insufficient for hydrological applications.

Methods using active microwaves have also been proposed (Bernier et al., 1999) to get much finer resolution (~100 m versus 25 km with passive microwave) at the expense of temporal resolution (monthly versus daily). Retrieval of SWE using the ERS altimeter including both active and passive measurements were also investigated

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(Papa et al., 2002).

3.3.6 Application 3: net accumulation

On ice sheets (Antarctica, Greenland), snow accumulates over years and the ground is far deeper than microwaves can reach. Annual net accumulation is thus a more suitable variable than SWE. The method to estimate SWE based on the presence of soil is not applicable. A dedicated method was developed for Greenland (Winebrenner et al., 2001) and Antarctica (Arthern et al., 2006) using passive microwaves at low frequency (6 GHz).

Snow is usually a stratified medium where density and grain size vary vertically. These variations induce changes of the dielectric constant that microwaves see as interfaces, and where wave reflection and transmission occur. The interesting point is that vertically (V) polarized waves are less reflected (more transmitted) than horizontally (H) polarised waves, especially near the Brewster angle (about 50°) where V polarized waves are not reflected at all. In passive mode, the waves emitted by the lowest layers and propagating upward are reflected backward and do not reach the upper layers. As a consequence, the V polarized waves escape more easily the medium than H polarized waves. The difference between H and V polarized brightness temperatures (all microwave sensors are polarisation-sensitive) is thus proportional to the number of layers within the penetration depth. Furthermore, using the observed inverse correlation between the accumulation rate and the number of layers in a given snow depth (Shuman et al., 2001) allows an estimation of the snow accumulation rate.

In active mode, a method to derive accumulation based on the difference between backscattering coefficients at various incidence angles was developed for Greenland (Drinkwater et al., 2001; Forster et al., 1999).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.3.7 Application 4: melting/refreezing and liquid water content

The presence of liquid water is easy to detect as it significantly increases the absorption. As absorption equals emissivity (Kirchoff's law), emissivity and in turn brightness temperature increase from dry to wet snow. In active mode, the increased absorption decreases the backscattering coefficient. Both active and passive sensors have been used to detect liquid water in snow and various techniques were developed for Greenland (Albdalati and Steffen, 1997 and 2001; Wiesmann, 2000), Antarctica (Torinesi et al., 2003; Picard and Fily, 2006; Liu et al., 2006) and boreal regions (Kimball et al., 2004). These methods have been successful and the next step in the future may be on deriving quantitatively the liquid water content rather than simply detecting the presence/absence of liquid water.

3.3.8 Application 5: other variables: density, grain size and temperature

While it has been known for a long time that microwaves were sensitive to snow variables such as density, grain size and temperature (e.g. Ulaby et al., 1981; Mätzler, 1994; Sherjal and Fily, 1994; Surdyk 2002), retrieving these variables is difficult even with a moderate accuracy. It is now clear that it is not possible to retrieve these variables individually, i.e. by stand-alone algorithms, as achieved for variables discussed in the previous sections. The simultaneous retrieval of many snow variables based on complex physical models and inversion techniques seems the most promising route. Earlier works on this route use neural network (NN) trained by an electromagnetic emission model (Tsang et al., 1992; Davis et al., 1993). Recently, various inversion techniques were used instead of the NN (Pulliainen et al., 2006; Flach et al., 2005; Guo et al., 2003; Wilson et al., 1999). In any case, the idea is the same: add physical information provided by forward models to constrain as much as possible the inversion. Future developments will likely make use of coupled forward models at multiple wavelengths from UV to microwaves, for instance models of the physical evolution of the snowpack coupled to a radiative model, to further constrain the inversions.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4 Developing snow photochemistry models from snow physics models

4.1 General considerations

Mathematical and numerical models are useful for testing hypotheses about snow-air interaction, the impact of snow photochemistry on atmospheric chemistry, and post-depositional process of snow properties and chemistry. The choice of conceptual approach, model development, and use of models depends strongly upon the desired application. There are literally thousands of different snow models, and their detailed description is beyond the scope of this paper. Here we discuss some of the existing models that simulate aspects of the physics relevant to air-snow exchange for the study of atmospheric chemistry. Similarly, there are at least hundreds of atmospheric chemistry models that simulate heat, momentum, and chemical exchange within the atmosphere, and it is beyond the scope of this chapter to describe all of them; none of them currently address processes in the underlying snow. Below, we will describe classes of existing models and stress where aspects of those models may be useful to the current topic.

The dimensionality of any model constrains the processes that it can explicitly address. Zero-dimension models cannot include effects of physical processes or movement, and are not used in physical modelling, however “box” models are used in atmospheric chemistry where they perform an accounting of the species for what is put into and taken out of the box. This is useful for problems involving complicated stoichiometry. One-dimensional models are needed for gradient-driven processes, such as diffusion, when the driving gradients act along a straight line. One-dimensional models are reasonable choices in snowmelt hydrology and in boundary layer atmospheric chemistry when the energy driving the exchange is primarily in the vertical direction, and in cases where movement in other directions can be ignored. Because situations involving flow and advection are rarely unidirectional, parameterizations of multidimensional phenomena must be developed if they are to be simulated in 1-D models. The use of eddy diffusivities is an example of a parameterization of a multidimensional sit-

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

uation for use in 1-D modeling. Multidimensional models are used when any geometry other than a straight line is important in the problem, and usually involve problems of convective flow and transport, although they are used in cases of diffusion when the geometry of the problem requires it. Examples where multidimensional models are needed include wind-driven interstitial air flow and transport over and through rough surfaces, investigations of flow and transport around sampling equipment, etc.

4.2 Snow physics model

A number of one-dimensional models exist for snow that are driven by a surface energy balance boundary condition and use data input for changing meteorological conditions to drive simulations of snow accumulation, densification, and melt. The domain of these models extends from the snow surface down to the ground or underlying strata. Examples of such models include SNTHERM (Jordan, 1991), CROCUS (Brun et al., 1992) and SNOWPACK (Lehning et al., 1999). The governing equation is 1-D heat conduction, snow is assumed to be a continuum, and parameterizations for effects of grain growth and changing snow properties over time are included. Models of this type are most often used for forecasting snow conditions for hydrological applications (e.g. Mote et al, 2003), solving heat transfer problems (e.g. Jordan and Andreas, 1999), and for use in modelling snow parameter evolution (e.g. Lefebre et al., 2003; Zappa et al., 2004).

Multidimensional snow physics models have been developed to study issues of flow and transport in snow. Like the models described above, their domain usually extends from the snow surface down to the ground or underlying strata, and they sometimes also have an upper boundary condition involving a surface energy balance using meteorological data to drive time-varying simulations. The governing equations involve a coupled set of partial differential equations, one that solves for the flow field using boundary conditions of pressure or prescribed flow, along with a partial differential equation describing advective-diffusive transport of a scalar (temperature or chemical species) with the flow. Examples of models in this category include the ventila-

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

tion model of Albert (1996), Albert and Shultz (2002), and Neumann and Waddington (2004). Modelling has shown that layering is a first-order effect in transport in the snow and that buried high-permeability layers can act as conduits for faster flowing air (Albert, 1999), that ventilation impacts the sublimation of snow (Albert, 2004) and the isotopic content of snow (Neumann and Waddington, 2004), and that sampling of firn air needs to account for surface air filtration to the subsurface for sampling chemistry in firn air (Albert et al., 2002). The snow properties that are required input for the snow physics models above include snow density, permeability, and thermal conductivity.

Since atmospheric input data cannot be obtained with the necessary resolution to make two- or three dimensional simulations of the snow in a real domain, the higher dimensional models have only been used for idealized process studies. Any real domain simulation of the snow cover will be with distributed one-dimensional models (Lehning et al., 2006) for any foreseeable future. The most advanced process representation has been achieved with physical models used for avalanche warning. The models SNOWPACK (Lehning et al., 1999) and CROCUS (Brun et al., 1991) have a full set of metamorphism parameters, which help describe not only the energy- and mass balance in snow but also the development of snow types, storing the full metamorphic history of deposited snow and linking the bulk properties conductivity and viscosity to snow microstructure (Lehning et al., 2002a). Lehning et al. (2002b) also show that the effect of snow ventilation can be included in a one-dimensional model using a parameterization of the thermal conductivity and diffusivity, although recent wind tunnel measurements show that this effect may have been overestimated previously.

4.3 Current approaches to model snow photochemistry

Models for investigating snow chemistry to date have each concentrated on one chemical species or one class of species, and have parameterized the creation and decay of that species in the code. Thus they have not included details of heterogeneous chemistry or photochemical processes. Models of this type include one-dimensional investigations of hydrogen peroxide in snow by McConnell et al. (1998), and formaldehyde

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

in snow by Hutterli et al. (1999) The models of McConnell and Hutterli incorporate the 1-D snow physics model SNTherm (Jordan, 1999) to calculate the snow temperature under varying meteorological conditions, but also include separate modules to calculate the diffusion of chemical species. Ventilation was accommodated in the McConnell model by increasing the gas diffusivity parameter. None of the 1-D models explicitly calculate the flow field, which is a multidimensional phenomenon. Two-dimensional models that calculate both the flow field and transport of chemical species have been used to investigate ozone destruction (Albert et al., 2002) and isotopic evolution in snow (Neumann and Waddington, 2004), both with parameterization of the chemical source/sink.

The aspects that these models need to describe include the transport of scalars such as heat, mass or chemical species. Such transports can be described by the governing transport equations for non-reactive gases, or reactive gases for which the reactions have been parameterized. For example, gas transport through snow is described by the following equation in one dimension, which allows for the parameterization of chemical reactions:

$$\phi_a \frac{\partial C_a}{\partial t} + \phi_a v \frac{\partial C_a}{\partial x} = \phi_a \frac{\partial}{\partial x} \left[D_a \frac{\partial C_a}{\partial x} \right] + S_c \quad (4.3-1)$$

Where C_a is the concentration of the species in the air space, t is time, x is spatial location, v is the interstitial air flow velocity (usually taken to be Darcy flow through snow), D_a is the diffusivity of the gas in snow, ϕ_a is the porosity, and S_c is the source term. If there is no air flow through the snow, then $v=0$, the velocity term on the left hand side of Eq. (4.3-1) drops out, and the interstitial transport in that case is by simple gas diffusion. The source term, S_c , may take on various forms depending on the type of chemistry modeling being used to estimate reversible transfer between the ice crystals and the interstitial air. Firn air chemistry models published to date can be considered parameterized models, in that they use parameterizations to relate gas concentrations in the ice to those in the firn air, rather than tackle the detailed stoichiometry involved in the reactions. A typical parameterized model estimates the

chemical exchange between the ice crystal and the firm air as :

$$-S_c = \theta_i k_b (C_a K_D - C_i) \quad (4.3-2)$$

Where k_b is a mass transfer coefficient, K_D is the ice-air equilibrium partition coefficient (both of which are temperature dependent), θ_i is the ice fraction, C_a is the concentration in air and C_i is the concentration in ice. Models of this type include those of McConnell et al. (1998) and Hutterli et al. (1999).

The interstitial air flow velocity is given by Darcy's law:

$$v = -\frac{K_p}{\mu} \frac{\partial P}{\partial x} \quad (4.3-3)$$

Where v is the interstitial air flow velocity, μ is the viscosity of air, K_p is the snow permeability, and $\partial P/\partial x$ is the spatial pressure gradient.

The analogous equations for heat transfer are given by Albert and McGilvary (1992). The transport of water vapor through the snow is described by

$$\frac{\partial \rho_v}{\partial t} + v \frac{\partial \rho_v}{\partial x} = \frac{\partial}{\partial x} \left[D_s \frac{\partial \rho_v}{\partial x} \right] + S_v \quad (4.3-4)$$

where ρ_v is water vapor density, v is the interstitial air flow (ventilation) velocity, D_s is the diffusion coefficient for vapor flow in snow, and S_v is the source of water vapor due to phase change, where

$$S_v = h_m SSA (\rho_{v,sat} - \rho_v) \quad (4.3-5)$$

where h_m is the mass transfer coefficient, SSA is the specific surface area of the snow, and $\rho_{v,sat}$ is the saturated vapor density (Albert and McGilvary, 1992). Experimental determination of the mass transfer coefficient in snow is given in Neumann et al. (2007)³.

³Neumann, T. A., Albert, M. R., Engel, C., Courville, Z., and Perron, F. E.: Experimental determination of snow sublimation Rate, J. Glaciol, submitted, 2007.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Albert and McGilvary (1992) demonstrated that the heat transfer associated with vapor transport is significant in the determination of the overall temperature profile of a ventilated snow sample, but that the major temperature effects are controlled by the balance between the heat carried by the dry air flow through the snow and heat conduction due to the temperatures imposed at the boundaries. For high porosity snows such as fresh snow or depth hoar, the thermal conductivity is sufficiently low that air flow through snow can cause advection-controlled temperature profiles. This was demonstrated in a field experiment (Hardy and Albert, 1995) where the immediate temperature effects from ventilation appeared when flow was induced in natural seasonal snow. For lower porosity, higher conductivity snow, such as wind-packed snow or firn, the thermal conductivity is sufficiently high that the temperature profile will be dominated by the heat conduction profile, even though there may be significant air flow through the snow (Albert and McGilvary, 1992).

Some simpler approaches have attempted to model snow chemistry without describing transport. In particular, persistent organic pollutants have a simple behavior in snow, because they are unreactive and are thought to undergo only adsorption/desorption processes, and box models have been used to describe their exchanges between the atmosphere and the snow (Daly and Wania, 2004). However, these models are not able to treat detailed snow processes and the numerical solution of a joint atmosphere-snow model with a detailed representation of snow models has not yet been achieved.

Next-generation snow photochemistry models will need to improve upon the 1D modeling above by including variables that are crucial to quantify chemistry, such as specific surface area (SSA) and light fluxes in the snow, and a more sophisticated description of the actual photochemical processes taking place in the snow. Coupling with 1D modeling of atmospheric chemistry will be required to permit investigations of the complicated phenomena across the atmosphere-snow interface.

Regarding SSA, this variable can change rapidly during metamorphism (Cabanes et al., 2003; Legagneux et al., 2004; Taillandier et al., 2007) and these changes must

be described. Since models such as SNOWPACK and CROCUS provide a detailed description of known snow metamorphism, it is possible that they can also be used to parameterize snow SSA. Domine et al. (2007) propose to determine SSA from density using empirical correlations, and show that also taking into account the snow crystal type, which is predicted by these models, leads to good predictions. Alternatively, it may be possible to use Eqs. (2.3-4) and (2.3-5) detailed in Sect. 2.3 to predict SSA from the age and metamorphic history of the snow. Today, however, the rate of SSA decay is not known for all common metamorphic scenarios, such as in high density windpacks, as mentioned in Taillandier et al. (2007). Integrating microphysical models of snow metamorphism (Flanner and Zender, 2006) that treat the grain to grain transfer of water vapor is another possibility to predict SSA evolution in snow. However, today, these models often treat snow crystals as spheres, and this is a clear source of concern for SSA prediction, as SSA is very shape-dependent.

Regarding the light flux in snow, models such as SNOWPACK and CROCUS also have versions that calculate the shortwave radiation flux in wavelength bands for each snow layer (Meirolid-Mautner and Lehning, 2004), and this may represent a first step towards predicting light fluxes in snow photochemistry models. However, as detailed in section 2e, the attenuation of UV and visible light in snow is mostly due to absorption by impurities. Since these absorbers mostly come from the atmosphere, an atmosphere-snow coupled model will probably be required to predict depth-resolved light fluxes in snow.

But the greatest difficulty in writing an integrated snow photochemistry model is doubtless the description of the photochemistry itself. Our current limitations in our understanding of this field is detailed in Grannas et al. (2007). In the next section, we briefly develop those aspects most related to snow physics.

4.4 The challenge of describing actual snow photochemistry in snow physics models

Snow forms a complex multiphase photocatalytic reactor under the influence of the solar radiation (e.g. Dominé and Shepson, 2002) where reactive species can be located

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

in the interstitial air, in aerosol particles trapped in the snow, within ice crystals, in the disordered surface layer of ice crystals, or at the ice-air interface. A full snow chemistry model has to consider chemical reactions in these different phases (often called compartments in models). Grain boundaries and triple junctions may also be considered an extra phase where reaction or physical fractionation occurs (e.g. Huthwelker et al., 2001). The difficulty in constructing a snow photochemistry model is that some of these compartments are poorly characterized. In particular, the nature of the surface layer of ice crystals has stirred much interest, interrogations and controversy (Baker and Nelson, 1996; Baker and Dash, 1996; Chen and Crutzen, 1994; Chen and Crutzen, 1996; Knight, 1996a; Knight, 1996b).

There is now clear experimental evidence that the ice surface is disordered and that the degree of disorder and the thickness of the disordered layer increases with temperature, so that it is often called the quasi-liquid layer (QLL) (e.g. Petrenko and Whitworth, 1999; Rosenberg, 2005). Its physical properties are intermediate between those of water and those of ice (Dash et al., 1995). Using ellipsometry, the existence of a surface layer down to about -10°C was shown (Beaglehole and Nason, 1980; Elbaum et al., 1993; Furukawa et al., 1987b). The value for the refractive index is between those of ice and of water (Furukawa et al., 1987a; Furukawa et al., 1987b). X-Ray diffraction revealed that the crystal structure is indeed disordered in the upper ~ 10 nm (Dosch, 1992; Dosch et al., 1995; Lied et al., 1994). Döppenschmidt and Butt (2000) used atomic force microscopy to determine that the thickness of the disordered layer ranged from a few nm near -25°C to a few tens of nm near the melting point. The diffusion constant value in the disordered region is in between those of ice and water (Mizuno and Hanafusa, 1987). Moreover, all observed effects increase with rising temperature, consistent with the theoretical prediction and molecular dynamics calculations that the disordered region should grow with rising temperature (Girardet and Toubin, 2001).

These various physical studies support the conclusion that the QLL is not simply a layer of supercooled water on top of the ice surface, because supercooled water would

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

freeze readily, once brought in contact with ice. Surface disorder occurs because it lowers the ice-air interfacial energy. Because the surface has liquid-like properties, a great simplification would be to use liquid phase reaction rate coefficients to model QLL chemistry, but there is at present no convincing basis that this is correct. Likewise, using the solubility of a trace gas in bulk water to estimate the overall ‘uptake into a QLL’ is probably incorrect, because the ‘solubility’ in such a layer is likely to differ strongly from that in bulk water. Moreover, as the surface disordered layer is not homogeneous, a net solubility is an ill-defined quantity.

Theoretical studies predict that surface disorder will increase in the presence of impurities (Beaglehole, 1991; Wettlaufer, 1999). Indeed, there is experimental evidence that the presence of HCl induces some surface disorder on the ice surface, even at temperatures as low as 200 K (McNeill et al., 2006) and a disordered phase forms in frozen brine solutions (Cho et al., 2002).

Grain boundaries and triple junctions that form during metamorphism form another poorly characterized compartment. Such interfacial reservoirs are possibly disordered. Moreover, since impurities are little soluble in ice, they are thought to concentrate in triple junctions. In these reservoirs, both the melting point depression induced by the dissolved impurities and the Kelvin effect on the curved surfaces in the triple junction thermodynamically stabilizes the liquid in the inclusion, as suggested theoretically (Nye, 1991) and studied experimentally (Mader, 1992a, b). Sulfates have indeed been found in triple junctions using different microscopic techniques (Baker et al., 2003; Barnes et al., 2002; Barnes and Wolff, 2004; Cullen and Baker, 2001; Cullen and Baker, 2002; Fukazawa et al., 1998; Mulvaney et al., 1988). Such reservoirs may be suspected to have a profound impact on chemical processes in snow. For example, it has been suggested that nitrate photolysis may occur there (Grannas et al., 2007).

In general, complex chemical transformations in multiphase systems are treated using differential equations of the following form:

$$\frac{dC_{l,n}}{dt} = P_{l,n} - L_{l,n} \cdot C_{l,n} + T_{l,n} \quad (4.4-1)$$

with $C_{l,n}$ concentration of the specie l in the compartment n , t time, $P_{l,n}$ chemical production of specie l in the compartment n , $L_{l,n}$ chemical sink of specie l in the compartment n , and $T_{l,n}$ transport loss or gain of specie l in the compartment n with respect to adjacent compartments. Because the chemical and transport terms depend on concentrations of other species or concentrations in different compartments, these equations constitute a set of dependent differential equations, which need to be solved using established integration schemes.

To apply the general equation, the concentrations of the species involved in the different compartments must be known. However, only very few concentration measurements in specific compartments are currently available. These include mainly firn air measurements of several reactive species (Grannas et al., 2007). Measurements of concentrations in the snow are restricted to stable compounds in melted snow samples, which is obviously the sum of the concentrations in condensed phases, possibly even including a fraction of the amount present in particles. It can be suggested that the fractionation of the impurities into the different snow compartments depends on the properties of the species and different methods can be used to estimate the partitioning. We expect that significant fractions of those species which fit easily into the ice structure are present in the solid compartment. For example, Jacobi et al. (2002) estimated from measurements above freshly deposited snow that more than 80% of the hydrogen peroxide in the fresh snow was located in the solid ice crystal. Ionic species are much more soluble in water than in ice (e.g. Gross and Svec, 1997; Thibert and Dominé, 1997 and 1998), and one would therefore expect ionic concentrations to be much higher in the QLL than in the solid, but there is not convincing evidence of this at present.

While soluble organic and inorganic compounds are likely present either in the QLL or in the solid ice crystal, significant fractions of insoluble compounds can also be adsorbed at the QLL-air interface. It is in fact at present not possible to deduce from adsorption measurements whether species taken up by the ice surface are located at the QLL-air interface or are dissolved in the QLL. Experimental studies of gas ad-

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

sorption on ice at temperatures relevant to most snowpacks ($T > -35^{\circ}\text{C}$) are limited (Huthwelker et al., 2006). Molecules studied at snowpack temperatures include nitric acid, sulfur dioxide, hydrogen peroxide, nitrogen oxides, 1-pentanol, acetic acid and phenanthrene on ice surfaces (e.g. Conklin et al., 1993; Clapsaddle and Lamb, 1989; Laird and Sommerfeld, 1995; Clegg and Abbatt, 2001; Bartels-Rausch et al., 2002; Sokolov and Abbatt, 2002; Domine et al., 2007²) and the sorption of a range of organic compounds on snow (Roth et al., 2004). Unfortunately, there is to our knowledge no study of adsorption over a wide enough temperature range to cover the region where the QLL is thin (of the order of one ice bi-layer) and thick ($> 10\text{ nm}$) so that the impact of the formation of the QLL on gas adsorption cannot be predicted. In particular, it is not known whether adsorption data obtained at low temperatures for species that can be expected to be soluble in the QLL (such as nitric acid and acetone) can be extrapolated to higher temperatures.

Despite these difficulties, Jacobi et al. (2004) attempted to determine the partitioning of hydrogen peroxide, nitric acid and acetic acid between the QLL and the QLL-air interface in the summertime snowpack at Summit for a snow sample at -19°C . They used chemical measurements of melted snow and of the gas phase and determined the concentration of the QLL-air interface by extrapolating low-temperature adsorption coefficients. They concluded that the amounts adsorbed at the QLL-air interface constituted insignificant fractions of hydrogen peroxide and nitric acid, but more than 20% of the snowpack acetic acid. While hydrogen peroxide and nitric acid were expected to be indeed mostly dissolved in the QLL, the result for acetic acid was somewhat expected from molecular dynamics calculations at 250 K (Compoint et al., 2002), which showed that that acid are trapped at the QLL-air interface.

Another possible approach that would avoid the difficulty of knowing where reactants are in the vicinity of the ice surface would be to use effective rate parameters that would include reactions both in the QLL and on the QLL-air interface. While many will not find this approach intellectually satisfying, it would enable the description of snow chemistry without requiring data that will probably not be available in the near future.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

One crucial parameter for the modeling of snowpack processes is the specific surface area (SSA). In equilibrium the amount of adsorbed species is directly proportional to the SSA according to the following equation:

$$n_{\text{ads}} = \theta \cdot \text{SSA} \cdot \rho_{\text{snow}} \quad (4.4-2)$$

5 with n_{ads} the number of adsorbed molecules per cm^3 , θ the surface coverage in molecules cm^{-2} , the SSA in $\text{cm}^2 \text{g}^{-1}$, and ρ_{snow} the density of the snow in g cm^{-3} . Moreover, the SSA also determines the volume of the thin QLL, V_{QLL} , which is proportional to the SSA.

$$V_{\text{QLL}} = d_{\text{QLL}} \cdot \text{SSA} \cdot \rho_{\text{snow}} \quad (4.4-3)$$

10 where d_{QLL} is the thickness of the QLL. Therefore, the thickness of the QLL is another crucial parameter for quantifying chemical processes occurring in the QLL.

Given the complexity of snow chemistry and the lack of detailed data on the reactivity of species in the different phases, several laboratory studies concerning photochemical reactions in snow and ice have been analyzed using the assumption that the reactions occur in the QLL, and that liquid phase rate coefficients could be applied to the QLL. For example, established aqueous phase reaction pathways and mechanisms were applied to the photolysis of nitrate in ice and snow (e.g. Honrath et al., 2000; Dubowski et al., 2001; Chu and Anastasio, 2003; Cotter et al., 2003; Boxe et al., 2005, 2006; Jacobi et al., 2006; Jacobi and Hilker, 2007). The authors justifies their approach a posteriori by the good agreement of the experimental results with the description using the liquid phase mechanism. Moreover, Chu and Anastasio (2003, 2005) demonstrated that the temperature dependent quantum yields of the nitrate and hydrogen peroxide photolysis in ice agree well with quantum yields obtained in aqueous solutions and extrapolated to sub-zero temperatures. More recently, Jacobi and Hilker (2007) proposed a simplified reaction mechanism including reaction rates for the photolysis of NO_3^- in natural snow based on the assumptions as described above. After adjusting the aqueous phase rate constants a good agreement between calculated and observed nitrate and nitrite

Snow physics as relevant to snow photochemistry

F. Domine et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

concentration-time-profiles was obtained. In summary, these laboratory studies indicate that for the chosen experimental conditions the investigated reactions seemed to proceed in a phase whose properties were similar to those of liquid water, and that according to the authors, was the QLL.

5 However, these results might be biased in some cases because higher concentrations than normally found in natural snow were applied leading to a thicker QLL with more 'liquid-like' properties than in natural snow, or even to the formation of a true aqueous phase as predicted by phase diagrams, if high enough concentrations were used. Nevertheless, some of these studies used impurity concentrations in the μM range typical of natural snow, and led to the same conclusions. A way to reconcile the conclusion of these studies that the QLL behaves as water with our earlier statements that the QLL and water are different may be to realize that the ice samples used in these laboratory experiments were dramatically different from natural snow. In particular, some studies used droplets of solution that were flash-frozen in liquid nitrogen as a snow mimic, but that resulted in small crystals with a large number of triple junctions, where a true liquid phase may have been present. This interpretation is consistent with the work of Huthwelker et al. (2001) who showed that the anomalous temperature dependence of the SO_2 uptake into packed ice bed would be inconsistent with uptake into a quasi-liquid layer on the ice surface, but with the dissolution into triple junctions. Hence, the observed chemistry could indeed be described using liquid phase rate coefficients. Clearly, understanding the microstructure of natural snow and snow mimics appears crucial to understand the phases in presence and the chemistry taking place. As alluded to above, other uncertainties include the exact nature of the QLL and its volume, the exact distribution of the impurities between the different phases, and in particular the location (particulate vs other phases) and speciation of organic impurities in the snow.

Snow physics as relevant to snow photochemistryF. Domine et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 Conclusions

This overview shows that snow is a complex multiphase medium whose physical properties change over time. Even though the physics governing these changes are not fully understood, many experimental studies have produced empirical relationships that have allowed the construction of snow physics model that reproduce in a satisfactory manner observations of snow evolution. Several complications arise if snow photochemistry is to be described by snow models.

The first one is to parameterize adequately variables of specific importance to photochemistry such as specific surface area and the light flux in the snow. Recent experimental and modeling studies of SSA (Legagneux et al., 2002; Flanner and Zender, 2006; Taillandier et al., 2007; Domine et al., 2007) provide some basis for the parameterization of SSA in future models. Regarding the light flux in snow, we have explained above that it could be predicted knowing snow scattering properties and absorption. Scattering is explained by snow physics, so that being able to predict snow density, SSA and grain shape should allow the parameterization of scattering. Absorption at the photochemically active wavelengths (UV and visible) are due mostly to deposited aerosols, whose amounts in snow depend on meteorology, atmospheric composition, snow permeability and surface structure. Therefore, coupling to an atmospheric model treating aerosols appears mandatory if light fluxes in snow are to be predicted with confidence.

The second one is to be able to describe the complex chemistry taking place in snow, and this is certainly the biggest challenge. At present, we do not understand where impurities are located in snow, so that writing the adequate chemical reactions is for the most part a guessing game. For example, is the nitrate ion that is photolyzed to produce nitrogen oxides present as a mineral salt in particles trapped in snow, as suggested by Beine et al. (2006), or as nitrate ions dissolved in the QLL at the surface of snow grains, as proposed in many laboratory studies and inferred from some field studies (e.g. Beine et al., 2002, Jacobi and Hilker, 2007). The nitrate ion could also be in triple junctions,

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

as suggested here in the proposed re-interpretation of the data of Jacobi and Hilker (2007). Clearly novel approaches are required to determine the nature and location of reactants in natural snow. Until a better understanding of this crucial aspect is reached, our progress in designing a predictive model of snow photochemistry will be severely limited.

Acknowledgements. This paper arose from a meeting held at LGGE, Grenoble, in May 2006. It was sponsored by the International Global Atmospheric Chemistry program (IGAC), the British Antarctic Survey, LGGE, Région Rhone-Alpes, Université Jopseph Fourier and the city of Grenoble. This paper is a contribution to the IGAC task on Air-Ice Chemical Interactions (AICI). We thank E. W. Wolff for his encouragements.

References

- Abdalati, W., and Steffen, K.: Snow melt on the Greenland ice sheet as derived from passive microwave satellite data, *J. Climate*, 10, 165–175, 1997.
- Abdalati, W. and Steffen, K.: Greenland ice sheet melt extent: 1979–1999, *J. Geophys. Res.*, 106, 33 983–33 989, 2001
- Albert, M. R. and McGilvary, W. R.: Thermal Effects Due to Air Flow and Vapor Transport in Dry Snow, *J. Glaciol.*, 38, 273–281, 1992.
- Albert, M. R.: Modeling heat, mass, and species transport in polar firn, *Ann. Glaciol.*, 23, 138–143, 1996.
- Albert, M. R., Koh, G., and Perron, F.: Radar investigations of melt pathways in a natural snowpack, *Hydrol. Processes*, 13, 2991–3000, 1999.
- Albert, M. R. and Perron Jr., F. E.: Ice layer and surface crust permeability in a seasonal snow pack. *Hydrological Processes*, 14(18), 3207–3214, 2000.
- Albert, M. R., Shultz, E., and Perron, F. E., Jr.: Snow and firn permeability measurements at Siple Dome, Antarctica. *Annals Glaciol.*, 31, 353–356, 2000.
- Albert, M. R., Grannas, A. M., Bottenheim, J. W., Shepson, P. B., and Perron Jr., F. E.: Processes and Properties of Snow-Air Transfer in the High Arctic with Application to Interstitial Ozone at Alert, Canada. *Atmos. Environ.*, 36, 2779–2787, 2002.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Albert, M. R. and Shultz, E.: Snow and firn properties and transport processes at Summit, Greenland, *Atmos. Environ.*, 36, 2789–2797, 2002.
- Albert, M. R., Shuman, C., Courville, Z., Bauer, R., Fahnestock, M., and Scambos, T.: Extreme Firn Metamorphism: Impact of Decades of Vapor Transport on Near-Surface Firn at a Low-Accumulation Glazed Site on the East Antarctic Plateau, *Ann. Glaciol.*, 39, 73–78, 2004.
- Alley, R. B. and Koci, B. R.: Ice-core analysis at site A, Greenland : preliminary results, *Ann. Glaciol.*, 10, 1–4, 1988.
- Alley, R. B., Saltzman, E. S., Cuffey, K. M., and Fitzpatrick, J. J.: Summertime formation of depth hoar in central Greenland, *Geophys. Res. Lett.*, 17, 2393–2396, 1990.
- Aoki, T., Aoki, T., Fukabori, M., Hachikubo, A., Tachibana, Y., and Nishio, F.: Effects of snow physical parameters on spectral albedo and bidirectional reflectance of snow surface, *J. Geophys. Res.*, 105D, 10219–10236, 2000.
- Armstrong, R. L. and Brodzik, M. J. : Recent Northern Hemisphere snow extent: a comparison of data derived from visible and microwave sensors, *Geophys. Res. Lett.*, 28(19), 3673–3676, 2001.
- Arthern, R. J., Winebrenner, R. J, and Vaughan, D. G.: Antarctic snow accumulation mapped using polarisation of 4.3 cm wavelength microwave emission, *J. Geophys. Res.*, 111, D06107, doi:10.1029/2004JD005667, 2006.
- Baker, I., Cullen, D., and Iliescu, D.: The microstructural location of impurities in ice, *Can. J. Phys.*, 81, 1–9, 2003.
- Baker, M. and Nelson, J.: Comment on "solute effects on the evaporation of ice particles" by Jen-Ping Chen and P.J. Crutzen, *J. Geophys. Res.*, 101(D17), 23 035–23 038, 1996.
- Baker, M. B. and Dash, J. G.: Surface layers on ice – Comment, *J. Geophys. Res.*, 101(D8), 12 929–12 931, 1996.
- Barnes, P. R. F., Mulvaney, R., Wolff, E. W., and Robinson, K.: A technique for the examination of polar ice using the scanning electron microscope, *J. Microsc.-Oxford*, 205, 118–124, 2002
- Barnes, P. R. F. and Wolff, E. W.: Distribution of soluble impurities in cold glacial ice, *J. Glaciol.*, 50(170), 311–324, 2004.
- Barnola, J. M., Pimienta P., Raynaud D., and Korotkevich Y. S.: CO₂ -climate relationship as deduced from the Vostok ice core: a re-examination based on new measurements and on a re-evaluation of the air dating, *Tellus*, 43B, 83–90, 1991.
- Bartels-Rausch, T., Eichler, B., Zimmermann, P., Gäggeler, H. W., and Ammann, M.: The adsorption enthalpy of nitrogen oxides on crystalline ice, *Atmos. Chem. Phys.* 2, 235–247,

Snow physics as relevant to snow photochemistry

F. Domine et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2002.

Bartelt, P. B. and Lehning, M.: A physical SNOWPACK model for Avalanche Warning Services. Part I: Numerical Model, *Cold Reg. Sci. Technol.*, 35/3, 123–145, 2002.

Beaglehole, D.: Surface Melting of Small Particles, and the Effects of Surface Impurities, *J. Crystal Growth*, 112(4), 663–669, 1991.

Beaglehole, D. and Nason, D.: Transition Layer on the Surface on Ice, *Surface Science*, 96(1–3), 357–363, 1980.

Beaglehole, D., Ramanathan, B., and Rumberg, J.: The UV and IR transmission of Antarctic snow, *J. Geophys. Res.*, 103(D8), 8849–8858, 1998.

Beine, H. J., Honrath, R. E., Domine, F., Simpson, W. R., and Fuentes, J. D.: NO_x During Background and Ozone Depletion Periods at Alert: Fluxes Above the Snow Surface, *J. Geophys. Res.*, 107 (D21), 4584, doi:10.1029/2002JD002082, 2002.

Beine, H. J., Domine, F., Simpson, W. R., Honrath, R. E., Sparapani, R., Zhou, X., and King, M. D.: Snow-pile and chamber experiments during the Polar Sunrise Experiment “Alert 2000”: Exploration of nitrogen chemistry, *Atmos. Env.*, 36, 2707–2719, 2002.

Beine, H. J., Domine, F., Ianniello, A., Nardino, M., Allegrini, I., Teinilä, K., and Hillamo, R.: Fluxes of Nitrates Between Snow Surfaces and the Atmosphere in the European High Arctic, *Atmos. Chem. Phys.*, 3, 335–346, 2003, <http://www.atmos-chem-phys.net/3/335/2003/>.

Beine, H. J., Amoroso, A., Dominé, F., M. D. King, M. D., Nardino M., Ianniello, A., and France, J. L.: Small HONO emissions from snow surfaces at browning pass, Antarctica, *Atmos. Chem. Phys.* 6, 2569–2580, 2006.

Bernier, M., Fortin, J. P., Gauthier, Y., Gauthier, R., Roy, R. and Vincent, P.: Determination of Snow Water Equivalent using RADARSAT in Eastern Canada, *Hydrol. Processes*, 13(18), 3031–3042, 1999.

Bingham, A. W. and Drinkwater, M.: Recent changes in the microwave scattering properties of the Antarctic ice sheet, *IEEE Trans, Geosci. Remote Sens.*, 38(4), 1810–1820, 2000.

Bohren, C. F. and Barkstrom, B. R.: Theory of the optical properties of snow, *J. Geophys. Res.*, 79, 4527–4535, 1974.

Bohren, C. F. and Beschta, R. L.: Snowpack albedo and snow density, *Cold Regions Sci. Technol.*, 1, 47–50, 1979.

Bottenheim, J. W., Barrie, L. W., Atlas, E., Heidt, L. E., Niki, H., Rasmussen, R. A., and Shepson, P. B.: Depletion of lower tropospheric ozone during Arctic spring: The Polar Sunrise

ACPD

7, 5941–6036, 2007

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

EGU

- Experiment 1988, *J. Geophys. Res.*, 95, 18 555–18 568, 1990.
- Bourdelles, B. and Fily, M.: Snow grain-size determination from Landsat imagery over Terre Adelie, Antarctica, *Ann. Glaciol.*, 17, 86–92, 1993.
- Boxe, C. S., Colussi, A. J., Hoffmann, M. R., Murphy, J. G., Wooldridge, P. J., Bertram, T. H., and Cohen, R. C.: Photochemical production and release of gaseous NO₂ from nitrate-doped water ice, *J. Phys. Chem. A*, 109, 8520–8525, 2005.
- Boxe, C. S., Colussi, A. J., Hoffmann, M. R., Perez, I. M., Murphy, J. G., and Cohen, R. C.: Kinetics of NO and NO₂ evolution from illuminated frozen nitrate solutions, *J. Phys. Chem. A*, 110, 3578–3583, 2006.
- Bremond, L., Alexandre, A., Hély, C., and Guiot, J.: A phytolith index as a proxy for tree cover density in tropical areas: calibration with Leaf area Index along a forest-savanna transect in southeastern Cameroon, *Global Planet. Change*, 45, 277–293, 2005.
- Brun, E. and Touvier, F.: Experimental study of thermal convection in snow *J. Phys. Colloque*. 48(C-1), 257–262, 1987.
- Brun, E., Martin, E., Simon, V., Gendre, C., and Coleou, C.: An energy and mass model of snow cover suitable for operational avalanche forecasting, *J. Glaciol.*, 121, 333–342, 1989.
- Brun, E., David, P., Sudul, M., and Brunot, G.: A numerical model to simulate snow-cover stratigraphy for operational avalanche forecasting, *J. Glaciol.*, 128, 13–22, 1992.
- Brunauer, S., Emmet, P. H., and Teller, E.: Adsorption of gases in multimolecular layers, *J. Am. Chem. Soc.*, 60, 309–319, 1938.
- Cabanes, A., Legagneux, L., and Dominé, F.: Rate of evolution of the specific surface area of surface snow layers, *Environ. Sci. Technol.*, 37(4), 661–666, 2003.
- Chen, J.-P. and Crutzen, P. J.: Solute effects on the evaporation of ice particles, *J. Geophys. Res.*, 99(D9), 18 847–18 859, 1994.
- Chang, A. T. C., Foster, J. L., and Hall, D. K.: Nimbus-7 SMMR derived global snow cover parameters, *Ann. Glaciol.*, 9, 39–44, 1987.
- Chang, A. T. C., Grody, N., Tsang, L., Basist, A., Goodison, B., Walker, A., Carroll, T., Armstrong, R., Josberger, E., and Sun, C.: Algorithm Theoretical Basis Document (ATBD) for AMSR-E snow water equivalent algorithm Greenbelt, MD: NASA Goddard Space Flight Center, Nov. 1997.
- Chen, J. P. and Crutzen, P. J.: Solute effects on the evaporation of ice particles – Reply, *J. Geophys. Res.*, 101(D17), 23 037–23 038, 1996.
- Cho, H., Shepson, P. B., Barrie L. A., Cowin J. P., and Zaveri, R.: NMR investigation of the

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- quasi-brine layer in ice/brine mixtures, *J. Phys. Chem. B*, 106, 11 226–11 232, 2002.
- Chu, L. and Anastasio, C.: Quantum yields of hydroxyl radical and nitrogen dioxide from the photolysis of nitrate on ice, *J. Phys. Chem. A*, 107, 9594–9602, 2003.
- Chu, L. and Anastasio, C.: Formation of hydroxyl radical from the photolysis of frozen hydrogen peroxide, *J. Phys. Chem. A*, 109, 6264–6271, 2005.
- Clapsaddle, C. and Lamb, D.: The sorption behaviour of SO₂ on ice at temperatures between –30°C and –5°C, *Geophys. Res. Lett.*, 16, 1173–1176, 1989.
- Clegg, S. M. and Abbatt, J. P. D.: Uptake of gas-phase SO₂ and H₂O₂ by ice surfaces: Dependence on partial pressure, temperature, and surface acidity, *J. Phys. Chem. A*, 105, 6630–6636, 2001.
- Cockell, C. S., Rettberg, P., Horneck, G., Wynn-Williams, D. D., Scherer, K., and Gugg-Helminger, A.: Influence of ice and snow covers on the UV exposure of terrestrial microbial communities: dosimetric studies, *J. Photochem. Photobiol. B-Biol.*, 68, 23–32, 2002.
- Colbeck, S. C.: An overview of seasonal snow metamorphism, *Rev. Geophys. Space Phys.*, 20, 45–61, 1982.
- Colbeck, S. C.: Theory of metamorphism of dry snow, *J. Geophys. Res.*, 88, 5475–5482, 1983.
- Colbeck, S., Akitaya, E., Armstrong, R., Gubler, H., Lafeuille, J., Lied, K., McClung, D., and Morris, E.: The International Classification for Seasonal Snow on the Ground, ICSI, IGS, 1990.
- Colbeck, S. C.: The layered character of snow covers, *Rev. Geophys.*, 29, 81–96, 1991.
- Coléou, C., Pieritz, R. A., Lesaffre, B., Brzoska, J. B., and Etchevers, P.: Isothermal metamorphism of a new snow layer: some measurements and simulation. In: *Proceedings ISSW 2004. International Snow Science Workshop, Jackson Hole WY, USA, 19–24 September 2004*, 2005.
- Compoin, M., Toubin, C., Picaud, S., Hoang, P. N. M., and Girardet, C.: Geometry and dynamics of formic and acetic acids adsorbed on ice, *Chem. Phys. Lett.*, 365, 1–7, 2002.
- Conklin, M. H., Sommerfeld, R. A., Laird, S. K., and Villinski, J. E.: Sulfur dioxide reactions on ice surfaces – implications for dry deposition to snow, *Atmos. Environ.*, 27, 159–166, 1993.
- Cotter, E. S. N., Jones, A. E. Wolff, E. W., and Baugitte S. J.-B.: What controls photochemical NO and NO₂ production from Antarctic snow? Laboratory investigation assessing the wavelength and temperature dependence, *J. Geophys. Res.*, 108, 4147, doi: 10.1029/2002JD002602, 2003.
- Courville, Z. R., Albert, M. R., Fahnstock, M. A., Cathles IV, L. M., Shuman, C. A.: Impacts of

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

an accumulation hiatus on the physical properties of firn at a low-accumulation polar site, in press, *J. Geophys. Res.*, 2007

Cullen, D. and Baker, I.: Observation of impurities in ice, *Microscopy Research and Technique*, 55(3), 198–207, 2001.

5 Cullen, D. and Baker, I.: Observation of sulfate crystallites in Vostok accretion ice, *Materials Characterization*, 48(4), 263–269, 2002.

Cunningham, J. and Waddington, E. D.: Air flow and dry deposition of non-sea salt sulfate in polar firn: paleoclimatic implications, *Atmos. Environ.*, 27A, 2943–2956, 1993.

10 Dash, J. G., Fu, H., and Wettlaufer, J. S.: The premelting of ice and its environmental consequences, *Rep. Prog. Phys.* 58, 115–167, 1995.

Davis, R. E., Dozier, J., and Perla, R.: Measurement of snow grain properties, in: *Seasonal Snowcovers: Physics, Chemistry, Hydrology*, edited by: Jones, H. G. and Orville-Thomas, W. J., D. Reidel Publishing Company, 63–74, 1987.

15 Davis, D. T., Chen, Z. Tsang, L. Hwang, J.-N., and Chang, A. T. C.: Retrieval of snow parameters by iterative inversion of a neural network, *IEEE Trans Geoscience Rem. Sens.*, 31(4), 842–852, 1993.

Dibb, J. E. and Arsenaault, M.: Shouldn't snowpacks be sources of monocarboxylic acids?, *Atmos. Environ.*, 36, 2513–2522, 2002.

20 Dominé, F., Thibert, E., Silvente, E., Legrand, M., and Jaffrezo, J.-L.: Determining past atmospheric HCl mixing ratios from ice core analyses, *J. Atmos. Chem.*, 21, 165–186, 1995.

Dominé, F., Cabanes, A., and Legagneux, L.: Structure, microphysics, and surface area of the Arctic snowpack near Alert during ALERT 2000, *Atmos. Environ.*, 36, 2753–2765, 2002.

Dominé, F. and Rey-Hanot, L.: Adsorption isotherms of acetone on ice between 193 and 213 K, *Geophys. Res. Lett.*, 29(18), 1873, doi:10.1029/2002GL015078, 2002.

25 Dominé, F. and Shepson P. B.: Air-snow interactions and atmospheric chemistry, *Science*, 297, 1506–1510, 2002.

Dominé, F., Lauzier, T., Cabanes, A., Legagneux, L. Kuhs, W. F., Techmer, K., and Heinrichs, T.: Snow metamorphism as revealed by scanning electron microscopy, *Microsc. Res. Tech.*, 62, 33–48, 2003.

30 Dominé, F. Sparapani, R., Ianniello, A., and Beine, H. J.: The origin of sea salt in snow on Arctic sea ice and in coastal regions, *Atmos. Chem. Phys.*, 4, 2259–2271, 2004,
<http://www.atmos-chem-phys.net/4/2259/2004/>.

Domine, F., Salvatori, R., Legagneux, L., Salzano, R., Fily, M., and Casacchia, R.:

Snow physics as relevant to snow photochemistry

F. Domine et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Correlation between the specific surface area and the short wave infrared (SWIR) reflectance of snow: preliminary investigation, *Cold Regions Sci. Technol.*, 6, 60–68, doi:10.1016/j.coldregions.2006.06.002, 2006.

Domine, F., Taillandier, A.-S., and Simpson, W. R.: A parameterization of the specific surface area of snow in models of snowpack evolution, based on 345 measurements, *J. Geophys. Res.*, in press, 2007.

Dosch, H.: Critical phenomena at surfaces and interfaces, *Springer Tracts in modern Physics*, 126, 1992.

Dosch, H., Lied, A., and Bilgram, J. H.: Glancing-Angle X-Ray-Scattering Studies Of The Pre-melting Of Ice Surfaces, *Surface Science*, 327(1–2), 145–164, 1995.

Döppenschmidt, A. and Butt, H.-J.: Measuring the thickness of the liquid-like layer on ice surfaces with atomic force microscopy, *Langmuir*, 16, 6709–6714, 2000.

Drinkwater, M. R., Long, D. G., and Bingham, A. W.: Greenland snow accumulation estimates from satellites radar scatterometer data, *J. Geophys. Res.*, 106, 33 935–33 950, 2001.

Dubowski, Y., Colussi, A. J., and Hoffmann, M. R.: Nitrogen dioxide release in the 302 nm band photolysis of spray-frozen aqueous nitrate solutions. Atmospheric implications, *J. Phys. Chem. A*, 105, 4928–4932, 2001.

Elbaum, M., Lipson, S. G., and Dash, J. G.: Optical Study of Surface Melting on Ice, *J. Cryst. Growth*, 129, 491–505, 1993.

Etchevers, P., Martin, E., Brown, R., et al.: Validation of the energy budget of an alpine snowpack simulated by several snow models (SnowMIP project), *Ann. Glaciol.*, 38, 150–158, 2004

Ewing, G. E.: Thin film water, *J. Phys. Chem. B*, 108(41), 15 953–15 961, 2004.

Fabre, A., Barnola, J.-M., Arnaud L., and Chappellaz, J.: Determination of gas diffusivity in polar firn: comparison between experimental measurements and inverse modeling, *Geophys. Res. Lett.*, 27, 557–560, 2000.

Fierz, C. and Lehning, M.: Assessment of the microstructure-based snow-cover model SNOWPACK: thermal and mechanical properties, *Cold Reg. Sci. Technol.*, 33, 123–213, 2001.

Fily, M., Bourdelles, B., Dedieu J. P., and Sergent, C.: Comparison of in situ and Landsat Thematic Mapper derived snow grain characteristics in the Alps, *Remote Sensing Environ.*, 59, 452–460, 1997.

Flach, J. D., Partington, K. C., Ruiz, C., Jeansou, E., and Drinkwater, M. R.: Inversion of the surface properties of ice sheets from satellite microwave IEEE Transactions on data,

- Geoscience and Remote Sensing, 43, 743–752, 2005.
- Flin, F., Brzoska, J. B., Lesaffre, B., Coléou, C., and Pieritz, R. A.: Full three-dimensional modelling of curvature-dependent snow metamorphism: first results and comparison with experimental tomographic data, *J. Phys. D: Appl. Phys.*, 36, 1–6, 2003.
- 5 Forster, R., Jezek, K. C., Bolzan, J., Baumgartner, F., and Gogineni, S. P.: Relationships between radar backscatter and accumulation rates on the Greenland ice sheet, *Int. J. Remote Sens.*, 20, 3131–3147, 1999.
- Franz, T. P. and Eisenreich, S. J.: Snow scavenging of polychlorinated biphenyls and polycyclic aromatic hydrocarbons in Minnesota, *Environ. Sci. Technol.*, 24, 1771–1778, 1998.
- 10 Fukazawa, H., Sugiyama, K., Mae, S., Narita, K., and Hondoh, T.: Acid ions at triple junction of Antarctic ice observed by Raman scattering, *Geophys. Res. Lett.*, 25, 2845–2848, 1998.
- Furukawa, Y., Yamamoto, M., and Kuroda, T.: Ellipsometric Study of the Ice Surface-Structure Just Below the Melting-Point, *J. Phys.*, 48(C-1), 495–501, 1987a.
- Furukawa, Y., Yamamoto, M., and Kuroda, T.: Ellipsometric Study of the Transition Layer on the
15 Surface of an Ice Crystal, *J. Cryst. Growth*, 82(4), 665–677, 1987b.
- George, C., Streckowski, R. S., Kleffmann, J., Stemmler, K., and Ammann, M.: Photoenhanced uptake of gaseous NO₂ on solid organic compounds: A photochemical source of HONO?, *Faraday Discuss.*, 130, 195–210, 2005.
- Giddings, J. C. and LaChapelle, E. R.: Diffusion theory applied to radiant energy distribution and albedo of snow, *J. Geophys. Res.*, 66, 181–189, 1961.
- 20 Girardet, C. and Toubin, C.: Molecular atmospheric pollutant adsorption on ice: a theoretical survey, *Surf. Sci. Reports*, 44, 159–238, 2001.
- Grannas, A. M., Shepson, P. B., Guimbaud, C., Sumner, A. L., Albert, M., Simpson, W., Dominé, F., Boudries, H., Bottenheim, J. W., Beine, H. J., Honrath, R., and Zhou, X.: A study of carbonyl compounds and photochemistry in the arctic atmospheric boundary layer, *Atmos. Environ.* 36, 2733–2742, 2002.
- 25 Grannas, A. M., Jones, A. E., Dibb, J., Ammann, M., Anastasio, C., Beine, H., Bergin, M., Bottenheim, J., Boxe, C.S., Carver, G., Crawford, J. H., Domine, F., Frey, M. M., Guzman, M. I., Heard, D., Helmig, D., Hoffmann, M. R., Honrath, R. E., Huey, L. G., Jacobi, H.-W., Klan, P., McConnell, J., Sander, R., Savarino, J., Shepson, P. B., Simpson, W. R., Sodeau, J., von Glasgow, R., Weller, R., Wolff, E., and Zhu, T.: Snow photochemistry: Current state of the science, *Atmos. Chem. Phys.*, 7, 4165–4283, 2007,
<http://www.atmos-chem-phys.net/7/4165/2007/>.

Snow physics as relevant to snow photochemistryF. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Grippa, M., Mognard, N. M., Letoan, T., and Josberger, E. G.: Siberia snow depth climatology from SSM/I data using a combined dynamic and static algorithm, *Rem. Sens. Environ.*, 93, 30–41, 2004.
- Grenfell, T. C., Perovich, D. K., and Ogren J. A.: Spectral albedos of an alpine snowpack, *Cold Regions Sci. Technol.*, 4, 121–127, 1981.
- Grenfell, T. C., Warren, S. G., and Mullen, P. C.: Reflection of solar radiation by the Antarctic snow surface at ultraviolet, visible, and near-infrared wavelengths, *J. Geophys. Res.*, 99, 18 669–18 684, 1994.
- Grenfell, T. C. and Warren S. G.: Representation of a nonspherical ice particle by a collection of independent spheres for scattering and absorption of radiation, *J. Geophys. Res.*, 104, 31 697–31 709, 1999.
- Grippa, M., Mognard, N. M., and Letoan, T.: Comparison between the interannual variability of snow parameters derived from SSM/I and the Ob river discharge, *Rem. Sens. Environ.*, 98, 35–44, 2005.
- Grody, N. C. and Basist, A. N.: Global Identification of snowcover using SSM/I measurements, *IEEE Transc. Geosc. Remote Sens.*, 34, 237–249, 1996.
- Gross, G. W. and Svec, R. K.: Effect of ammonium on anion uptake and dielectric relaxation in laboratory-grown ice columns, *J. Phys. Chem. B*, 101, 6282–6284, 1997.
- Guo, J., Tsang L., Josberger, E. G. Wood, A. W. Hwang, J.-N., and Lettenmaier, D. P.: Mapping the spatial distribution and time evolution of snow water equivalent with passive microwave measurements, *IEEE Trans. Geosci. Rem. Sens.*, 41(3), 612–621, 2003.
- Guo, J., Tsang, L., Josberger, E. G., Wood, A. W., and Hwang, J.-N.: Mapping the spatial distribution and time evolution of snow water equivalent with passive microwave measurements, *IEEE TGARS*, 41, 612–621, 2003.
- Hall, D. K., Riggs, G. A., and Salomonson, V.: Algorithm Theoretical Basis Document (ATBD) for the MODIS Snow and Sea Ice-Mapping Algorithms, available at http://modis.gsfc.nasa.gov/data/atbd/atbd_mod10.pdf, 2001.
- Hansen, J. and Nazarenko, L.: Soot climate forcing via snow and ice albedos, *Proc. Nat. Acad. Sci.*, 101, 423–428, 2004.
- Hanson, D. R. and Ravishankara, A. R.: Comment on Porosities of Ice Films Used to Simulate Stratospheric Cloud Surfaces – Response, *J. Phys. Chem.*, 97, 2802–2803, 1993.
- Harder, S., Warren, S. G., and Charlson, R. J.: Sulfate in air and snow at the South Pole: Implications for transport and deposition at sites with low snow accumulation, *J. Geophys.*

Snow physics as relevant to snow photochemistryF. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Res., 105D, 22 825–22 832, 2000.
- Hardy, J. P. and Albert, M. R.: Snow-Induced Thermal Variations Around a Single Conifer Tree, *Hydrol. Processes*, 9, 923–934, 1995.
- Herbert, B. M. J., Halsall, C. J., Jones, K. C., and Kallenborn, R.: Field investigation into the diffusion of semi-volatile organic compounds into fresh and aged snow, *Atmos. Environ.*, 40, 1385–1393, 2006.
- Honrath, R. E., Peterson, M. C., Guo, S., Dibb, J. E., Shepson, P. B., and Campbell, B.: Evidence of NO_x production within or upon ice particles in the Greenland snowpack, *Geophys. Res. Lett.*, 26, 695–698, 1999.
- Honrath, R. E., Guo, S., Peterson, M. C., Dziobak, M. P., Dibb, J. E., and Arsenault, M. A.: Photochemical production of gas phase NO_x from ice crystal NO_3^- , *J. Geophys. Res.* 105, 24 183–24 190, 2000.
- Honrath, R., Peterson, M. C., Lu, Y., Dibb, J. E., Arsenault, M. A., Cullen, N. J., and Steffen, K.: Vertical fluxes of nitrogen oxides above the snowpack at Summit, Greenland, *Atmos. Environ.*, 36, 2629–2640, 2002.
- Hudson, S. R., Warren, S. G., Brandt, R. E., Grenfell, T. C., and Six, D.: Spectral bi-directional reflectance of Antarctic snow: measurements and parameterization, *J. Geophys. Res.*, 111, D18106, doi:10.1029/2006JD007290, 2006.
- Huthwelker, T., Lamb, D., Baker, M., Swanson, B., and Peter, T.: Uptake of SO_2 by polycrystalline water ice, *J. Colloid. Interf. Sci.*, 238, 147–159, 2001.
- Huthwelker, T., Ammann, M., and Peter, T.: The uptake of acidic gases on ice, *Chem. Rev.*, 106, 1375–1444, 2006.
- Hutterli, M. A., Rothlisberger, R., and Bales, R. C.: Atmosphere-to-snow-to-firn transfer studies of HCHO at Summit, Greenland, *Geophys. Res. Lett.*, 26, 1691–1694, 1999.
- Hutterli, M. A., Bales, R. C., McConnell, J. R., and Stewart, R. W.: HCHO in Antarctic snow: preservation in ice cores and air-snow exchange, *Geophys. Res. Lett.*, 29, 76-1–76-4, 2002.
- Hutterli, M. A., McConnell, J. R., Chen, G., Bales, R. C., Davis, D. D., and Lenschow, D. H.: Formaldehyde and hydrogen peroxide in air, snow and interstitial air at South Pole, *Atmos. Environ.*, 38, 5439–5450, 2004.
- Jacobi, H.-W., Frey, M. M., Hutterli, M. A., Bales, R. C., Schrems, O., Cullen, N. J., Steffen, K., and Koehler, C.: Measurements of hydrogen peroxide and formaldehyde exchange between the atmosphere and surface snow at Summit, Greenland, *Atmos. Environ.*, 36, 2619–2628, 2002.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Jacobi, H.-W., Bales, R. C., Honrath, R. E., Peterson, M. C., Dibb, J. E., Swanson, A. L., and Albert, M. R.: Reactive trace gases measured in the interstitial air of surface snow at Summit, Greenland, *Atmos. Environ.*, 38, 1687–1697, 2004.

Jacobi, H.-W., Annor, T., and Quansah, E.: Investigation of the photochemical decomposition of nitrate, hydrogen peroxide, and formaldehyde in artificial snow, *J. Photochem. Photobiol. A*, 179, 330–338, 2006.

Jacobi, H.-W. and Hilker, B.: A mechanism for the photochemical transformation of nitrate in snow, *J. Photochem. Photobiol. A*, 185, 371–382, 2007.

Jones, A. E., Weller, R., Wolf, E. W., and Jacobi, H.-W.: Speciation and rate of photochemical NO and NO₂ production in Arctic Snow, *Geophys. Res. Lett.*, 27(3), 345–348, 2000.

Jones, A. E., Weller, R. Anderson, P. S., Jacobi, H.-W., Wolff, E. W., Schrems, O., and Miller, H.: Measurements of NO_x emissions from the Antarctic snowpack, *Geophys. Res. Lett.*, 28, 1499–1502, 2001.

Jordan, R.: A one-dimensional temperature model for a snow cover, Technical Documentation for SNTHERM.89, CRREL special report 91-16, U.S. Army Corps of Engineers, 64, 1991.

Jordan, R. E., Hardy, J. P., Perron, F. E., and Fisk, D. J.: Air permeability and capillary rise as measures of the pore structure of snow: an experimental and theoretical study, *Hydrol. Process.* 13, 1733–1753, 1999.

Jordan, R., Albert, M., and Brun, E.: Physical Processes in Snow and their Parameterization, in *Snow and Climate*, eds, Elsevier Press, in press, 2007.

Josberger, E. G. and Mognard, N.: A passive microwave snow depth algorithm with a proxy for snow metamorphism, *Hydrol. Processes*, 16, 1557–1568, 2002.

Judson, A. and Doesken, N.: Density of freshly fallen snow in the Central Rocky Mountains, *Bull. Am. Meteorol. Soc.*, 81, 1577–1587, 2000.

Kappen L.: Plant activity under snow and ice, with particular reference to lichens, *Arctic*, 46, 297-302, 1993.

Keyser, L. F., Moore, S. B., and Leu, M. T.: Surface-Reaction and Pore Diffusion in Flow-Tube Reactors, *J. Phys. Chem.*, 95, 5496–5502, 1991.

Keyser, L. F., Leu, M. T., and Moore, S. B.: Comment on Porosities of Ice Films Used to Simulate Stratospheric Cloud Surfaces, *J. Phys. Chem.*, 97, 2800–2801, 1993.

Kimball, J. S., McDonald, K. C., Frolking, S., and Running, S. W.: Radar remote sensing of the spring thaw transition across a boreal landscape, *Rem. Sens. Env.*, 89, 163–175, 2004.

King, M. D. and Simpson, W. R.: Extinction of UV radiation in Arctic snow at Alert, Canada

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- (82° N), *J. Geophys. Res.*, 106(D12), 12 499–12 508, 2001.
- Knight, C. A.: Surface layers on ice, *J. Geophys. Res.-Atmos.*, 101(D8), 12 921–12 928, 1996a.
- Knight, C. A.: Surface layers on ice – Reply, *J. Geophys. Res.-Atmos.*, 101(D8), 12 933–12 936, 1996b.
- 5 Kojima, K.: A field experiment on the rate of densification, *Proceedings of the Grindelwald Symposium*, IAHS Publication, 114, 298–308, 1975.
- Kokhanovsky, A. A. and Zege, E. P.: Scattering optics of snow, *Appl. Opt.*, 43, 1589–1602, 2004.
- Kokhanovsky, A. A.: Scaling constant and its determination from simultaneous measurements
10 of light reflection and methane adsorption by snow samples, *Opt. Lett.*, 31, 3282–3284, 2006.
- Kubelka, P. and Munk, F.: Ein Beitrag zur Optik der Farbanstriche, *Z. Tech. Phys. (Leipzig)* 12, 593-601, 1931.
- Kubelka, P.: New contribution to the optics of intensely light-scattering materials, Part I, *J. Opt. Soc. Am.*, 38, 448-457, 1948.
- 15 Kuhn, M. and Siogas, L.: Spectroscopic studies at McMurdo, South Pole and Siple stations during the austral summer 1977–78, *Antarctic Journal of the United States*, 13, 178–179, 1978.
- Laird, S. K. and Sommerfeld, R. A.: Nitric acid adsorption on ice: A preliminary study, *Geophys. Res. Lett.*, 22, 921–924, 1995.
- 20 Lange, M.: Measurement of thermal parameters in Antarctic snow and firn, *Ann. Glaciol.*, 6, 100–104, 1985.
- Lee-Taylor, J. and Madronich, S.: Application of TUV Tropospheric Radiative Transfer Model to Snowpack Photochemistry, *J. Geophys. Res.*, 107, 4796, 2002.
- 25 Legagneux, L., Cabanes, A., and Domine, F.: Measurement of the specific surface area of 176 snow samples using methane adsorption at 77 K, *J. Geophys. Res.*, 107(D17), 4335, doi:10.1029/2001JD001016, 2002.
- Legagneux, L., Lauzier, T., Dominé, F., Kuhs, W. F., Heinrichs, T., and Techmer, K.: Rate of decay of the specific surface area of snow during isothermal experiments and morphological changes studied by scanning electron microscopy, *Can. J. Phys.* 81, 459–468, 2003.
- 30 Legagneux, L., Taillandier, A.-S., and Domine, F.: Grain growth theories and the isothermal evolution of the specific surface area of snow, *J. Appl. Phys.*, 95, 6175–6184, 2004.
- Legrand, M., A. Léopold, A., and Dominé, F.: Acidic gases (HCl, HF, HNO₃, HCOOH and

Snow physics as relevant to snow photochemistryF. Domine et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- CH₃COOH): a review of ice core data and some preliminary discussions on their air-snow relationships, in: NATO ASI Series, vol I 43 "Chemical exchange between the atmosphere and polar snow", edited by: Wolff, E. and Bales, R. C., Springer, Berlin, 19–44, 1996.
- 5 Lehning, M., Bartelt, P., Brown, R. L., Russi, T., Stöckli, U., and Zimmerli, M.: Snowpack Model Calculations for Avalanche Warning based upon a new Network of Weather and Snow Stations, *Cold Reg. Sci. Technol.*, 30, 145–157, 1999.
- Lehning, M., Bartelt, P., Brown, B., Fierz, C., and Satyawali, P.: A physical SNOWPACK model for the Swiss avalanche warning Part II: Snow microstructure, *Cold Regions Sci. Technol.*, 35, 147–167, 2002.
- 10 Lehning, M., Bartelt, P. B., Brown, R. L., Fierz, C., and Satyawali, P.: A physical SNOWPACK model for the Swiss Avalanche Warning Services. Part III: Meteorological Boundary Conditions, Thin Layer Formation and Evaluation, *Cold Reg. Sci. Technol.*, 35, 169–184, 2002.
- Lehning, M., Völsch, I., Gustafsson, D., Nguyen, T. A., Stähli, M., and Zappa, M.: ALPINE3D: A detailed model of mountain surface processes and its application to snow hydrology, *Hydrol. Processes*, 20, 2111–2128, 2006.
- 15 Leu, M. T., Keyser, L. F., and Timonen, R. S.: Morphology and surface areas of thin ice films, *J. Phys. Chem. B*, 101, 6259–6262, 1997.
- Li, W., Stamnes K., and Chen, B.: Snow grain size retrieved from near-infrared radiances at multiple wavelengths, *Geophys. Res. Lett.*, 28, 1699–1702, 2001.
- 20 Lied, A., Dosch, H., and Bilgram, J. H.: Surface Melting Of Ice I(H) Single-Crystals Revealed By Glancing Angle X-Ray-Scattering, *Phys. Rev. Lett.*, 72(22), 3554–3557, 1994.
- Liljquest, G. H.: Energy exchange of an Antarctic snow-field, in *Norwegian-British-Swedish Antarctic Expedition 1949–1952, Scientific results vol. 2*, 93–103, Norsk Polarinstitut, Oslo, 1956.
- 25 Liu, H., Wang, L., and Jezek, K. C.: Spatiotemporal variations of snowmelt in Antarctica derived from satellite scanning multichannel microwave radiometer and Special Sensor Microwave Imager data (1978–2004), *J. Geophys. Res.*, 111, F01003, doi:10.1029/2005JF000318, 2006
- Luciano, G. L. and Albert, M. R.: Bi-directional Permeability Measurements of Polar Firn, *Ann. Glaciol.*, 35, 63–66, 2002.
- 30 Macelloni, G., Paloscia, S., Pampaloni, P., and Tedesco, M.: Microwave emission from dry snow: a comparison of experimental and model results, *IEEE Trans. Geosci. Rem. Sens.*, 39(12), 2649–2656, 2001.

Snow physics as relevant to snow photochemistryF. Domine et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Mader, H. M.: Observations of the Water-Vein System in Polycrystalline Ice, *J. Glaciol.*, 38, 333–347, 1992a.
- Mader, H. M.: The Thermal-Behavior of the Water-Vein System in Polycrystalline Ice, *J. Glaciol.*, 38, 359–374, 1992b.
- 5 Madronich, S.: Photodissociation in the atmosphere 1. Actinic flux and the effects of ground reflections and clouds, *J. Geophys. Res.*, 92, 9740–9752, 1987.
- Marbouty, D.: An experimental study of temperature gradient metamorphism, *J. Glaciol.*, 26, 303–312, 1980.
- Marsh, P. and Woo, M.-K.: Wetting Front Advance and Freezing of Meltwater Within a Snow
10 Cover, 1. Observations in the Canadian Arctic, *Water Resour. Res.*, 20, 1853–1864, 1984.
- Massman, W. J.: A review of the molecular diffusivities of H₂O, CO₂, CH₄, CO, O₃, SO₂, NH₃, N₂O, NO, and NO₂ in air, O₂ and N₂ near STP, *Atmos. Environ.*, 32, 1111–1127, 1998.
- Matzl, M. and Schneebeli, M.: Measuring specific surface area of snow by near-infrared photography, *J. Glaciol.*, 52, 558–564, 2006.
- 15 Mätzler, C.: Passive microwave signatures of landscapes in winter, *Meteorol. Atmos. Phys.*, 54, 241–260, 1994.
- McConnell, J. R., Bales, R. C., Stewart, R. W., Thompson, A. M., Albert, M. R., and Ramos R.: Physical based modeling of atmosphere-to-snow-to-firn transfer of H₂O₂ at South Pole, *J. Geophys. Res.*, 103(D9), 10561–10570, 1998.
- 20 McNeill, V. F., Loerting, T., Geiger, F. M., Trout, B. L., and Molina, M. J.: Hydrogen chloride-induced surface disordering on ice. *Proceedings of the National Academy of Sciences of the United States of America*, 103(25), 9422–9427, 2006.
- Meier, R. R., Anderson, G. P., Cantrell, C. A., Hall L. A., Lean, J., Minschwaner, K., Shetter, R. E., Shettle, E. P., and Stamnes K.: Actinic radiation in the terrestrial atmosphere, *J. Atmos. Solar-Terr. Phys.*, 59, 2111–2157, 1997.
- 25 Meirolid, I. and Lehning, M.: Measurements and model calculations of the solar shortwave fluxes in snow on Summit, Greenland, *Ann. Glaciol.*, 38, 279–284, 2004.
- Mellor, M.: Engineering properties of snow, *J. Glaciol.*, 19, 15–66, 1977.
- Mialon, A., Fily, M., and Royer, A.: Seasonal snow cover extent from microwave remote sensing data: Comparison with existing ground and satellite based measurements, *EARSel eProceedings*, 4(2), 215–225, 2005.
- 30 Mie, G.: Bitrage zur Optik trüber Medien speziell kolloidaler Metallosungen, *Annalen der Physik*, 25, 377–445, 1908.

Snow physics as relevant to snow photochemistryF. Domine et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Mishchenko, M. I., Dlugach, J. M., Yanovitskij, E. G., and Zakharova N. T.: Bidirectional reflectance of flat, optically thick particulate layers : an efficient radiative transfer solution and applications to snow and soil surfaces, *J. Quant. Spectrosc. Radiat. Transfer*, 63, 409–432, 1999.

5 Mizuno, Y. and Hanafusa, N.: Studies of Surface-Properties of Ice Using Nuclear-Magnetic-Resonance, *J. Phys.*, 48(C-1), 511–517, 1987.

Mulvaney, R., Wolff, E., and Oates, K.: Sulphuric acid at grain boundaries in Antarctic ice, *Nature*, 331, 247–249, 1988.

Nakaya, U.: *Snow Crystals: Natural and Artificial*. Harvard: Harvard University Press, 1954.

10 Narita, H.: Specific surface of deposited snow II, *Low Temp. Sci. A29*, 69–81, 1971.

Navarre, J. P.: Modèle unidimensionnel d'évolution de la neige déposée. *Modèle perce-neige, Météorologie*, 4, 103–120, 1975.

Neshyba, S., Grenfell, T. C., and Warren, S. G.: Representation of a nonspherical ice particle by a collection of independent spheres for scattering and absorption of radiation: II. Hexagonal columns and plates, *J. Geophys. Res.*, 108, 4448, doi:10.1029/2002JD003302, 2003.

15 Neumann, T. A. and Waddington, E. D.: Effects of firn ventilation on isotopic exchange, *J. Glaciol.*, 50, 183–194, 2004.

Nolin, A. W. and Dozier, J.: Estimating snow grain size using AVIRIS data, *Remote Sensing Environ.*, 44, 231–238, 1993.

20 Nolin, A. W. and Dozier, J.: A hyperspectral method for remotely sensing the grain size of snow, *Remote Sensing Environ.*, 74, 207–216, 2000.

Nye, J. F.: Thermal behavior of glacier and laboratory ice, *J. Glaciol.*, 37, 401–413, 1991.

Painter, T. H., Dozier, J., Roberts, D. A., Davis, R. E., and Greene, R. O.: Retrieval of sub-pixel snow-covered area and grain size from imaging spectrometer data, *Remote Sensing Environ.*, 85, 64–77, 2003.

25 Papa, F., Legresy, B., Mognard, N. M., Josberger, E. G., and Remy, F.: Estimating terrestrial snow depth with the TOPEX-Poseidon altimeter and radiometer, *IEEE Transactions on Geoscience and Remote Sensing*, 40(10), 2162–2169, 2002.

Perla, R., Dozier, J., and Davis, R. E.: Preparation of serial sections in dry snow specimens, *J. Microsc.* 141, 111–114, 1986.

30 Perovich, D. K. and Govoni, J. W.: Absorption coefficients of ice from 250 to 400 nm, *Geophys. Res. Lett.*, 18, 1233–1235, 1991.

Petrenko, V. F. and Whitworth, R. W.: *Physics of ice*, Oxford University Press, Oxford, UK,

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1999.

Picard, G. and Fily, M., Surface melting observations in Antarctica by microwave radiometers: correcting 26-year long time series from changes in acquisition hours, *Remote Sensing Environ.*, 104, 325–336, 2007.

5 Pielke, R. A., Liston, G. E., Chapman, W. L., and Robinson, D. A.: Actual and insolation-weighted northern hemisphere snow cover and sea ice between 1973–2002, *Clim. Dyn.*, 22, 591–595, 2004.

Phillips, G. J. and Simpson, W. R.: Verification of snowpack radiation transfer models using actinometry, *J. Geophys. Res.*, 110, D08306, doi:10.1029/2004JD005552, 2005.

10 Pulliainen, J. T.: Mapping of snow water equivalent and snow depth in boreal and sub-arctic zones by assimilating space-borne microwave radiometer data and ground-based observations, *Rem. Sens. of Environ.*, 101, 257–269, 2006.

Pulliainen, J. T. Grandell, J., and Hallikainen, M. T.: HUT snow emission model and its applicability to snow water equivalent retrieval, *IEEE Trans Geoscience and Rem. Sens.*, 37(3), 1378–1390, 1999.

15 Qiu, R., Green, S. A., Honrath, R. E., Peterson, M., Lu, C. Y., and Dziobak M.: Measurements of $J(\text{NO}_3^-)$ in snow by nitrate-based chemical actinometry, *Atmos. Env.*, 36, 2751, 2002.

Raymond, C. F. and Tusima, K.: Grain coarsening of water-saturated snow, *J. Glaciol.*, 86, 83–105, 1979.

20 Rick, U. and Albert, M. R.: Microstructure and Permeability in the Near-Surface Firn Near a Potential U.S. Deep Drilling Site in West Antarctica, *Ann. Glaciol.*, 39, 62–66, 2004.

Romanov, P., Gutman, G., and Csiszar, I.: Automated Monitoring of Snow Cover over North America with Multispectral Satellite Data, *J. Appl. Meteorol.*, 39, 1866–1880, 2000.

Rosenberg, R.: Why is ice slippery?, *Physics Today*, 58, 50–55, 2005.

25 Rosenfeld, S. and Grody, N.: Anomalous microwave spectra of snow cover observed from Special Sensor Microwave/Imager measurements, *J. Geophys. Res.*, 105, 14913–14926, 2000.

Roth, C. M., Goss, K.-U., and Schwarzenbach, R. P.: Sorption of diverse organic vapors to snow, *Environ. Sci. Technol.*, 38, 4078–4084, 2004.

30 Rozanov, V. V. and Kokhanovsky, A. A.: Semianalytical cloud retrieval algorithm as applied to the cloud top altitude and the cloud geometrical thickness determination from top-of-atmosphere reflectance measurements in the oxygen A band, *J. Geophys. Res.*, 109, D05202, doi:10.1029/2003JD004104, 2004.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Roy, V., Goïta, K., Royer, A., Walker, A. E., and Goodison, B. E.: Snow Water Equivalent Retrieval in a Canadian Boreal Environment From Microwave Measurements Using the HUT Snow Emission Model, *IEEE Trans. Geosci. Remote Sensing*, 42(9), 1850–1859, 2004.
- Schneebeli, M. and Sokratov, S. A.: Tomography of temperature gradient metamorphism of snow and associated changes in heat conductivity, *Hydrol. Processes*, 18, 3655–3655. doi:10.1002/hyp.5800, 2004.
- Schwander, J., Stauffer, B., and Sigg, A.: Air mixing in firn and the age of the air at pore close-off, *Ann. Glaciol.*, 10, 141–145, 1988.
- Sherjal, I. and Fily, M.: Temporal variations of microwave brightness temperatures over Antarctica, *Ann. Glaciol.*, 20, 19–25, 1994.
- Shimizu, H.: Air permeability of deposited snow. Institute of low temperature science: Sapporo, Japan. Contribution N° 1053. English translation, 1970.
- Shuman, C. A., Bromwich, D. H., Kipfstuhl, J., and Schwager, M.: Multiyear accumulation and temperature history near the North Greenland Ice Core Project site, north central Greenland, *J. Geophys. Res.*, 106D, 33 935–33 950, 2001.
- Simpson, W. R., King, M. D., Beine, H. J., Honrath, R. E., and Zhou, X.: Radiation-transfer modeling of snow pack photochemical processes during ALERT2000, *Atmos. Environ.*, 36, 2663–2670, 2002a.
- Simpson, W. R., King, M. D., Beine, H. J., Honrath, R. E., and Zhou, X.: Radiation-transfer modeling of snow pack photochemical processes during ALERT2000., *Atmos. Environ.*, 36, 2663–2670, 2002b.
- Simpson, W. R., King, M. D., Beine, H. J., Honrath, R. E., and Peterson, M. C.: Corrigendum to ‘Atmospheric photolysis rates during the Polar Sunrise Experiment ALERT2000’ (*Atmospheric Environment* 36, 2471–2480, 2002), *Atmos. Environ.*, 36, 5749, 2002.
- Sokolov, O. and Abbatt, J. P. D.: Adsorption to ice of n-alcohols (ethanol to 1-hexanol), acetic acid, and hexanal, *J. Phys. Chem. A*, 106, 775–782, 2002.
- Sommerfeld, R. A. and LaChapelle, E.: The classification of snow metamorphism, *J. Glaciol.*, 9, 3–17, 1970.
- Spicer, C. W., Plastridge, R. A., Foster, K. L., Finlayson-Pitts, B. J., Bottenheim, J. W., Grannas, A. M., and Shepson P. B.: Molecular halogens before and during ozone depletion events in the Arctic at polar sunrise: concentrations and sources, *Atmos. Environ*, 36, 2721–2731, 2002.
- Stamnes, K., Tsay, S. C., Wiscombe, W., and Jayaweera K.: Numerically stable algorithm for

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

discrete-ordinate method radiative transfer in multiple scattering and emitting layered media, *Appl. Opt.*, 27, 2502–2510, 1988.

Steffen, A., Schroeder, W., Bottenheim, J., Narayan, J., and Fuentes J. D.: Atmospheric mercury concentrations: measurements and profiles near snow and ice surfaces in the Canadian Arctic during Alert 2000, *Atmos. Environ.*, 36, 2653–2661, 2002.

Stogryn, A.: A study of the microwave brightness temperature of snow from the point of view of strong fluctuation theory, *IEEE Trans. Geosci. Remote Sensing*, GE-24, 220–231, 1986.

Sturm, M. and Johnson J.: Natural convection in the subarctic snow cover. *J. Geophys. Res.*, 96, 11 657–11 671, 1991.

Sturm, M. and Johnson, J.: Thermal conductivity measurements of depth hoar, *J. Geophys. Res.*, 97, 2129–2139, 1992.

Sturm, M., Holmgren, J., König M., and Morris. K.: The thermal conductivity of snow, *J. Glaciol.*, 43(143), 26–41, 1997.

Sturm, M. and Benson, C. S.: Vapor transport, grain growth and depth hoar development in the subarctic snow, *J. Glaciol.*, 43, 42–59, 1997.

Sumner, A. L. and Shepson, P. B.: Snowpack production of formaldehyde and its effect on the Arctic troposphere, *Nature*, 398, 230–233, 1999.

Surdyk, S. and Fily, M.: Results of a stratified snow emissivity model based on the wave approach: application to the Antarctic Ice Sheet, *J. Geophys. Res.*, 100C, 8837–8848, 1994.

Surdyk S. and Fily, Comparison of microwave spectral signature of the Antarctic Ice Sheet with traverse ground data, *Ann. Glaciol.*, 17, 161–166, 1993.

Surdyk, S.: Using microwave brightness temperature to detect short-term surface air temperature changes in Antarctica: An analytical approach, *Rem. Sens. Env.*, 80, 256–271, 2002.

Swanson, A. L., Blake, N. J., Blake, D. R., Rowland, F. S., and Dibb, J. E.: Photochemically induced production of CH₃Br, CH₃I, C₂H₅I, ethene and propene within surface snow, *Atmos. Environ.*, 36, 2671–2682, 2002.

Taillandier, A.-S., Domine, F., Simpson, W. R., Sturm, M., Douglas, T. A., and Severin, K.: Evolution of the Snow Area Index (SAI) of the subarctic snowpack in Central Alaska over a whole season. Consequences for the air to snow transfer of pollutants, *Environ. Sci. Technol.*, 40, 7521–7527, doi:10.1021/es060842j, 2006

Taillandier, A.-S., Domine, F., Simpson, W. R., Sturm, M., and Douglas, T. A.: The rate of decrease of the specific surface area of dry snow: isothermal versus temperature gradient conditions, *J. Geophys. Res.*, in press, 2007.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Tanikawa, T., Aoki, T., and Nishio, F.: Remote sensing of snow grain-size and impurities from Airborne Multispectral Scanner data using a snow bidirectional reflectance distribution function model, *Ann. Glaciol.*, 34, 74–80, 2002.
- 5 Tedesco, M., Pampaloni, P., Pulliainen, J., and Hallikainen, M.: Classification and retrieval of dry snow parameters by means of SMM/I data and artificial neural networks, *Proc. Geoscience and Remote Sensing Symposium*, 1, 330–332, 2003.
- Tedesco, M. and Kokhanovsky, A. A.: The semi-analytical snow retrieval algorithm and its application to MODIS data, *Rem. Sens. Env.*, in press, 2007.
- 10 Thibert, E. and Dominé, F.: Thermodynamics and kinetics of the solid solution of HCl in ice, *J. Phys. Chem. B*, 101, 3554–3565, 1997.
- Thibert, E. and Dominé, F.: Thermodynamics and kinetics of the solid solution of HNO₃ in ice, *J. Phys. Chem. B*, 102, 4432–4439, 1998.
- Thomas, G. E. and Stamnes, K.: Shortwave Radiative Transfer, in *Radiative Transfer in the Atmosphere and Ocean*, edited, 335–383, Cambridge University Press, Cambridge, 2002.
- 15 Torinesi O., Fily, M., and Genthon, C.: Interannual variability and trend of the Antarctic summer melting period from 20 years of spaceborne microwave data, *J. Climate*, 16, 1047–1060, 2003.
- Tsang, L., Chen, Z., Oh, S., Marks, R. J., and Chang, A. T. C.: Inversion of Snow Parameters from Passive Microwave Remote Sensing Measurements by a Neural Network Trained with a Multiple Scattering Model, *IEEE Trans. Geoscience Rem. Sens.*, 30, 1015–1024, 1992.
- 20 Tsang, L., Chen, C., Chang, A. T. C., Guo, J., and Ding, K.: Dense media radiative transfer theory based on quasicrystalline approximation with applications to passive microwave remote sensing of snow, *Radio Sci.*, 35, 731–749, 2000.
- Ulaby, F. T., Moore, R. K., and Fung, A. K.: *Microwave remote sensing: active and passive. Volume 1. Microwave remote sensing fundamentals and radiometry*, Norwood, M.A. Artech House, 1981.
- 25 Underwood, E. E.: *Quantitative stereology*, Addison-Wesley, Reading, MA, 1970.
- Vermote, E. and Vermeulen, A.: Atmospheric correction algorithm: Spectral reflectances (MOD09), ATBD version 4.0 available at http://eosps0.gsfc.nasa.gov/ftp_ATBD/REVIEW/MODIS/ATBD-MOD-08/atbd-mod-08.pdf, 1999.
- 30 Waddington, E. D., Cunningham, J., and Harder, S.: The effects of snow ventilation on chemical concentrations. In *Chemical Exchange Between the Atmosphere and Polar Snow*, edited by: Wolff, E. W. and Bales, R. C., Springer, 403–451, 1996.

Snow physics as relevant to snow photochemistryF. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Warren, S. G.: Optical properties of snow, *Rev. Geophys.*, 20, 67–89, 1982.
- Wen, B. Tsang, L., Winebrenner, D. P., and Ishimaru, A.: Dense medium radiative transfer theory: comparison with experiment and application to microwave remote sensing and polarimetry, *IEEE Trans Geoscience and Rem. Sens.*, 28, 1 46–59, 1990.
- 5 Wergin, W. P., Rango A., and Erbe, E. F.: Observations of Snow Crystals Using Low-Temperature Scanning Electron Microscopy, *Scanning*, 17, 41–49, 1995.
- Wergin, W. P., Rango A., Erbe, E. F., and Murphy, C. A.: Low Temperature S.E.M. of Precipitated and Metamorphosed Snow Crystals Collected and Transported from Remote Sites, *J. Microsc. Soc. Am.*, 2, 99–112, 1996.
- 10 Wettlaufer, J. S.: Impurity effects in the premelting of ice, *Phys. Rev. Lett.* 82, 2516–2519, 1999.
- Wiesmann A. and Mätzler C.: Microwave emission model of layered snowpacks, *Remote Sens. Env.*, 70, 307-316, 1999.
- Wiesmann A., Fierz, C., and Matzler, C.: Simulation of microwave emission from physically modeled snowpacks, *Ann. Glaciol.*, 31, 397–405, 2000.
- 15 Wilson, L., Tsang, L., Hwang, J.-N., and Chen, C.-T.: Mapping snow water equivalent in mountainous areas by combining a spatially distributed snow hydrology model with passive microwave remote sensing data, *IEEE Trans. Geosci. Remote Sensing*, 37, 690–704, 1999.
- Wilson, L. L., Tsang, L., and Hwang, J.-N.: Mapping SWE by combining a spatially distributed snow hydrology model with passive microwave remote-sensing data, *IEEE TGARS*, 37, 690–704, 1999.
- 20 Winebrenner, D. P., Arthern, R. J., and Shuman, C. A.: Mapping Greenland accumulation rates using observations of thermal emission at 4.5-cm wavelength, *J. Geophys. Res.*, 106D, 33 919–33 934, 2001.
- 25 Wiscombe, W. J.: The delta-Eddington approximation for a vertically inhomogeneous atmosphere, *NCAR Technical Note*, NCAR/TN-121+STR, 1-66, 1977.
- Wiscombe, W. and Warren, S.: A model for the spectral albedo of snow. 1: pure snow, *J. Atmos. Sci.*, 37, 2712–2733, 1980.
- Wismann, V.: Monitoring of seasonal snowmelt on Greenland with ERS scatterometer data, *IEEE Trans. Geosci. Remote Sensing*, 38, 1821–1826, 2000.
- 30 Xie, Y., Yang, P., Gao, B.-C., Kattawar, G. W., and Mishchenko M. I.: Effect of ice crystal shape and effective size on snow bidirectional reflectance, *J. Quantitative Spectr. Radiative Transfer*, 100, 457–469, 2006.

Snow physics as relevant to snow photochemistryF. Domine et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Zappa, M., Pos, F., Strasser, U., Warmerdam, P., and Gurtz, J.: Seasonal water balance of an Alpine catchment as evaluated by different methods for spatially distributed snowmelt modelling, *Nordic Hydrol.*, 34, 179–202, 2003.

5 Zege, E. P., Ivanov A. P., and Katsev I. L.: Image transfer through a scattering medium, Berlin: Springer, 1991.

Zhou, X., Beine, H. J., Honrath, R. E., Fuentes, J. D., Simpson, W., and Shepson, P. B.: Snowpack photochemical production of HONO: a major source of OH in the Arctic boundary layer in spring time, *Geophys. Res. Lett.*, 28, 4087–4090, 2001.

10 Zurk, L. M., Tsang, L., and Winebrenner, D. P.: Scattering properties of dense media from Monte Carlo simulations with application to active remote sensing of snow, *Radio Sci.*, 31, 4, 803–820, 1996.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.

Table 1. Typical ranges of values of selected physical variables for the most frequent snow crystal types in surface snow (top 50 cm). Data from Sturm et al. (1997), Domine et al. (2002), Albert et al. (2002), Simpson et al. (2002), Legagneux et al. (2002).

Physical variable	Fresh snow	Decomposing particles	Rounded grains, ET metamorphism	Rounded grains, wind-pack	Faceted crystals	Depth hoar	Melt-freeze crusts
Density, g cm^{-3}	0.01–0.2	0.06–0.25	0.15–0.35	0.3–0.6	0.13–0.3	0.15–0.28	0.15–0.4
Specific surface area, $\text{cm}^2 \text{g}^{-1}$	350–1600	150–900	110–300	150–380	80–450	70–220	20–50
e-folding depth, cm	3–10	2–10	5–15	3–10	10–25	15–30	20–40
Permeability, 10^{-10}m^2	10–50	10–90	10–80	5–30	50–200	100–600	1–20
Heat conductivity, $\text{W m}^{-1} \text{K}^{-1}$	0.03–0.12	0.06–0.2	0.08–0.3	0.2–0.6	0.05–0.12	0.03–0.15	0.1–0.5

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.

Table 2. Typical values of selected physical variables encountered in the top 20 cm (photic zone) of three surface snow types in late winter. Data from Sturm and Benson (1997), Sturm et al. (1997), Simpson et al. (2002 and unpublished results), Albert et al. (2002), Taillandier et al. (2006), Domine et al. (unpublished results).

Physical variable	Tundra snowpack	Taiga snowpack	Alpine snowpack
Density, g cm^{-3}	0.4	0.18	0.2
Specific surface area, $\text{cm}^2 \text{g}^{-1}$	240	190	380
e-folding depth, cm	5	15	8
Permeability, 10^{-10}m^2	8	200	50
Heat conductivity, $\text{W m}^{-1} \text{K}^{-1}$	0.4	0.05	0.2

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.




Fig. 1. Schematic illustrating the physical variables that determine the snow-to-atmosphere flux of NO_2 due to the photolysis of adsorbed NO_3^- . These variables include light e -folding depth, snow specific surface area, density, permeability and diffusivity (see text). Figure in part taken from Domine and Shepson (2002).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Snow physics as
relevant to snow
photochemistry

F. Domine et al.




Fig. 2. Photomicrographs of metamorphic crystals frequently observed in snowpacks. **(a)** Depth hoar crystal; **(b)** faceted crystals; **(c)** dense melt-freeze crust; **(d)** melt-freeze polycrystal in low density (0.11 g cm^{-3}) melt-freeze layer; **(e)** medium-size rounded grains in a near-basal layer subjected to near-isothermal metamorphism, of density 0.35 g cm^{-3} ; **(f)** crystals with mixed forms, showing facets and rounded shapes.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 3. Simplified stratigraphies illustrating the main seasonal snowpacks: the tundra snowpack near Barrow, Alaska, on 4 April 2004 (Domine, Taillandier and Simpson, unpublished results); the taiga snowpack near Fairbanks, Alaska, on 12 February 2004 (after Taillandier et al., 2006); the Alpine snowpack in the ablation zone of the Argentières glacier, near Chamonix, French Alps, on 31 January 2002 (Domine and Lauzier, unpublished results).

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as
relevant to snow
photochemistry

F. Domine et al.




Fig. 4. Measured and computed (with SNOWPACK9.0) snow stratification at Weissfluhjoch, Davos, during the winter 92/93. The upper envelope represents the measured, total snow height. The solid lines show the heights above ground of 12 markers placed on the snow surface at different times of the winter and subsequently buried by precipitation. The dotted lines show the computed heights of the corresponding layer.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Snow physics as relevant to snow photochemistry

F. Domine et al.




Fig. 5. SSA-density correlation, using the 297 coupled SSA-density measurements of Domine et al. (2007). SSA was measured using CH₄ adsorption at 77 K. The graph also shows average values and standard deviations of some of the snow types mentioned in Table 1: Fresh snow, decomposing particles, rounded grains (encompassing both ET metamorphism grains and windpacked grains), faceted crystals and depth hoar (after Domine et al., 2007).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Snow physics as relevant to snow photochemistry

F. Domine et al.




Fig. 6. A simulation of optical transmission of snow. In this simulation, the ratio of the in-snow actinic flux, F , to the downwelling incident flux, F_0 , is shown as a function of depth in the snow. The depth in the snow is listed as the optical depth, τ_{ext} , which is proportional to the physical depth in the snowpack. The relationship to physical depth depends upon the equivalent spherical radius of the snow (related to SSA) and density, as described in the text. For the example described in the text (aged snow, density ($\sim 0.3 \text{ g/cm}^3$), and equivalent spherical grain radius of $\sim 0.5 \text{ mm}$), the extinction coefficient is $\sim 1 \text{ mm}^{-1}$, and in this case, the depth would be equivalent to millimeters within the snowpack. The asymmetry factor, g , is 0.886. The single scattering albedo of the snow is $\omega = 0.99995$, indicating that the absorption coefficient is 0.00005 times smaller than the total extinction, which is typical of snow in the UV. Four cases of incident light are shown. Cases **(a)**–**(c)** have purely direct sunlight at three solar zenith angles (a) $\theta_0 = 0^\circ$, (b) $\theta_0 = 60^\circ$, (c) $\theta_0 = 84^\circ$, while case **(d)** is for purely diffuse incoming radiation. One observes non-exponential behavior in the near surface region, generally with enhancement of photolysis rates in the snowpack ($F/F_0 > 1$ indicates light enhancement). Deeper in the snowpack, in the asymptotic region, the light is completely diffused and decays exponentially, with an e-folding depth of approximately 20 cm (based upon the example in the text). Reprinted from Simpson et al. (2002). The triangles overlapping the curve for the 0.19 mm grain size are measurements by Hudson et al. (2006), showing the excellent agreement between theory and experiments in the visible and near IR. In the mid IR, the greater absorption limits the validity of the theory.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.




Fig. 7. Calculated spherical albedo of snow with different grain sizes. The triangles overlapping the curve for the 0.19 mm grain size are measurements by Hudson et al. (2006), showing the excellent agreement between theory and experiments in the visible and near IR. In the mid IR, the greater absorption limits the validity of the theory.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 8. The dependence of the diffuse reflectance at 400 nm on the soot concentration (in ppm by volume, i.e. the volume fraction of the snow occupied by soot particles). The solar zenith angle is 60° . Snow is composed of non-absorbing grains with the effective grain diameter 200 or $500 \mu\text{m}$, the snow density is 0.275 g cm^{-3} in both cases, the asymmetry factor g is 0.75. Soot particles are treated as Rayleigh scatterers with the refractive index equal to $1.8-0.5i$.

Snow physics as relevant to snow photochemistry

F. Domine et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as relevant to snow photochemistry

F. Domine et al.




Fig. 9. The main processes responsible for heat transport through snow. The snow structure represented is columnar depth hoar. Processes shown are conduction through interconnected snow crystals (the heat conductivity of ice $k_{\text{ice}} = 2.3 \text{ W m}^{-1} \text{ K}^{-1}$, is recalled), conduction by air in the pore space (the heat conductivity of air is $k_{\text{air}} = 0.023 \text{ W m}^{-1} \text{ K}^{-1}$), latent heat transport by sublimation/condensation cycles induced by the temperature gradient. These three processes are measured as k_{eff} , for example using a heated needle probe. Additionally, air flow through snow can be induced by surface wind or air convection and this represents an extra process which, when present, can transport heat much faster than either conduction through the ice or void space.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Snow physics as
relevant to snow
photochemistry

F. Domine et al.




Fig. 10. Impact of the value of K_{ia} on the trace gas diffusion in snow, calculated after Herbert et al. (2006), for a snow density of 0.34 g cm^{-3} ($r_{sp}=0.2 \text{ g cm}^{-3}$) and a SSA of $200 \text{ cm}^2 \text{ g}^{-1}$. The factor f is the ratio of the effective diffusion coefficient of an adsorbing trace gas in snow over the diffusion coefficient in snow in the absence of adsorption (see text).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)