

**Modelling Pliocene
climate using the
Hadley Centre Model**

F. J. Bragg et al.

**Mid-Pliocene climate modelled using the
UK Hadley Centre Model: PlioMIP
Experiments 1 and 2**

F. J. Bragg¹, D. J. Lunt¹, and A. M. Haywood²

¹BRIDGE, School of Geographical Sciences, University of Bristol, University Road, Bristol, BS8 1SS, UK

²School of Earth and Environment, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT, UK

Received: 13 March 2012 – Accepted: 27 March 2012 – Published: 18 April 2012

Correspondence to: F. J. Bragg (fran.bragg@bristol.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

The Pliocene Model Intercomparison Project (PlioMIP) project is a sub-project of the Paleoclimate Modelling Intercomparison Project (PMIP) whose objective is to compare predictions of the mid-Pliocene climate from the widest possible range of general circulation models. The mid-Pliocene (3.3–3.0 Ma) is the most recent sustained period of greater warmth and atmospheric carbon dioxide concentration than the pre-industrial times and as such has potential to inform predictions of our warming climate in the coming century. This paper describes the UK contribution to PlioMIP using the Hadley Centre Model both in atmosphere-only mode (HadAM3, PlioMIP Experiment 1) and atmosphere-ocean coupled mode (HadCM3, PlioMIP Experiment 2). The coupled model predicts a greater overall warming (3.3°C) relative to the control than the atmosphere-only (2.5°C). The Northern Hemisphere latitudinal temperature gradient is greater in the coupled model with a warmer equator and colder Arctic than the atmosphere-only model, which is constrained by sea surface temperatures from Pliocene proxy reconstructions. The atmosphere-only model predicts a reduction in equatorial precipitation and south Asian monsoon intensity whereas the coupled models shows and increase in the intensity of these systems. Sensitivity studies using alternative boundary conditions for both the Pliocene and the control simulations are presented, which indicate the sensitivity of the mid-Pliocene warming to uncertainties in both pre-industrial and mid-Pliocene climate.

1 Introduction

This paper describes the Pliocene Model Intercomparison Project (PlioMIP) simulations carried out using the UK Meteorological Office Hadley Centre Model (HadAM3 and HadCM3).

General circulation models (GCMs) are one of the main tools used for studying the climate system of the present day and the past and to predict likely climate changes

GMDD

5, 837–871, 2012

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Modelling Pliocene
climate using the
Hadley Centre Model**

F. J. Bragg et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

that can be expected in the future. These models vary significantly in the way they parameterise certain complex processes and as such vary in their simulation of the modern climate as well as in their past and future predictions. The Paleoclimate Modelling Project (PMIP, see <http://pmip3.lsce.ipsl.fr/>) exists to create intercomparisons of the widest possible range of models forced as nearly as possible with identical palaeo boundary conditions. The intercomparisons focus on different time slices of particular interest to the scientific community, including the Last Glacial Maximum (21 kyr), Last Interglacial (130 kyr, 125 kyr and 115 kyr) and mid-Holocene (6 kyr) and more recently extended to 8.2 kyr and Pliocene, specifically the mid-Pliocene warm period (MPWP) from 3.29–2.97 Ma (Haywood et al., 2010).

The Pliocene is of particular interest for the development of the GCMs used in future climate predictions as it is the most recent sustained period which is significantly warmer than the present day and thus the climate system may operate in a similar manner to potential climates of the coming century. A substantial dataset describing the MPWP time slab has been assembled by the US Geological Survey PRISM project (Pliocene Research, Interpretation and Synoptic Mapping, <http://geology.er.usgs.gov/eespteam/prism/>) including topography data, sea surface temperatures, vegetation reconstruction and ice sheet extents and topography (Dowsett et al., 2010). These data are used to force the models to achieve consistent simulations of how the Pliocene climate differs from the pre-industrial control.

This paper describes the implementation of the PRISM3 boundary conditions for use with the UK Hadley Centre Model according to the PlioMIP protocols for atmosphere-only models (“Experiment 1”, Haywood et al., 2010) and coupled ocean-atmosphere models (“Experiment 2”, Haywood et al., 2011), a summary of the basic results and some initial analysis including model-data comparison.

2 Model description

We use the UK Meteorological Office Hadley Centre Model, also known as the Unified Model (UM) for these experiments, specifically the atmosphere-only version, HadAM3 (Pope et al., 2000), for Experiment 1 and the coupled ocean-atmosphere version, HadCM3 (Gordon et al., 2000), for Experiment 2.

2.1 Atmosphere

The atmosphere module has a resolution of 96×73 grid points ($3.75^\circ \times 2.5^\circ$) and 19 vertical levels using a hybrid σ -pressure grid with a timestep of 0.5 h. An Arakawa B-grid (Arakawa and Lamb, 1977) is used in the horizontal plane to improve accuracy, with thermodynamic variables stored at the centre of the grid and wind components at the corners (Johns et al., 1997).

The atmosphere-only model uses specifies 12 mid-monthly fields of sea-surface temperature as boundary conditions, which are interpolated to daily values at run-time.

A convection scheme due to Gregory et al. (1997) is included which accounts for the direct effects of convection on momentum. A first order scheme for turbulent vertical mixing of momentum and thermodynamic quantities is used within the boundary layer, which can occupy up to the first 5 layers of the model (Smith, 1990). Sub grid-scale gravity wave and orographic drag parameterisations include the impact of orographic variance anisotropy (Milton and Wilson, 1996; Gregory et al., 1998).

Clouds are modelled as either water, ice or mixed-phase between 0 and -9°C . Clouds are aggregated into 3 layers (low, medium and high) and form when the cell moisture level standard deviation exceeds a critical level of relative humidity, $\text{RH}_{\text{crit}} = 0.7$. The threshold of total water content for precipitation to occur is varied between land and ocean cells to account for the different levels of available cloud condensation nuclei.

GMDD

5, 837–871, 2012

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



2.2 Radiation and land-surface energy schemes

The radiation scheme of Edwards and Slingo (1996) has 6 short-wave and 8 long-wave bands and represents the effects of water vapour, carbon dioxide, ozone and minor trace gases. A background aerosol climatology following Cusack et al. (1998) increases the atmospheric absorption of short-wave radiation relative to previous versions representing a significant improvement. The land-surface energy scheme, MOSES I (Cox et al., 1999) accounts for the effects of freezing and melting of soil moisture in 4 soil layers and includes the impact of atmospheric concentration of carbon dioxide, water vapour and temperature on stomatal resistance to evapo-transpiration.

2.3 Ocean

The ocean module has a horizontal resolution of 288×144 grid points ($1.25^\circ \times 1.25^\circ$): that is 6 ocean cells correspond to each atmosphere cell. The land-sea mask is defined at the atmosphere resolution to simplify coupling. There are 20 vertical levels with finer definition at the ocean surface: the first cell is 10 m deep. The ocean timestep is 1 h. The ocean and atmosphere modules are coupled once a day with no flux adjustment being necessary.

Modern bathymetry is derived from the ETOPO5 reconstruction (Edwards, 1989) using a simple smoothing algorithm. Behaviour in some significant channels is modified from the resulting coarse interpolation to ensure a more realistic model performance (Gordon et al., 2000). The Greenland-Scotland ridge and Denmark Strait have significant sub-gridscale channels which are lost in the smoothing and so have been re-created by deepening single cell width channels in 3 locations along the ridge to reproduce a mean outflow matching observations. The resolution of the Gibraltar Strait leaves the Mediterranean isolated, so a partial mixing of the closest cells at each depth down to 1200 m is carried out to represent the actual mixing that occurs across this opening. The region around Indonesia is modified to ensure that flow occurs between

GMDD

5, 837–871, 2012

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Indonesia and Papua New Guinea and not between Indonesia and the mainland of Asia.

A rigid lid approach is used meaning there is no variation in volume of the ocean. Freshwater flux from land runoff is therefore converted to a salinity flux on entering the ocean. Ice sheets are not modelled dynamically in HadCM3, therefore the snow accumulation on each ice sheet is balanced by a notional equivalent loss through iceberg calving represented as a freshwater flux distributed around the edge of the ice sheet and polar oceans.

The ocean mixed layer mixing of tracers (potential temperature and salinity) is represented by the Kraus and Turner (1967) model which assigns 15% of gravitational potential energy and 70% of wind-stress energy to turbulent kinetic energy, which is mixed out exponentially with depth. At all depths, 5 iterations of convective mixing of tracers are carried out each timestep. Horizontal mixing of tracers is carried out using the isopycnal parameterisation of Gent and McWilliams (1990). Horizontal mixing of momentum is performed using a latitudinally varying formulation which, coupled with the finer resolution of the ocean grid, enables western boundary currents to be resolved.

2.4 Sea ice

Sea ice is calculated as a zero layer model on top of the ocean grid. Partial cell coverage of sea ice is possible in all high latitude cells, up to 0.995 in the Arctic and 0.98 in the Antarctic according to the parameterisation of sea ice concentration due to Hibler (1979). Ice forms primarily by freezing in leads, also ice can form from snow falling on existing ice and by freezing at the base at the freezing point of -1.8°C . A constant salinity is assumed for ice, excess salt from freezing is rejected into the ocean. Ice drift follows the ocean currents in the top layer, converging ice has a limit of 4 m in depth. Ice albedo is set at 0.8 below -10°C and 0.5 above 0°C with a linear variation between.

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.5 Model validation

The Hadley Centre model validation is documented for HadAM3 in Pope et al. (2000) and for HadCM3 in Gordon et al. (2000). The model has been shown to reproduce the main features of modern climate observations.

3 Experimental design

Table 1 summarises the experimental design for both Experiment 1 (Haywood et al., 2010) and Experiment 2 (Haywood et al., 2011) Pliocene and control simulations. Also included are details for an additional Pliocene simulation, a fully coupled experiment based on the previous PRISM2 version of boundary conditions (Dowsett, 2007) which will be referred to in the discussion below.

3.1 Land-sea mask

The PlioMIP protocols define two possible land-sea masks: a “preferred” mask which differs from modern (primarily due to sea-level change and glacial erosion) and an “alternate” mask which is the same as modern. For Experiment 1, the Pliocene simulation land-sea mask was derived from the PlioMIP preferred fractional dataset, being first interpolated onto the UM $3.75^\circ \times 2.5^\circ$ grid, then those cells with a land-fraction greater than 0.5 being set to be land. The principal differences from the modern mask are in the Hudson Bay which is filled in at low altitude and the regions of West Antarctica where the modern ice shelf is absent in the Pliocene. The Panama Seaway is post-edited to be closed as the interpolation process renders this region as ocean in the coarser grid. The standard UM land-sea mask was used for the control run. All of the coupled Experiment 2 simulations also use the standard UM land-sea mask without modification due to the difficulty of changing the land-sea mask in the ocean module and achieving a stable solution. This mask is derived from the US Navy 10' topography (NCAR/Navy, 1984) and can be seen in the boundary condition plots in Fig. 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Topography including ice sheets

For both experiments, Pliocene topography was created using the anomaly method specified in the PlioMIP protocol. Anomalies were calculated as the difference between the PlioMIP Pliocene and control datasets. For Experiment 1, the UM standard topography was then extended to cover the different land-sea mask using an iterative expansion algorithm before applying the anomaly values. Ice sheet topography is included simply as part of the overall topography. The resultant field was then masked with the required final land-sea mask ensuring that all ocean points have an elevation of 0 m and no land points are below sea-level. For the pre-industrial control dataset, calculation of derived fields required for the model describing the orographic variance was performed on the high resolution topography data prior to regridding onto the UM resolution. These data were also used for the Pliocene in both experiments as topographic data on the same resolution as the modern data are not available to make the equivalent variance calculations. In the case of Experiment 1, the data were expanded to match the preferred land-sea mask.

3.3 Land surface properties including ice sheet extent

Properties required by the land surface-energy scheme for the Pliocene simulations were derived from the PlioMIP BIOME4 dataset by means of the lookup table of Hayward et al. (2010). This table relates the 28 biome types describing land cover to the MOSES I input parameters (Cox et al., 1999) via the land-use classifications of Wilson and Henderson-Sellers (1985). These data include ice as a possible surface type, hence the extent of ice sheets is naturally incorporated into this process. In land regions which are specified as ocean in the PlioMIP dataset, these derived parameter fields were expanded out from neighbouring land points, most notably in the West Antarctic region. This conversion is illustrated using snow-free albedo which is shown in Fig. 1.

GMDD

5, 837–871, 2012

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



For the control experiments, the standard UM dataset of vegetation was applied which is based on the Wilson and Henderson-Sellers (1985) archive of land cover.

3.4 Soil properties

Soil properties in both Pliocene and control experiments are those used in the standard UM setup derived from Wilson and Henderson-Sellers (1985), expanded in the case of Experiment 1 Pliocene simulation to match the preferred land-sea mask. In reality, soil properties could be different at the Pliocene compared to modern, but a lack of Pliocene palaeosol data precludes meaningful changes to be implemented in the model.

3.5 River routing (Experiment 2 only)

River routing in both Pliocene and control experiments is the same as the UM standard catchments.

3.6 Sea surface temperatures and sea ice (Experiment 1 only)

Sea surface temperatures (SST) for the Pliocene simulation were created using the specified PlioMIP anomaly method (Haywood et al., 2010, Sect. 2.2). The difference between the Pliocene and modern fields was calculated and applied to the UM standard SST climatology for each month. Temperatures were not allowed to fall below -1.7°C . The data were expanded into regions which are land in the standard UM mask using an iterative process, in particular into the Hudson Bay. The SST fields for January and July are shown in Fig. 2.

Sea ice is specified on a monthly basis where $\text{SST} < -1.7^{\circ}\text{C}$ with a depth of 2 m in the Northern Hemisphere and 1 m in the Southern Hemisphere, as in the control.

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.7 Solar constant and orbital parameters

The solar constant and orbital parameters were unchanged from the standard UM values (Ingram et al., 1997) and are summarised in Table 1. The perihelion is specified as day 2.5 from a 360 day year giving an angle of 281.5° from the spring equinox.

3.8 Trace gases and aerosols

The atmospheric CO_2 concentration was set to 280 ppmv for the control simulations and 405 ppmv for the Pliocene in accordance with the PlioMIP protocol. Levels of methane (761 ppb) and nitrous oxide (269 ppb) were held constant in all simulations at standard pre-industrial levels. Ozone is specified as a fixed annual climatology of 12 monthly fields which is the same for all simulations. No aerosols are specified other than the background climatology of the scheme of Cusack et al. (1998).

3.9 Initialisation

For Experiment 1 simulations, the atmosphere was initialised using a standard UM pre-industrial restart file. For Experiment 2, the atmosphere and ocean modules were initialised from the respective Pliocene and control simulations based on PRISM2 boundary conditions (Lunt et al., 2010) with cumulative integration lengths of over 1000 yr. Time series of temperature evolutions of the global mean of 1.5 m air temperature are shown in Fig. 3 along with intermediate and deep ocean temperature evolution for the coupled simulations.

Other parameters were initialised as follows:

- for coupled experiments, sea ice is initialised using the method of Sect. 3.6.
- soil temperature is initialised with a constant value of 14°C in ice free regions and -6°C with ice cover.
- soil moisture is initialised with constant default values for each of the 4 layers.

GMDD

5, 837–871, 2012

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



– snow depth is initialised with flat profile of 50 m over ice and none elsewhere.

3.10 Integration length, spinup and climatological means

Both Experiment 1 simulations were run for 50 yr, as required by the PlioMIP protocol, enabling the simulations to come to equilibrium. The Experiment 2 Pliocene simulation was run for 500 yr as specified in the protocol. The control model was run for 200 yr, but this simulation is a continuation from that of Lunt et al. (2010) with no change to the boundary conditions.

Average climatologies for the Experiment 1 simulations were calculated over the last 30 yr and for Experiment 2 over the final 50 yr.

4 Results

The results are presented here firstly for Experiment 1, then Experiment 2.

4.1 Global

Global means for the Experiment 1 and 2 simulations are listed in Table 2. The coupled model predicts a significantly greater level of warming in the Pliocene relative to the control than the atmosphere-only version: 3.3°C compared with 2.5°C. There is also a small rise in total precipitation in the Pliocene relative to the control of around 4 % in Experiment 1 and 5.5 % in Experiment 2.

4.2 Experiment 1

Figure 4 shows the zonal average surface air temperatures for Experiment 1, for land only and ocean only, both absolute values and Pliocene minus control difference. The profile over the non-polar oceans falls to close to 0°C around the equator; this profile is strongly constrained by the imposed sea surface temperature boundary condition.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The latitudinal temperature gradient, especially in the Northern Hemisphere is significantly reduced in the Pliocene relative to the control. Polar amplification in the Pliocene relative to the control is clear in the zonal profiles.

These results are apparent in greater detail in Fig. 5 which shows annual, DJF and JJA mean surface air temperature patterns for both Pliocene and control simulations and the difference between them. The polar amplification is most pronounced in winter for both hemispheres. In the Antarctic, the majority of the warming maxima occur in regions which are ocean in the Pliocene and land in the control simulation due to the change in albedo and heat capacity, combined with smaller areas on land where the ice sheet is at considerably lower altitude in the Pliocene model. Pliocene Arctic warming is also associated with regions of change in the altitude or extent of the Greenland icesheet in a similar manner to the Antarctic. There is also a warming evident in the North Atlantic driven by the sea surface temperature boundary conditions which are significantly warmer than in the control.

Figure 6 summarises the precipitation patterns for the Pliocene and control experiments and the difference between them, showing the annual, DJF and JJA means. There is a reduction in equatorial rainfall in the Pliocene, especially the extent and intensity of the south Asian summer monsoon systems.

4.3 Experiment 2

Figure 7 shows the zonal mean surface air temperatures globally, for land only and ocean only along with Pliocene minus control difference for Experiment 2. Polar amplification is seen as in Experiment 1, though with reduced magnitude. Experiment 2 also exhibits noticeable warming at lower latitudes, typically at levels of around 2 °C over water and 4 °C on land which are consistent throughout most of the non-polar latitudes. The difference in Pliocene-control changes are shown in Fig. 11 as Experiment 2 minus Experiment 1. Figure 11a–c again demonstrate the increased warming at lower latitudes in Experiment 2 and the reduction in polar warming, especially apparent in the Northern Hemisphere. Taken together, the warmer equator and colder (less warm)

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



poles lead to an increased latitudinal temperature gradient, which is broadly similar in shape to that predicted for the control experiment (see Fig. 7a), in contrast to the reduction in latitudinal gradient shown by the PRISM reconstruction, implicit in Fig. 4c.

This global shift in temperature is also apparent in greater spatial detail in Fig. 8, which shows global temperature patterns for Experiment 2. Outside of the polar regions, there is very little variation in temperature shift with latitude, only the marked difference between land and sea noted above. The difference in Experiment 1 and Experiment 2 patterns of temperature change are shown in Fig. 11d–f and confirm the previous observations, the equatorial ocean is broadly warmer in Experiment 2, the poles are cooler, especially in winter and the land warms more than the oceans. The most striking difference is in the far north Atlantic where there is apparent cooling in Experiment 2, more meaningfully described as a lack of the warming seen in the sea surface temperature data which are imposed in Experiment 1.

Figure 9 shows precipitation patterns for Experiment 2. The Pliocene minus control differences for Experiment 2 are very different here from those seen for Experiment 1 in Fig. 6. In this case, there is an increase in equatorial precipitation and an intensification of the Indian monsoon. There is also a significant drying over equatorial South and Central America.

Figure 10 summarises the coupled model predicted sea surface temperatures and salinity for the Pliocene, control and the difference between them. Also shown are Atlantic (Fig. 10g,h) and Pacific (Fig. 10i,j) zonally averaged meridional overturning streamfunction. The change in sea surface temperature between the Pliocene and control broadly parallels that seen in surface air temperature over the oceans, a rise in the order of 2 °C in a fairly uniform distribution. There is a distinct change in circulation south of Greenland with adjacent warming and cooling zones. Salinity increases in the Atlantic and reduces in the rest of the world ocean, with the greatest decrease being in the Arctic ocean. The Pacific streamfunction shows a significant increase in the formation of Antarctic Bottom Water.

GMDD

5, 837–871, 2012

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Discussion

5.1 Comparison with data

The model results are compared with data in Fig. 12. The Pliocene minus control sea surface temperatures from the coupled experiments are shown in the background contours. The data are shown as the difference between the PRISM3 data points and the annual mean of the Hadley Centre HadISST compilation of sea surface temperature observations (Rayner et al., 2003) averaged over the years 1870–1899 at the PRISM3 locations. Data sites along the Eastern Pacific coast show high levels of warming over and above the general Pliocene increased warmth, phenomena which are not reproduced by the model. Warming of the Kuroshio current is present in the model. The relatively dense sampling of the North Atlantic shows consistent warming north of 50° N. This extended “hotspot” is not reproduced by the model, however there is a distinct circulation change south of Greenland evident in the sea surface temperature change. HadCM3 is known to be sensitive in the modern to the detailed submarine topography of the Greenland-Scotland Ridge in this region (Gordon et al., 2000) and recent work has also suggested that this region is sensitive to uncertainties in the Pliocene bathymetry (Robinson et al., 2011).

5.2 Impact of revised PRISM3 boundary conditions

The PRISM3 boundary conditions applied in the model are updated from the previous version, PRISM2 (Dowsett, 2007). Previous simulations has been carried out with the coupled model, using these older boundary conditions (Lunt et al., 2010) which we continued for 200 yr, Fig. 13 shows the change in Pliocene-control differences for PRISM3 compared with PRISM2 boundary conditions. See Table 1 for details of the PRISM2 model simulation. There is very little difference in the simulations in terms of PRISM3-PRISM2 global means: surface air temperature anomaly falls by 0.05 °C and precipitation is unchanged to 3 significant figures. At the regional level however, there

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



is a distinct increase in seasonality of temperature over much of the land in the Northern Hemisphere. There are also temperature differences where ice sheet topography has been updated and in regions of significant change to the orographic boundary conditions, notably the Rockies which show cooling with PRISM3 topography even in summer when the rest of the Northern Hemisphere is has a fairly uniform warming trend. Recent work has interrogated the impact of each set of boundary conditions (CO₂, orography, ice and vegetation) in the PRISM2 simulations (Lunt et al., 2012), and a similar study is required for the PRISM3 simulations in order to fully understand the changes shown in Fig. 13, but this result serves to highlight the significance of the uncertainty in boundary conditions on model predictions.

5.3 Impact of alternative vegetation data in control experiment

For the control Experiment 1 simulation, an alternative vegetation dataset was used to derive the land surface properties required by the model. PlioMIP supplies a modern biomes dataset (BAS_Observ_BIOME.nc) whose data represent potential natural vegetation created by running the BIOME4 vegetation model (Haxeltine and Prentice, 1996; Kaplan et al., 2003) forced with a pre-industrial climate. This differs from the WHS dataset which represents modern conditions and thus includes an element of human land use changes. The same lookup process as for the Pliocene data was applied to derive surface characteristics from biomes. The differences due to the change in the control model is shown in Fig. 14. As with the change in Pliocene boundary conditions discussed above, this change in the control model also results in significant changes to the results. There is a direct impact over land with the standard pre-industrial control data resulting in predicted temperatures around 2 °C colder over much of the Northern Hemisphere land in Eurasia and North America, corresponding to a reduction in global annual mean temperature of 0.26 °C or a reduction of around 10 % of the total warming in the standard experiment (0.26 in 2.5 °C). It is possible that the Pliocene minus control anomalies here are a closer representation of the true natural mid-Pliocene warming, as the vegetation changes include only naturally forced changes. Furthermore, the

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



underlying biome vegetation types and their conversion to model boundary condition parameters is more consistent between the Pliocene and control simulations than in the standard pre-industrial dataset.

6 Conclusions

- The predicted global mean temperature increase for the Pliocene is greater for the coupled model (3.3°C) than the atmosphere-only version (2.5°C).
- Polar amplification is evident in the atmosphere-only model, which is highly constrained by the PRISM3 SST dataset, reducing the mean latitudinal temperature gradient relative to the control. The coupled model has higher levels of warming in equatorial regions and lower levels in polar areas, relative to the atmosphere-only version, resulting in a higher mean latitudinal gradient similar to that in the control simulation.
- There is a marked difference in predicted precipitation patterns between the atmosphere-only and coupled models. The atmosphere-only model predicts reduced equatorial and Asian monsoon intensity whereas in the coupled model, these systems increase in intensity.
- Features of the PRISM3 data which could be interpreted as suggesting changes in ocean circulation are partially supported by the coupled model results. There is good agreement in the region of the Kuroshio current which is predicted to warm significantly, no evidence of Eastern Pacific coastal warming in present day cold upwelling zones. There is evidence of a distinct circulation change in the North Atlantic south of Greenland in the model, in the data there is also a significant feature in the North Atlantic, but considerably further north.
- Sensitivity to boundary conditions was demonstrated using alternative datasets for both Pliocene and control simulations. Whilst the change from PRISM2 to

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



PRISM3 mid-Pliocene conditions did not significantly alter the global mean climate, the change from standard pre-industrial to potential natural modern vegetation reduced the predicted Pliocene minus control warming by 10 %.

References

- 5 Arakawa, A. and Lamb, V. R.: Computational design of the basic dynamical processes of the UCLA general circulation model, in: *Methods in Computational Physics*, edited by: Chang, J., Academic Press, New York, 173–265, 1977.
- Cox, P. M., Betts, R. A., Bunton, C. B., Essery, R. L. H., Rowntree, P. R., and Smith, J.: The impact of new land surface physics on the GCM simulation of climate and climate sensitivity, *Clim. Dynam.*, 15, 183–203, 1999.
- 10 Cusack, S., Slingo, A., Edwards, J. M., and Wild, M.: The radiative impact of a simple aerosol climatology on the Hadley Centre atmospheric GCM, *Q. J. Roy. Meteor. Soc.*, 124, 2517–2526, 1998.
- Dowsett, H. J.: The PRISM palaeoclimate reconstruction and Pliocene sea-surface temperature, in: *Deep-Time Perspectives on Climate Change: Marrying the Signal from Computer Models and Biological Proxies*, edited by: Williams, M., Haywood, A. M., Gregory, F. J., and Schmidt, D. N., Geological Soc Publishing House, Brassmill Enterprise Ctr, Brassmill Lane, Bath, BA1 3JN, UK, 459–480, 2007.
- 15 Dowsett, H., Robinson, M., Haywood, A., Salzmann, U., Hill, D., Sohl, L., Chandler, M., Williams, M., Foley, K., and Stoll, D.: The PRISM3D paleoenvironmental reconstruction, *Stratigraphy*, 7, 123–139, 2010.
- Edwards, M.: *Global gridded elevation and bathymetry on 5-minute geographic grid (ETOPO5)*, NOAA, National Geophysical Data Center, Boulder, Colorado, USA, 1989.
- Edwards, J. M. and Slingo, A.: Studies with a flexible new radiation code, 1. Choosing a configuration for a large-scale model, *Q. J. Roy. Meteor. Soc.*, 122, 689–719, 1996.
- 25 Gent, P. R. and McWilliams, J. C.: Isopycnal mixing in ocean circulation models, *J. Phys. Oceanogr.*, 20, 150–155, 1990.
- Gordon, C., Cooper, C., Senior, C. A., Banks, H., Gregory, J. M., Johns, T. C., Mitchell, J. F. B., and Wood, R. A.: The simulation of SST, sea ice extents and ocean heat transports in a ver-

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sion of the Hadley Centre coupled model without flux adjustments, *Clim. Dynam.*, 16, 147–168, 2000.

Gregory, D., Kershaw, R., and Inness, P. M.: Parametrization of momentum transport by convection, 2. Tests in single-column and general circulation models, *Q. J. Roy. Meteor. Soc.*, 123, 1153–1183, 1997.

Gregory, D., Shutts, G. J., and Mitchell, J. R.: A new gravity-wave-drag scheme incorporating anisotropic orography and low-level wave breaking: impact upon the climate of the UK Meteorological Office Unified Model, *Q. J. Roy. Meteor. Soc.*, 124, 463–493, 1998.

Haxeltine, A. and Prentice, I. C.: BIOME3: An equilibrium terrestrial biosphere model based on ecophysiological constraints, resource availability, and competition among plant functional types, *Global Biogeochem. Cy.*, 10, 693–709, 1996.

Haywood, A. M., Dowsett, H. J., Otto-Bliesner, B., Chandler, M. A., Dolan, A. M., Hill, D. J., Lunt, D. J., Robinson, M. M., Rosenbloom, N., Salzmann, U., and Sohl, L. E.: Pliocene Model Intercomparison Project (PlioMIP): experimental design and boundary conditions (Experiment 1), *Geosci. Model Dev.*, 3, 227–242, doi:10.5194/gmd-3-227-2010, 2010.

Haywood, A. M., Dowsett, H. J., Robinson, M. M., Stoll, D. K., Dolan, A. M., Lunt, D. J., Otto-Bliesner, B., and Chandler, M. A.: Pliocene Model Intercomparison Project (PlioMIP): experimental design and boundary conditions (Experiment 2), *Geosci. Model Dev.*, 4, 571–577, doi:10.5194/gmd-4-571-2011, 2011.

Hibler, W. D.: Dynamic thermodynamic sea ice model, *J. Phys. Oceanogr.*, 9, 815–846, 1979.

Ingram, W. J., Woodward, S., and Edwards, J.: Unified Model Documentation Paper No 23: Radiation Climate Research, Meteorological Office, London Road, Bracknell, Berkshire, RG12 2SY, UK, 1997.

Johns, T. C., Carnell, R. E., Crossley, J. F., Gregory, J. M., Mitchell, J. F. B., Senior, C. A., Tett, S. F. B., and Wood, R. A.: The second Hadley Centre coupled ocean-atmosphere GCM: model description, spinup and validation, *Clim. Dynam.*, 13, 103–134, 1997.

Kaplan, J. O., Bigelow, N. H., Prentice, I. C., Harrison, S. P., Bartlein, P. J., Christensen, T. R., Cramer, W., Matveyeva, N. V., Mcguire, A. D., Murray, D. F., Razzhivin, V. Y., Smith, B., Walker, D. A., Anderson, P. M., Andreev, A. A., Brubaker, L. B., Edwards, M. E., and Lozhkin, A. V.: Climate change and Arctic ecosystems: 2. Modeling, paleodata-model comparisons, and future projections, *J. Geophys. Res.-Atmos.*, 108, 8171, doi:10.1029/2002JD002559, 2003.

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kraus, E. B. and Turner, J. S.: A one-dimensional model of seasonal thermocline, 2. General theory and its consequences, *Tellus*, 19, 98–106, doi:10.1111/j.2153-3490.1967.tb01462.x, 1967.
- Lunt, D. J., Haywood, A. M., Schmidt, G. A., Salzmann, U., Valdes, P. J., and Dowsett, H. J.: Earth system sensitivity inferred from Pliocene modelling and data, *Nat. Geosci.*, 3, 60–64, 2010.
- Lunt, D. J., Haywood, A. M., Schmidt, G. A., Salzmann, U., Valdes, P. J., Dowsett, H. J., and Loptson, C. A.: On the causes of mid-Pliocene warmth and polar amplification, *Earth Planet. Sc. Lett.*, 321–322, 128–138, 2012.
- Milton, S. F. and Wilson, C. A.: The impact of parameterized subgrid-scale orographic forcing on systematic errors in a global NWP model, *Mon. Weather Rev.*, 124, 2023–2045, 1996.
- Ncar/Navy: Global 10-minute elevation data, Digital Tape Available through National Oceanic and Atmospheric Administration, N. G. D. C., Boulder, Co., 1984.
- Pope, V. D., Gallani, M. L., Rowntree, P. R., and Stratton, R. A.: The impact of new physical parametrizations in the Hadley Centre climate model: HadAM3, *Clim. Dynam.*, 16, 123–146, 2000.
- Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., Rowell, D. P., Kent, E. C., and Kaplan, A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century, *J. Geophys. Res.-Atmos.*, 108, 4407, doi:10.1029/2002jd002670, 2003.
- Robinson, M. M., Valdes, P. J., Haywood, A. M., Dowsett, H. J., Hill, D. J., and Jones, S. M.: Bathymetric controls on Pliocene North Atlantic and Arctic sea surface temperature and deepwater production, *Palaeogeogr. Palaeoclimatol.*, 309, 92–97, 2011.
- Smith, R. N. B.: A scheme for predicting layer clouds and their water-content in a general-circulation model, *Q. J. Roy. Meteor. Soc.*, 116, 435–460, 1990.
- Wilson, M. F. and Henderson-Sellers, A.: A global archive of land cover and soils data for use in general-circulation climate models, *J. Climatol.*, 5, 119–143, 1985.

Table 1. Summary of boundary conditions for Pliocene simulations.

	EXPT 1			EXPT 2		
	Pliocene	Control	Pre-Ind: Alt Veg	Pliocene	Control	PRISM2
Land-sea mask	PlioMIP preferred	UM standard	UM standard	UM standard	UM standard	UM standard
Topography including ice sheets	PlioMIP anomaly method	UM standard	UM standard	PlioMIP anomaly method	UM standard	PRISM2 absolute
Land-surface properties including ice	PlioMIP BIOME4 vegetation	UM standard	BAS-Observ modern vegetation	PlioMIP BIOME4 vegetation	UM standard	PRISM2 vegetation
Soils	UM standard	UM standard	UM standard	UM standard	UM standard	UM standard
River routing	n/a	n/a	n/a	UM standard	UM standard	UM standard
SST	PlioMIP anomaly method	UM standard	UM standard	n/a	n/a	n/a
Bathymetry	n/a	n/a	n/a	UM standard	UM standard	UM standard
Solar constant ($W m^{-2}$)	1365	1365	1365	1365	1365	1365
Sine obliquity (angle)	0.397789 (23.440°)	0.397789 (23.440°)	0.397789 (23.440°)	0.397789 (23.440°)	0.397789 (23.440°)	0.397789 (23.440°)
Eccentricity	0.0167	0.0167	0.0167	0.0167	0.0167	0.0167
Perihelion (day/360)	2.5	2.5	2.5	2.5	2.5	2.5
CO ₂ (ppm)	405	280	280	405	280	401
N ₂ O (ppb)	269	269	269	269	269	269
CH ₄ (ppb)	761	761	761	761	761	761
Ozone	standard UM climatology	standard UM climatology	standard UM climatology	standard UM climatology	standard UM climatology	standard UM climatology
Integration length (yr)	50	50	50	500	200	200
Climatological means	30	30	30	50	50	30

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[⏪](#) [⏩](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.

Table 2. Summary of overall global means and differences from control for Experiments 1 and 2.

	Expt 1		Expt 2	
	Absolute Pliocene	Absolute control	Absolute Pliocene	Absolute control
TOA global energy imbalance (Wm^{-2})	6.01	1.50	0.22	-0.12
	Absolute Pliocene	Delta (Plio-control)	Absolute Pliocene	Delta (Plio-control)
Global mean annual surface air temp ($^{\circ}\text{C}$)	14.93	2.53	16.34	3.27
Global mean total precipitation rate (mm day^{-1})	2.94	0.12	3.05	0.17

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 1. Summary of land surface boundary conditions for PlioMIP simulations: panels (a) to (c) show orography projected onto the appropriate land-sea mask for control, Experiment 1 and Experiment 2 Pliocene simulations. Panels (d) to (f) show snow-free albedo for control, Experiment 1 and Experiment 2 Pliocene simulations – values above 0.3 shown in beige represent ice-sheets.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 2. Summary of sea surface temperature boundary conditions for atmosphere-only PlioMIP simulations: panels (a) and (b) show January temperatures for control and Pliocene, panels (c) and (d) show July temperatures for control and Pliocene.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 3. Time series of global mean temperatures.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 4. Summary of zonal mean of predicted surface air temperatures for Experiment 1: panels (a) to (c) show absolute values for Pliocene (red) and control (blue) for global, land and ocean means, respectively. Panels (d) to (f) show the corresponding differences in temperature between the 2 simulations.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 5. Summary of surface air temperatures for Experiment 1: panels (a) to (c) show annual mean temperatures for Pliocene and control simulations and the difference between them. Panels (d) to (f) similarly show the DJF means and panels (g) to (i) the JJA means.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 6. Summary of mean precipitation (mm day^{-1}) for Experiment 1: panels (a) to (c) show annual mean precipitation for Pliocene and control simulations and the difference between them. Panels (d) to (f) similarly show the DJF means and panels (g) to (i) the JJA means.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling Pliocene
climate using the
Hadley Centre Model

F. J. Bragg et al.




Fig. 7. Summary of zonal mean of predicted surface air temperatures for Experiment 2: panels (a) to (c) show absolute values for Pliocene (red) and control (blue) for global, land and ocean means, respectively. Panels (d) to (f) show the corresponding differences in temperature between the 2 simulations.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 8. Summary of surface air temperatures for Experiment 2: panels (a) to (c) show annual mean temperatures for Pliocene and control simulations and the difference between them. Panels (d) to (f) similarly show the DJF means and panels (g) to (i) the JJA means.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 9. Summary of mean precipitation (mm day^{-1}) for Experiment 2: panels (a) to (c) show annual mean precipitation for Pliocene and control simulations and the difference between them. Panels (d) to (f) similarly show the DJF means and panels (g) to (i) the JJA means.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 10. Summary of predicted ocean behaviour for Experiment 2: panels (a) to (c) show sea surface temperatures for Pliocene, control and the difference between them; panels (d) to (f) similarly show salinity. Panels (g) and (h) show Atlantic overturning streamfunction for the Pliocene and control and (i) and (j) show the Pacific overturning streamfunction.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 11. “Difference of difference” showing the difference in Pliocene – control anomalies predicted by Experiments 1 and 2. Differences are shown as Experiment 2 – Experiment 1. Panels (a) to (c) show zonal mean difference globally, for land only and ocean only. Panels (d) to (f) show surface air temperature change as annual mean, DJF mean and JJA mean. Panels (g) to (i) show precipitation change as annual mean, DJF mean and JJA mean.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 12. Model data comparison. Contours show Pliocene – control sea surface temperatures predicted for Experiment 2. Filled circles show PRISM3 – HadISST pre-industrial climatology annual means taken over 1870–1899.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 13. “Difference of difference” showing the difference in Pliocene – control anomalies predicted using PRISM3 and PRISM2 boundary conditions. Differences are shown as PRISM3–PRISM2. Panels (a) to (c) show zonal mean difference globally, for land only and ocean only. Panels (d) to (f) show surface air temperature change as annual mean, DJF mean and JJA mean. Panels (g) to (i) show precipitation change as annual mean, DJF mean and JJA mean.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Modelling Pliocene climate using the Hadley Centre Model

F. J. Bragg et al.




Fig. 14. “Difference of difference” showing the difference in Pliocene – control anomalies predicted using standard UM pre-industrial vegetation and using PlioMIP “BAS-Observ” vegetation to derive land surface properties. Differences are shown as standard UM pre-industrial – BAS Modern. Panels (a) to (c) show zonal mean difference globally, for land only and ocean only. Panels (d) to (f) show surface air temperature change as annual mean, DJF mean and JJA mean. Panels (g) to (i) show precipitation change as annual mean, DJF mean and JJA mean.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)