



1

## Global Carbon Budget 2017

2 Corinne Le Quéré<sup>1</sup>, Robbie M. Andrew<sup>2</sup>, Pierre Friedlingstein<sup>3</sup>, Stephen Sitch<sup>4</sup>, Julia Pongratz<sup>5</sup>, Andrew C.  
3 Manning<sup>6</sup>, Jan Ivar Korsbakken<sup>2</sup>, Glen P. Peters<sup>2</sup>, Josep G. Canadell<sup>7</sup>, Robert B. Jackson<sup>8</sup>, Thomas A. Boden<sup>9</sup>,  
4 Pieter P. Tans<sup>10</sup>, Oliver D. Andrews<sup>1</sup>, Vivek K. Arora<sup>11</sup>, Dorothee C. E. Bakker<sup>6</sup>, Leticia Barbero<sup>12,13</sup>, Meike  
5 Becker<sup>14,15</sup>, Richard A. Betts<sup>16,4</sup>, Laurent Bopp<sup>17</sup>, Frédéric Chevallier<sup>18</sup>, Louise P. Chini<sup>19</sup>, Philippe Ciais<sup>18</sup>,  
6 Catherine E. Cosca<sup>20</sup>, Jessica Cross<sup>20</sup>, Kim Currie<sup>21</sup>, Thomas Gasser<sup>22</sup>, Ian Harris<sup>23</sup>, Judith Hauck<sup>24</sup>, Vanessa  
7 Haverd<sup>25</sup>, Richard A. Houghton<sup>26</sup>, Christopher W. Hunt<sup>27</sup>, George Hurtt<sup>19</sup>, Tatiana Ilyina<sup>5</sup>, Atul K. Jain<sup>28</sup>,  
8 Etsushi Kato<sup>29</sup>, Markus Kautz<sup>30</sup>, Ralph F. Keeling<sup>31</sup>, Kees Klein Goldewijk<sup>32</sup>, Arne Körtzinger<sup>33</sup>, Peter  
9 Landschützer<sup>5</sup>, Nathalie Lefèvre<sup>34</sup>, Andrew Lenton<sup>35,36</sup>, Sebastian Lienert<sup>37,38</sup>, Ivan Lima<sup>39</sup>, Danica  
10 Lombardozzi<sup>40</sup>, Nicolas Metzler<sup>34</sup>, Frank Millero<sup>41</sup>, Pedro M. S. Monteiro<sup>42</sup>, David R. Munro<sup>43</sup>, Julia E. M. S.  
11 Nabel<sup>5</sup>, Shin-ichiro Nakaoka<sup>44</sup>, Yukihiro Nojiri<sup>44</sup>, X. Antonio Padin<sup>45</sup>, Anna Peregon<sup>18</sup>, Benjamin Pfeil<sup>14,15</sup>,  
12 Denis Pierrot<sup>12,13</sup>, Benjamin Poulter<sup>46,47</sup>, Gregor Rehder<sup>48</sup>, Janet Reimer<sup>49</sup>, Christian Rödenbeck<sup>50</sup>, Jörg  
13 Schwinger<sup>51</sup>, Roland Séférian<sup>52</sup>, Ingunn Skjelvan<sup>51</sup>, Benjamin D. Stocker<sup>53</sup>, Hanqin Tian<sup>54</sup>, Bronte  
14 Tilbrook<sup>35,36</sup>, Ingrid T. van der Laan-Luijkx<sup>55</sup>, Guido R. van der Werf<sup>56</sup>, Steven van Heuven<sup>57</sup>, Nicolas Viovy<sup>18</sup>,  
15 Nicolas Vuichard<sup>18</sup>, Anthony P. Walker<sup>58</sup>, Andrew J. Watson<sup>4</sup>, Andrew J. Wiltshire<sup>16</sup>, Sönke Zaehle<sup>50</sup>, Dan  
16 Zhu<sup>18</sup>

17

18 <sup>1</sup>Tyndall Centre for Climate Change Research, University of East Anglia, Norwich Research Park,  
19 Norwich NR4 7TJ, UK

20 <sup>2</sup>CICERO Center for International Climate Research, Oslo, Norway

21 <sup>3</sup>College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter EX4 4QF, UK

22 <sup>4</sup>College of Life and Environmental Sciences, University of Exeter, Exeter EX4 4RJ, UK

23 <sup>5</sup>Max Planck Institute for Meteorology, Hamburg, Germany

24 <sup>6</sup>Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East  
25 Anglia, Norwich Research Park, Norwich NR4 7TJ, UK

26 <sup>7</sup>Global Carbon Project, CSIRO Oceans and Atmosphere, GPO Box 1700, Canberra, ACT 2601, Australia

27 <sup>8</sup>Department of Earth System Science, Woods Institute for the Environment, and Precourt Institute for  
28 Energy, Stanford University, Stanford, CA 94305, USA

29 <sup>9</sup>Climate Change Science Institute, Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA

30 <sup>10</sup>National Oceanic & Atmospheric Administration, Earth System Research Laboratory (NOAA/ESRL),  
31 Boulder, CO 80305, USA

32 <sup>11</sup>Canadian Centre for Climate Modelling and Analysis, Climate Research Division, Environment and  
33 Climate Change Canada, Victoria, BC, Canada

34 <sup>12</sup>Cooperative Institute for Marine and Atmospheric Studies, Rosenstiel School for Marine and  
35 Atmospheric Science, University of Miami, Miami, FL 33149, USA

36 <sup>13</sup>National Oceanic & Atmospheric Administration/Atlantic Oceanographic & Meteorological Laboratory  
37 (NOAA/AOML), Miami, FL 33149, USA

38 <sup>14</sup>Geophysical Institute, University of Bergen, 5020 Bergen, Norway

39 <sup>15</sup>Bjerknes Centre for Climate Research, 5007 Bergen, Norway

40 <sup>16</sup>Met Office Hadley Centre, FitzRoy Road, Exeter EX1 3PB, UK

41 <sup>17</sup>Laboratoire de Météorologie Dynamique, Institut Pierre-Simon Laplace, CNRS-ENS-UPMC-X,  
42 Département de Géosciences, Ecole Normale Supérieure, 24 rue Lhomond, 75005 Paris, France

43 <sup>18</sup>Laboratoire des Sciences du Climat et de l'Environnement, Institut Pierre-Simon Laplace, CEA-CNRS-  
44 UVSQ, CE Orme des Merisiers, 91191 Gif sur Yvette Cedex, France

45 <sup>19</sup>Department of Geographical Sciences, University of Maryland, College Park, Maryland 20742, USA

46 <sup>20</sup>Pacific Marine Environmental Laboratory, National Oceanic and Atmospheric Administration, Seattle,  
47 WA 98115, USA

48 <sup>21</sup>National Institute of Water and Atmospheric Research (NIWA), Dunedin 9054, New Zealand

49 <sup>22</sup>International Institute for Applied Systems Analysis (IIASA), 2361 Laxenburg, Austria





1 **Abstract**  
2 Accurate assessment of anthropogenic carbon dioxide (CO<sub>2</sub>) emissions and their redistribution  
3 among the atmosphere, ocean, and terrestrial biosphere – the ‘global carbon budget’ – is  
4 important to better understand the global carbon cycle, support the development of climate  
5 policies, and project future climate change. Here we describe data sets and methodology to  
6 quantify the five major components of the global carbon budget and their uncertainties. CO<sub>2</sub>  
7 emissions from fossil fuels and industry ( $E_{FF}$ ) are based on energy statistics and cement production  
8 data, respectively, while emissions from land-use change ( $E_{LUC}$ ), mainly deforestation, are based  
9 on land-cover change data and bookkeeping models. The global atmospheric CO<sub>2</sub> concentration is  
10 measured directly and its rate of growth ( $G_{ATM}$ ) is computed from the annual changes in  
11 concentration. The ocean CO<sub>2</sub> sink ( $S_{OCEAN}$ ) and terrestrial CO<sub>2</sub> sink ( $S_{LAND}$ ) are estimated with  
12 global process models constrained by observations. The resulting carbon budget imbalance ( $B_{IM}$ ),  
13 the difference between the estimated total emissions and the estimated changes in the  
14 atmosphere, ocean, and terrestrial biosphere, is a measure of our imperfect data and  
15 understanding of the contemporary carbon cycle. All uncertainties are reported as  $\pm 1\sigma$ . For the  
16 last decade available (2007-2016),  $E_{FF}$  was  $9.4 \pm 0.5 \text{ GtC yr}^{-1}$ ,  $E_{LUC}$   $1.3 \pm 0.7 \text{ GtC yr}^{-1}$ ,  $G_{ATM}$   $4.7 \pm 0.1$   
17  $\text{GtC yr}^{-1}$ ,  $S_{OCEAN}$   $2.4 \pm 0.5 \text{ GtC yr}^{-1}$ , and  $S_{LAND}$   $3.0 \pm 0.8 \text{ GtC yr}^{-1}$ , with a budget imbalance  $B_{IM}$  of  $0.6$   
18  $\text{GtC yr}^{-1}$  indicating overestimated emissions and/or underestimated sinks. For year 2016 alone, the  
19 growth in  $E_{FF}$  was approximately zero and emissions remained at  $9.9 \pm 0.5 \text{ GtC yr}^{-1}$ . Also for 2016,  
20  $E_{LUC}$  was  $1.3 \pm 0.7 \text{ GtC yr}^{-1}$ ,  $G_{ATM}$  was  $6.1 \pm 0.2 \text{ GtC yr}^{-1}$ ,  $S_{OCEAN}$  was  $2.6 \pm 0.5 \text{ GtC yr}^{-1}$  and  $S_{LAND}$  was  
21  $2.7 \pm 1.0 \text{ GtC yr}^{-1}$ , with a small  $B_{IM}$  of  $-0.3 \text{ GtC}$ .  $G_{ATM}$  continued to be higher in 2016 compared to  
22 the past decade (2007-2016), reflecting in part the higher fossil emissions and smaller  $S_{LAND}$  for  
23 that year consistent with El Niño conditions. The global atmospheric CO<sub>2</sub> concentration reached  
24  $402.8 \pm 0.1 \text{ ppm}$  averaged over 2016. For 2017, preliminary data indicate a renewed growth in  $E_{FF}$   
25 of +2.0% (range of 0.8% to 3.0%) based on national emissions projections for China, USA, and  
26 India, and projections of Gross Domestic Product corrected for recent changes in the carbon  
27 intensity of the economy for the rest of the world. For 2017, initial data indicate an increase in  
28 atmospheric CO<sub>2</sub> concentration of around 5.3 GtC (2.5 ppm), attributed to a combination of  
29 increasing emissions and receding El Niño conditions. This living data update documents changes  
30 in the methods and data sets used in this new global carbon budget compared with previous  
31 publications of this data set (Le Quéré et al., 2016; 2015b; 2015a; 2014; 2013). All results  
32 presented here can be downloaded from <https://doi.org/10.18160/GCP-2017>.



## 1 **1 Introduction**

2 The concentration of carbon dioxide (CO<sub>2</sub>) in the atmosphere has increased from approximately  
3 277 parts per million (ppm) in 1750 (Joos and Spahni, 2008), the beginning of the Industrial Era, to  
4 402.8 ± 0.1 ppm in 2016 (Dlugokencky and Tans, 2016; Fig. 1). The atmospheric CO<sub>2</sub> increase  
5 above preindustrial levels was, initially, primarily caused by the release of carbon to the  
6 atmosphere from deforestation and other land-use change activities (Ciais et al., 2013). While  
7 emissions from fossil fuels started before the Industrial Era, they only became the dominant  
8 source of anthropogenic emissions to the atmosphere from around 1920 and their relative share  
9 has continued to increase until present. Anthropogenic emissions occur on top of an active natural  
10 carbon cycle that circulates carbon between the reservoirs of the atmosphere, ocean, and  
11 terrestrial biosphere on time scales from sub-daily to millennia, while exchanges with geologic  
12 reservoirs occur at longer timescales (Archer et al., 2009).

13 The global carbon budget presented here refers to the mean, variations, and trends in the  
14 perturbation of CO<sub>2</sub> in the atmosphere, referenced to the beginning of the Industrial Era. It  
15 quantifies the input of CO<sub>2</sub> to the atmosphere by emissions from human activities, the growth rate  
16 of atmospheric CO<sub>2</sub> concentration, and the resulting changes in the storage of carbon in the land  
17 and ocean reservoirs in response to increasing atmospheric CO<sub>2</sub> levels, climate change and  
18 variability, and other anthropogenic and natural changes (Fig. 2). An understanding of this  
19 perturbation budget over time and the underlying variability and trends of the natural carbon  
20 cycle are necessary to understand the response of natural sinks to changes in climate, CO<sub>2</sub> and  
21 land-use change drivers, and the permissible emissions for a given climate stabilization target.

22 The components of the CO<sub>2</sub> budget that are reported annually in this paper include separate  
23 estimates for the CO<sub>2</sub> emissions from (1) fossil fuel combustion and oxidation and cement  
24 production ( $E_{FF}$ ; GtC yr<sup>-1</sup>) and (2) the emissions resulting from deliberate human activities on land  
25 leading to land-use change ( $E_{LUC}$ ; GtC yr<sup>-1</sup>); and their partitioning among (3) the growth rate of  
26 atmospheric CO<sub>2</sub> concentration ( $G_{ATM}$ ; GtC yr<sup>-1</sup>), and the uptake of CO<sub>2</sub> (the 'CO<sub>2</sub> sinks') in (4) the  
27 ocean ( $S_{OCEAN}$ ; GtC yr<sup>-1</sup>) and (5) on land ( $S_{LAND}$ ; GtC yr<sup>-1</sup>). The CO<sub>2</sub> sinks as defined here conceptually  
28 include the response of the land (including inland waters and estuaries) and ocean (including  
29 coasts and seaward edge) to elevated CO<sub>2</sub> and changes in climate, rivers, and other environmental  
30 conditions, although in practice not all processes are accounted for (see Section 2.7). The global  
31 emissions and their partitioning among the atmosphere, ocean and land are in reality in balance,



1 however due to imperfect spatial and/or temporal data coverage, errors in each estimate and due  
2 to smaller terms not included in our budget estimate (discussed in Section 2.7), their sum does  
3 not necessarily add up to zero. We introduce here a budget imbalance ( $B_{IM}$ ), which is a measure of  
4 the mismatch between the estimated emissions and the estimated changes in the atmosphere,  
5 land and ocean. This is an important change in the calculation of the global carbon budget. With  
6 this change, the full global carbon budget now reads:

$$E_{FF} + E_{LUC} = G_{ATM} + S_{OCEAN} + S_{LAND} + B_{IM}. \quad (1)$$

7  $G_{ATM}$  is usually reported in  $\text{ppm yr}^{-1}$ , which we convert to units of carbon mass per year,  $\text{GtC yr}^{-1}$ ,  
8 using  $1 \text{ ppm} = 2.12 \text{ GtC}$  (Table 1). We also include a quantification of  $E_{FF}$  by country, computed  
9 with both territorial and consumption based accounting (see Sect. 2). Equation (1) partly omits the  
10 net input of  $\text{CO}_2$  to the atmosphere from the chemical oxidation of reactive carbon-containing  
11 gases from sources other than the combustion of fossil fuels (discussed in Sect. 2.7).

12 The  $\text{CO}_2$  budget has been assessed by the Intergovernmental Panel on Climate Change (IPCC) in all  
13 assessment reports (Ciais et al., 2013; Denman et al., 2007; Prentice et al., 2001; Schimel et al.,  
14 1995; Watson et al., 1990), and by others (e.g. Ballantyne et al., 2012). The IPCC methodology has  
15 been adapted and used by the Global Carbon Project (GCP, [www.globalcarbonproject.org](http://www.globalcarbonproject.org)), which  
16 has coordinated a cooperative community effort for the annual publication of global carbon  
17 budgets up to year 2005 (Raupach et al., 2007; including fossil emissions only), year 2006  
18 (Canadell et al., 2007), year 2007 (published online; GCP, 2007), year 2008 (Le Quéré et al., 2009),  
19 year 2009 (Friedlingstein et al., 2010), year 2010 (Peters et al., 2012b), year 2012 (Le Quéré et al.,  
20 2013; Peters et al., 2013), year 2013 (Le Quéré et al., 2014), year 2014 (Friedlingstein et al., 2014;  
21 Le Quéré et al., 2015b), year 2015 (Jackson et al., 2016; Le Quéré et al., 2015a), and most recently  
22 year 2016 (Le Quéré et al., 2016). Each of these papers updated previous estimates with the latest  
23 available information for the entire time series.

24 We adopt a range of  $\pm 1$  standard deviation ( $\sigma$ ) to report the uncertainties in our estimates,  
25 representing a likelihood of 68% that the true value will be within the provided range if the errors  
26 have a Gaussian distribution. This choice reflects the difficulty of characterising the uncertainty in  
27 the  $\text{CO}_2$  fluxes between the atmosphere and the ocean and land reservoirs individually,  
28 particularly on an annual basis, as well as the difficulty of updating the  $\text{CO}_2$  emissions from land-  
29 use change. A likelihood of 68% provides an indication of our current capability to quantify each  
30 term and its uncertainty given the available information. For comparison, the Fifth Assessment



1 Report of the IPCC (AR5) generally reported a likelihood of 90% for large data sets whose  
2 uncertainty is well characterised, or for long time intervals less affected by year-to-year variability.  
3 Our 68% uncertainty value is near the 66% which the IPCC characterises as ‘likely’ for values falling  
4 into the  $\pm 1\sigma$  interval. The uncertainties reported here combine statistical analysis of the  
5 underlying data and expert judgement of the likelihood of results lying outside this range. The  
6 limitations of current information are discussed in the paper and have been examined in detail  
7 elsewhere (Ballantyne et al., 2015; Zscheischler et al., 2017).

8 All quantities are presented in units of gigatonnes of carbon (GtC,  $10^{15}$  gC), which is the same as  
9 petagrams of carbon (PgC; Table 1). Units of gigatonnes of  $\text{CO}_2$  (or billion tonnes of  $\text{CO}_2$ ) used in  
10 policy are equal to 3.664 multiplied by the value in units of GtC.

11 This paper provides a detailed description of the data sets and methodology used to compute the  
12 global carbon budget estimates for the period preindustrial (1750) to 2016 and in more detail for  
13 the period 1959 to 2016. We also provide decadal averages starting in 1960 including the last  
14 decade (2007-2016), results for the year 2016, and a projection for year 2017. Finally we provide  
15 cumulative emissions from fossil fuels and land-use change since year 1750, the preindustrial  
16 period, and since year 1870, the reference year for the cumulative carbon estimate used by the  
17 IPCC (AR5) based on the availability of global temperature data (Stocker et al., 2013). This paper is  
18 updated every year using the format of ‘living data’ to keep a record of budget versions and the  
19 changes in new data, revision of data, and changes in methodology that lead to changes in  
20 estimates of the carbon budget. Additional materials associated with the release of each new  
21 version will be posted at the Global Carbon Project (GCP) website  
22 (<http://www.globalcarbonproject.org/carbonbudget>), with fossil fuel emissions also available  
23 through the Global Carbon Atlas (<http://www.globalcarbonatlas.org>). With this approach, we aim  
24 to provide the highest transparency and traceability in the reporting of  $\text{CO}_2$ , the key driver of  
25 climate change.

## 26 **2 Methods**

27 Multiple organizations and research groups around the world generated the original  
28 measurements and data used to complete the global carbon budget. The effort presented here is  
29 thus mainly one of synthesis, where results from individual groups are collated, analysed and  
30 evaluated for consistency. We facilitate access to original data with the understanding that



1 primary data sets will be referenced in future work (See Table 2 for ‘How to cite’ the data sets).  
2 Descriptions of the measurements, models, and methodologies follow below and in depth  
3 descriptions of each component are described elsewhere.  
4 This is the 12<sup>th</sup> version of the global carbon budget and the sixth revised version in the format of a  
5 living data update. It builds on the latest published global carbon budget of Le Quéré et al. (2016).  
6 The main changes are: (1) the inclusion of data to year 2016 (inclusive) and a projection for the  
7 global carbon budget for year 2017; (2) the use of two bookkeeping models to assess  $E_{LUC}$  (instead  
8 of one), (3) the use of Dynamic Global Vegetation Models (DGVMs) to assess  $S_{LAND}$ , (4) the  
9 introduction of the budget imbalance  $B_{IM}$  as the difference between the estimated emissions and  
10 sinks, thus removing the assumption in previous global carbon budgets that the main  
11 uncertainties are primarily on the land sink ( $S_{LAND}$ ), and recognising uncertainties in the estimate  
12 of  $S_{ocean}$ , particularly on decadal time-scales, (5) the addition of a table presenting the major  
13 known sources of uncertainties, and (6) the expansion of the model descriptions. The main  
14 methodological differences between annual carbon budgets are summarised in Table 3.

## 15 **2.1 CO<sub>2</sub> emissions from fossil fuels and industry ( $E_{FF}$ )**

### 16 **2.1.1 Emissions estimates**

17 The estimates of global and national CO<sub>2</sub> emissions from fossil fuels, including gas flaring and  
18 cement production ( $E_{FF}$ ), relies primarily on energy consumption data, specifically data on  
19 hydrocarbon fuels, collated and archived by several organisations (Andres et al., 2012). We use  
20 four main datasets for historical emissions (1751-2016):

- 21 1. Global and national emission estimates from CDIAC for the time period 1751-2014 (Boden  
22 et al., 2017), as it is the only data set that extends back to 1751 by country.
- 23 2. Official UNFCCC national inventory reports for 1990-2015 for the 42 Annex I countries in  
24 the UNFCCC (UNFCCC, 2017), as we assess these to be the most accurate estimates and  
25 are periodically reviewed.
- 26 3. The BP Statistical Review of World Energy (BP, 2017), to project the emissions forward to  
27 2016 to ensure the most recent estimates possible.
- 28 4. The US Geological Survey estimates of cement production (USGS, 2017), to estimate  
29 cement emissions.



1 In the following we provide more details in each dataset and additional modifications that are  
2 required to make the dataset consistent and usable.

3 *CDIAC*: The CDIAC estimates have been updated annually to include the most recent year (2014)  
4 and to include statistical revisions to recent historical data (UN, 2017). Fuel masses and volumes  
5 are converted to fuel energy content using country-level coefficients provided by the UN, and  
6 then converted to CO<sub>2</sub> emissions using conversion factors that take into account the relationship  
7 between carbon content and energy (heat) content of the different fuel types (coal, oil, gas, gas  
8 flaring) and the combustion efficiency (Marland and Rotty, 1984).

9 *UNFCCC*: Estimates from the UNFCCC national inventory reports follow the IPCC guidelines (IPCC,  
10 2006), but have a slightly larger system boundary than CDIAC by including emissions coming from  
11 carbonates other than in cement manufacture. We reallocate the detailed UNFCCC estimates to  
12 the CDIAC definitions of coal, oil, gas, cement, and other to allow consistent comparisons over  
13 time and between countries.

14 *BP*: For the most recent period when the UNFCCC (2016) and CDIAC (2015-2016) estimates are not  
15 available, we generate preliminary estimates using the BP Statistical Review of World Energy  
16 (Andres et al., 2014; Myhre et al., 2009). We apply the BP growth rates by fuel type (coal, oil, gas)  
17 to estimate 2016 emissions based on 2015 estimates (UNFCCC), and to estimate 2015 and 2016  
18 based on 2014 estimates (CDIAC). BP's dataset explicitly covers about 70 countries (96% of global  
19 emissions), and for the remaining countries we use growth rates from the sub-region the country  
20 belongs to. For the most recent years, flaring is assumed constant from the most recent available  
21 year of data (2015 for countries that report to the UNFCCC, 2014 for the remainder).

22 *USGS*: Estimates of emissions from cement production are based on USGS (USGS, 2017), applying  
23 the emission factors from CDIAC (Marland and Rotty, 1984). The CDIAC cement emissions are  
24 known to be high, and are likely to be revised downwards next year (Andrew, 2017). Some  
25 fraction of the CaO and MgO in cement is returned to the carbonate form during cement  
26 weathering but this is omitted here (Xi et al., 2016).

27 *Country mappings*: The published CDIAC data set includes 256 countries and regions. This list  
28 includes countries that no longer exist, such as the USSR and Yugoslavia. We reduce the list to 220  
29 countries by reallocating emissions to the currently defined territories, using mass-preserving  
30 aggregation or disaggregation. Examples of aggregation include merging East and West Germany



1 to the currently defined Germany. Examples of disaggregation include reallocating the emissions  
2 from former USSR to the resulting independent countries. For disaggregation, we use the emission  
3 shares when the current territories first appeared, and thus historical estimates of disaggregated  
4 countries should be treated with extreme care.

5 *Global total:* Our global estimate is based on CDIAC, and this is greater than the sum of emissions  
6 from all countries. This is largely attributable to emissions that occur in international territory, in  
7 particular, the combustion of fuels used in international shipping and aviation (bunker fuels). The  
8 emissions from international bunker fuels are calculated based on where the fuels were loaded,  
9 but we do not include them in the national emissions estimates. Other differences occur 1)  
10 because the sum of imports in all countries is not equal to the sum of exports, and 2) because of  
11 inconsistent national reporting, differing treatment of oxidation of non-fuel uses of hydrocarbons  
12 (e.g. as solvents, lubricants, feedstocks, etc.), and 3) changes in fuel stored (Andres et al., 2012).

### 13 **2.1.2 Uncertainty assessment for $E_{FF}$**

14 We estimate the uncertainty of the global emissions from fossil fuels and industry at  $\pm 5\%$  (scaled  
15 down from the published  $\pm 10\%$  at  $\pm 2\sigma$  to the use of  $\pm 1\sigma$  bounds reported here; Andres et al.,  
16 2012). This is consistent with a more detailed recent analysis of uncertainty of  $\pm 8.4\%$  at  $\pm 2\sigma$   
17 (Andres et al., 2014) and at the high-end of the range of  $\pm 5\text{-}10\%$  at  $\pm 2\sigma$  reported by Ballantyne et  
18 al. (2015). This includes an assessment of uncertainties in the amounts of fuel consumed, the  
19 carbon and heat contents of fuels, and the combustion efficiency. While we consider a fixed  
20 uncertainty of  $\pm 5\%$  for all years, the uncertainty as a percentage of the emissions is growing with  
21 time because of the larger share of global emissions from emerging economies and developing  
22 countries (Marland et al., 2009). Generally, emissions from mature economies with good  
23 statistical processes have an uncertainty of only a few per cent (Marland, 2008), while developing  
24 countries such as China have uncertainties of around  $\pm 10\%$  (for  $\pm 1\sigma$ ; Gregg et al., 2008).  
25 Uncertainties of emissions are likely to be mainly systematic errors related to underlying biases of  
26 energy statistics and to the accounting method used by each country.

27 We assign a medium confidence to the results presented here because they are based on indirect  
28 estimates of emissions using energy data (Durant et al., 2011). There is only limited and indirect  
29 evidence for emissions, although there is a high agreement among the available estimates within  
30 the given uncertainty (Andres et al., 2014; Andres et al., 2012), and emission estimates are



1 consistent with a range of other observations (Ciais et al., 2013), even though their regional and  
2 national partitioning is more uncertain (Francey et al., 2013).

### 3 **2.1.3 Emissions embodied in goods and services**

4 CDIAC, UNFCCC, and BP national emission statistics ‘include greenhouse gas emissions and  
5 removals taking place within national territory and offshore areas over which the country has  
6 jurisdiction’ (Rypdal et al., 2006), and are called territorial emission inventories. Consumption-  
7 based emission inventories allocate emissions to products that are consumed within a country,  
8 and are conceptually calculated as the territorial emissions minus the ‘embodied’ territorial  
9 emissions to produce exported products plus the emissions in other countries to produce  
10 imported products (Consumption = Territorial – Exports + Imports). Consumption-based emission  
11 attribution results (e.g. Davis and Caldeira, 2010) provide additional information to territorial-  
12 based emissions that can be used to understand emission drivers (Hertwich and Peters, 2009) and  
13 quantify emission transfers by the trade of products between countries (Peters et al., 2011b). The  
14 consumption-based emissions have the same global total, but reflect the trade-driven movement  
15 of emissions across the Earth's surface in response to human activities.

16 We estimate consumption-based emissions from 1990-2015 by enumerating the global supply  
17 chain using a global model of the economic relationships between economic sectors within and  
18 between every country (Andrew and Peters, 2013; Peters et al., 2011a). Our analysis is based on  
19 the economic and trade data from the Global Trade and Analysis Project (GTAP; Narayanan et al.,  
20 2015), and we make detailed estimates for the years 1997 (GTAP version 5), 2001 (GTAP6), and  
21 2004, 2007, and 2011 (GTAP9.2), covering 57 sectors and 141 countries and regions. The detailed  
22 results are then extended into an annual time-series from 1990 to the latest year of the Gross  
23 Domestic Product (GDP) data (2015 in this budget), using GDP data by expenditure in current  
24 exchange rate of US dollars (USD; from the UN National Accounts main Aggregates database; UN,  
25 2016) and time series of trade data from GTAP (based on the methodology in Peters et al., 2011b  
26 ). We estimate the sector-level CO<sub>2</sub> emissions using the GTAP data and methodology, include  
27 flaring and cement emissions from CDIAC, and then scale the national totals (excluding bunker  
28 fuels) to match the emission estimates from the carbon budget. We do not provide a separate  
29 uncertainty estimate for the consumption-based emissions, but based on model comparisons and  
30 sensitivity analysis, they are unlikely to be significantly different than for the territorial emission  
31 estimates (Peters et al., 2012a).



#### 1 **2.1.4 Growth rate in emissions**

2 We report the annual growth rate in emissions for adjacent years (in percent per year) by  
3 calculating the difference between the two years and then comparing to the emissions in the first  
4 year:  $(E_{FF}(t_{0+1}) - E_{FF}(t_0)) / E_{FF}(t_0) \times 100\% \text{yr}^{-1}$ . We apply a leap-year adjustment to ensure valid  
5 interpretations of annual growth rates. This affects the growth rate by about  $0.3\% \text{yr}^{-1}$  ( $1/365$ ) and  
6 causes growth rates to go up approximately 0.3% if the first year is a leap year and down 0.3% if  
7 the second year is a leap year.

8 The relative growth rate of  $E_{FF}$  over time periods of greater than one year can be re-written using  
9 its logarithm equivalent as follows:

$$\frac{1}{E_{FF}} \frac{dE_{FF}}{dt} = \frac{d(\ln E_{FF})}{dt} \quad (2)$$

10 Here we calculate relative growth rates in emissions for multi-year periods (e.g. a decade) by  
11 fitting a linear trend to  $\ln(E_{FF})$  in Eq. (2), reported in percent per year.

#### 12 **2.1.5 Emissions projections**

13 To gain insight on emission trends for the current year (2017), we provide an assessment of global  
14 fossil fuel and industry emissions,  $E_{FF}$ , by combining individual assessments of emissions for China,  
15 USA, India (the three countries with the largest emissions), and the rest of the world. Although the  
16 EU in aggregate emits more than India, neither official forecasts nor monthly energy statistics are  
17 available for the EU as a whole. In consequence, we use GDP projections to infer the emissions for  
18 this region.

19 Our 2017 estimate for China uses: (1) estimates of coal consumption, production, imports and  
20 inventory changes from the China Coal Industry Association (CCIA) and the National Energy  
21 Agency of China (NEA) for January through June (CCIA, 2017; NEA, 2017) (2) estimated  
22 consumption of natural gas and petroleum for January through June from NEA (CCIA, 2017; NEA,  
23 2017) and (3) production of cement reported for January through August (NBS, 2017). Using these  
24 data, we estimate the change in emissions for the corresponding months in 2017 compared to  
25 2016 assuming no change in the energy and carbon content of coal for 2017. We then use a  
26 central estimate for the growth rate of the whole year that is adjusted down somewhat relative to  
27 the first half of the year, to account for a slowing trend in industrial growth observed since July  
28 and qualitative statements from the NEA saying that they expect oil and coal consumption to be



1 relatively stable for the second half of the year. The main sources of uncertainty are from  
2 inconsistencies between available data sources, incomplete data on inventory changes, the  
3 carbon content of coal and the assumptions for the behaviour for the rest of the year. These are  
4 discussed further in Sect. 3.2.1.

5 For the USA, we use the forecast of the U.S. Energy Information Administration (EIA) for emissions  
6 from fossil fuels (EIA, 2017). This is based on an energy forecasting model which is revised  
7 monthly, and takes into account heating-degree days, household expenditures by fuel type,  
8 energy markets, policies, and other effects. We combine this with our estimate of emissions from  
9 cement production using the monthly U.S. cement data from USGS for January-June, assuming  
10 changes in cement production over the first part of the year apply throughout the year. While the  
11 EIA's forecasts for current full-year emissions have on average been revised downwards, only nine  
12 such forecasts are available, so we conservatively use the full range of adjustments following  
13 revision, and additionally assume symmetrical uncertainty to give  $\pm 2.7\%$  around the central  
14 forecast.

15 For India, we use (1) coal production and sales data from the Ministry of Mines, Coal India Limited  
16 (CIL, 2017; Ministry of Mines, 2017) and Singareni Collieries Company Limited (SCCL, 2017),  
17 combined with imports data from the Ministry of Commerce and Industry (MCI, 2017) and power  
18 station stocks data from the Central Electricity Authority (CEA, 2017), (2) oil production and  
19 consumption data from the Ministry of Petroleum and Natural Gas (PPAC, 2017b), (3) natural gas  
20 production and import data from the Ministry of Petroleum and Natural Gas (PPAC, 2017a), and  
21 (4) cement production data from the Office of the Economic Advisor (OEA, 2017). The main source  
22 of uncertainty in the projection of India's emissions is the assumption of persistent growth for the  
23 rest of the year.

24 For the rest of the world, we use the close relationship between the growth in GDP and the  
25 growth in emissions (Raupach et al., 2007) to project emissions for the current year. This is based  
26 on a simplified Kaya Identity, whereby  $E_{FF}$  ( $\text{GtC yr}^{-1}$ ) is decomposed by the product of GDP ( $\text{USD yr}^{-1}$ )  
27 and the fossil fuel carbon intensity of the economy ( $I_{FF}$ ;  $\text{GtC USD}^{-1}$ ) as follows:

$$E_{FF} = \text{GDP} \times I_{FF} \quad (3)$$

28 Taking a time derivative of Equation (3) and rearranging gives:



$$\frac{1}{E_{FF}} \frac{dE_{FF}}{dt} = \frac{1}{GDP} \frac{dGDP}{dt} + \frac{1}{I_{FF}} \frac{dI_{FF}}{dt} \quad (4)$$

1 where the left-hand term is the relative growth rate of  $E_{FF}$ , and the right-hand terms are the  
2 relative growth rates of GDP and  $I_{FF}$ , respectively, which can simply be added linearly to give the  
3 overall growth rate.

4 The growth rates are reported in percent by multiplying each term by 100. As preliminary  
5 estimates of annual change in GDP are made well before the end of a calendar year, making  
6 assumptions on the growth rate of  $I_{FF}$  allows us to make projections of the annual change in  $CO_2$   
7 emissions well before the end of a calendar year. The  $I_{FF}$  is based on GDP in constant PPP  
8 (purchasing power parity) from the IEA up to 2014 (IEA/OECD, 2016) and extended using the IMF  
9 growth rates for 2015 and 2016 (IMF, 2017). Interannual variability in  $I_{FF}$  is the largest source of  
10 uncertainty in the GDP-based emissions projections. We thus use the standard deviation of the  
11 annual  $I_{FF}$  for the period 2006-2016 as a measure of uncertainty, reflecting a  $\pm 1\sigma$  as in the rest of  
12 the carbon budget. This is  $\pm 1.1\% \text{ yr}^{-1}$  for the rest of the world (global emissions minus China, USA,  
13 and India).

14 The 2017 projection for the world is made of the sum of the projections for China, USA, India, and  
15 the rest. The uncertainty is added in quadrature among the three regions. The uncertainty here  
16 reflects the best of our expert opinion.

## 17 **2.2 CO<sub>2</sub> emissions from land use, land-use change and forestry ( $E_{LUC}$ )**

18 Land-use change emissions reported here ( $E_{LUC}$ ) include  $CO_2$  fluxes from deforestation,  
19 afforestation, logging (forest degradation and harvest activity), shifting cultivation (cycle of cutting  
20 forest for agriculture, then abandoning), and regrowth of forests following wood harvest or  
21 abandonment of agriculture. Only some land management activities are included in our land-use  
22 change emissions estimates (Table 4a). Some of these activities lead to emissions of  $CO_2$  to the  
23 atmosphere, while others lead to  $CO_2$  sinks.  $E_{LUC}$  is the net sum of all anthropogenic activities  
24 considered. Our annual estimate for 1959-2016 is provided as the average of results from two  
25 bookkeeping models (Sect. 2.2.1): the estimate published by Houghton and Nassikas (2017;  
26 hereafter H&N2017) extended here to 2016, and the average of two simulations done with the  
27 BLUE model (“bookkeeping of land use emissions”; Hansis et al., 2015). In addition, we use results  
28 from DGVMs (see Sect. 2.2.3 and Table 4a), to help quantify the uncertainty in  $E_{LUC}$ , and to explore



1 the consistency of our understanding. The three methods are described below, and differences  
2 are discussed in Sect. 3.2.

### 3 **2.2.1 Bookkeeping models**

4 Land-use change CO<sub>2</sub> emissions and uptake fluxes are calculated by two bookkeeping models.  
5 Both are based on the original bookkeeping approach of Houghton (2003) that keeps track of the  
6 carbon stored in vegetation and soils before and after a land-use change (transitions between  
7 various natural vegetation types, croplands and pastures). Literature-based response curves  
8 describe decay of vegetation and soil carbon, including transfer to product pools of different  
9 lifetimes, as well as carbon uptake due to regrowth. Additionally, it represents permanent  
10 degradation of forests by lower vegetation and soil carbon stocks for secondary as compared to  
11 the primary forests and forest management such as wood harvest.

12 The bookkeeping models do not include land ecosystems' transient response to changes in  
13 climate, atmospheric CO<sub>2</sub> and other environmental factors, and the carbon densities are based on  
14 contemporary data reflecting stable environmental conditions at that time. Since carbon densities  
15 remain fixed over time in bookkeeping models, the additional sink capacity that ecosystems  
16 provide in response to CO<sub>2</sub>-fertilization and other environmental changes is not captured by these  
17 models (Pongratz et al., 2014; see Section 2.7.3).

18 The H&N and BLUE models differ in (1) computational units (country-level vs spatially explicit  
19 treatment of land-use change), (2) processes represented (see Table 4a), and (3) carbon densities  
20 assigned to vegetation and soil of each vegetation type. A notable change of H&N over the  
21 original approach by Houghton et al. (2003) used in earlier budget estimates is that no shifting  
22 cultivation or other back- and forth-transitions at a level below country level are included. Only a  
23 decline in forest area in a country as indicated by the Forest Resource Assessment of the FAO that  
24 exceeds the expansion of agricultural area as indicated by FAO is assumed to represent a  
25 concurrent expansion and abandonment of cropland. In contrast, the BLUE model includes sub-  
26 grid-scale transitions at the grid level between all vegetation types as indicated by the harmonized  
27 land-use change data (LUH2) dataset (Hurtt et al., in prep.). Furthermore, H&N assume conversion  
28 of natural grasslands to pasture, while BLUE allocates pasture proportionally on all natural  
29 vegetation that exist in a gridcell. This is one reason for generally higher emissions in BLUE. H&N  
30 add carbon emissions from peat burning based on the Global Fire Emission Database (GFED4s; van



1 der Werf et al. (2017)), and peat drainage, based on estimates by Hooijer et al. (2010) to the  
2 output of their bookkeeping model for the countries of Indonesia and Malaysia. Peat burning and  
3 emissions from the organic layers of drained peat soils, which are not captured by bookkeeping  
4 methods directly, need to be included to represent the substantially larger emissions and  
5 interannual variability due to synergies of land-use change and climate variability in South East  
6 Asia, in particular during El-Niño events. Similarly to H&N, peat burning and drainage-related  
7 emissions are also added to the BLUE estimate based on GFED4s (van der Werf et al., 2017),  
8 adding the peat burning for the GFED region of equatorial Asia, and the peat drainage for  
9 Southeast Asia from Hooijer et al (2010).

10 The two bookkeeping estimates used in this study also differ with respect to the land-cover  
11 change data used to drive the models. H&N base their estimates directly on the Forest Resource  
12 Assessment of the FAO which provides statistics on forest-cover change and management at  
13 intervals of five years (FAO, 2015). The data is based on countries' self-reporting, some of which  
14 include satellite data in more recent assessments. Changes in land cover other than forests are  
15 based on annual, national changes in cropland and pasture areas reported by the FAO Statistics  
16 Division (FAOSTAT, 2015). BLUE uses the harmonized land-use change data LUH2 (Hurtt et al., in  
17 prep.) which describes land cover change, also based on the FAO data, but downscaled at a  
18 quarter-degree spatial resolution, considering sub-grid-scale transitions between primary forest,  
19 secondary forest, cropland, pasture and rangeland. The new LUH2 data provides a new distinction  
20 between rangelands and pasture. This is implemented by assuming rangelands are treated either  
21 all as pastures, or all as natural vegetation. These two assumptions are then averaged to provide  
22 the BLUE result that is closest to the expected real value.

23 The estimate of H&N was extended here by one year (to 2016) by adding the anomaly of total  
24 peat emissions (burning and drainage) from GFED4s over the previous decade (2006-2015) to the  
25 decadal average of the bookkeeping result. A small correction to their 2015 value was also made  
26 based on the updated peat burning of GFED4s.

### 27 **2.2.2 Dynamic Global Vegetation Models (DGVMs)**

28 Land-use change CO<sub>2</sub> emissions have also been estimated using an ensemble of 12 DGVM  
29 simulations. The DGVMs account for deforestation and regrowth, the most important  
30 components of E<sub>LUC</sub>, but they do not represent all processes resulting directly from human



1 activities on land (Table 4a). All DGVMs represent processes of vegetation growth and mortality,  
2 as well as decomposition of dead organic matter associated with natural cycles, and include the  
3 vegetation and soil carbon response to increasing atmospheric CO<sub>2</sub> levels and to climate variability  
4 and change. Some models explicitly simulate the coupling of carbon and nitrogen cycles and  
5 account for atmospheric N deposition (Table 4a). The DGVMs are independent from the other  
6 budget terms except for their use of atmospheric CO<sub>2</sub> concentration to calculate the fertilization  
7 effect of CO<sub>2</sub> on plant photosynthesis.

8 The DGVMs used the HYDE land-use change data set (Klein Goldewijk et al., in press.; Klein  
9 Goldewijk et al., 2017), which provides annual, half-degree, fractional data on cropland and  
10 pasture. These data are based on annual FAO statistics of change in agricultural area available to  
11 2012 (FAOSTAT, 2015). For the years 2015 and 2016, the HYDE data were extrapolated by country  
12 for pastures and cropland separately based on the trend in agricultural area over the previous 5  
13 years. Some models also use an update of the more comprehensive harmonised land-use data set  
14 (Hurtt et al., 2011), that further includes fractional data on primary vegetation and secondary  
15 vegetation, as well as all underlying transitions between land-use states (Hurtt et al., in prep.).  
16 This new dataset is of quarter degree fractional areas of land use states and all transitions  
17 between those states, including a new wood harvest reconstruction, new representation of  
18 shifting cultivation, crop rotations, management information including irrigation and fertilizer  
19 application. The land-use states now include two different pasture/grazing types, and 5 different  
20 crop types. Wood harvest patterns are constrained with Landsat forest loss data.

21 DGVMs implement land-use change differently (e.g. an increased cropland fraction in a grid cell  
22 can either be at the expense of grassland or shrubs, or forest, the latter resulting in deforestation;  
23 land cover fractions of the non-agricultural land differ between models). Similarly, model-specific  
24 assumptions are applied to convert deforested biomass or deforested area, and other forest  
25 product pools into carbon, and different choices are made regarding the allocation of rangelands  
26 as natural vegetation or pastures.

27 The DGVM model runs were forced by either 6 hourly CRU-NCEP or by monthly CRU temperature,  
28 precipitation, and cloud cover fields (transformed into incoming surface radiation) based on  
29 observations and provided on a 0.5°x0.5° grid and updated to 2016 (Harris et al., 2014; Viovy,  
30 2016). The forcing data include both gridded observations of climate and global atmospheric CO<sub>2</sub>,



1 which change over time (Dlugokencky and Tans, 2017), and N deposition (as used in some models;  
2 Table 4a).

3 Two sets of simulations were performed with the DGVMs. The first forced initially with historical  
4 changes in land cover distribution, climate, atmospheric CO<sub>2</sub> concentration, and N deposition, and  
5 the second, as further described below, with a time-invariant preindustrial land cover distribution,  
6 allowing the models to estimate, by difference with the first simulation, the dynamic evolution of  
7 biomass and soil carbon pools in response to prescribed land-cover change. E<sub>LUC</sub> is diagnosed in  
8 each model by the difference between these two simulations. We only retain model outputs with  
9 positive E<sub>LUC</sub> during the 1990s. Using the difference between these two DGVM simulations to  
10 diagnose E<sub>LUC</sub> means the DGVMs account for the loss of additional sink capacity (around 0.3 GtC  
11 yr<sup>-1</sup>; see Section 2.7.3), while the bookkeeping models do not.

### 12 **2.2.3 Uncertainty assessment for E<sub>LUC</sub>**

13 Differences between the bookkeeping models and DGVM models originate from three main  
14 sources: the land cover change data set, the different approaches used in models, and the  
15 different processes represented (Table 4a). We examine the results from the DGVM models and  
16 of the bookkeeping method to assess the uncertainty in E<sub>LUC</sub>.

17 The E<sub>LUC</sub> estimate from the DGVMs multi-model mean is consistent with the average of the  
18 emissions from the bookkeeping models (Table 6). However there are large differences among  
19 individual DGVMs (standard deviation at around 0.5-0.6 GtC yr<sup>-1</sup>; Table 6), between the two  
20 bookkeeping models (average of 0.5 GtC yr<sup>-1</sup>), and between the current estimate of H&N and its  
21 previous model version (Houghton et al., 2012) as used in past Global Carbon Budgets (Le Quéré  
22 et al. 2016; average of 0.3 GtC yr<sup>-1</sup>). Given the large spread in new estimates we raise our  
23 assessment of uncertainty in E<sub>LUC</sub> to ±0.7 GtC yr<sup>-1</sup> (from 0.5 GtC yr<sup>-1</sup>) as a semi-quantitative  
24 measure of uncertainty for annual emissions. This reflects our best value judgment that there is at  
25 least 68% chance (±1σ) that the true land-use change emission lies within the given range, for the  
26 range of processes considered here. Prior to 1959, the uncertainty in E<sub>LUC</sub> was taken from the  
27 standard deviation of the DGVMs. We assign low confidence to the annual estimates of E<sub>LUC</sub>  
28 because of the inconsistencies among estimates and of the difficulties to quantify some of the  
29 processes in DGVMs.



#### 1 **2.2.4 Emissions projections**

2 We provide an assessment of  $E_{LUC}$  for 2017 by adding the anomaly of fire emissions in  
3 deforestation areas, including those from peat fires, from GFED4s (van der Werf et al., 2017) over  
4 the last year available. Emissions are estimated using active fire data (MCD14ML; Giglio et al.  
5 (2003)), which are available in near-real time, and correlations between those and GFED4s  
6 emissions for the 2001-2016 period for 12 the corresponding months. Emissions during January-  
7 October cover most of the fires season in the Amazon and Southeast Asia, where a large part of  
8 the global deforestation takes place.

#### 9 **2.3 Growth rate in atmospheric CO<sub>2</sub> concentration ( $G_{ATM}$ )**

##### 10 **2.3.1 Global growth rate in atmospheric CO<sub>2</sub> concentration**

11 The rate of growth of the atmospheric CO<sub>2</sub> concentration is provided by the US National Oceanic  
12 and Atmospheric Administration Earth System Research Laboratory (NOAA/ESRL; Dlugokencky  
13 and Tans, 2017), which is updated from Ballantyne et al. (2012). For the 1959-1980 period, the  
14 global growth rate is based on measurements of atmospheric CO<sub>2</sub> concentration averaged from  
15 the Mauna Loa and South Pole stations, as observed by the CO<sub>2</sub> Program at Scripps Institution of  
16 Oceanography (Keeling et al., 1976). For the 1980-2016 time period, the global growth rate is  
17 based on the average of multiple stations selected from the marine boundary layer sites with well-  
18 mixed background air (Ballantyne et al., 2012), after fitting each station with a smoothed curve as  
19 a function of time, and averaging by latitude band (Masarie and Tans, 1995). The annual growth  
20 rate is estimated by Dlugokencky and Tans (2017) from atmospheric CO<sub>2</sub> concentration by taking  
21 the average of the most recent December-January months corrected for the average seasonal  
22 cycle and subtracting this same average one year earlier. The growth rate in units of ppm yr<sup>-1</sup> is  
23 converted to units of GtC yr<sup>-1</sup> by multiplying by a factor of 2.12 GtC per ppm (Ballantyne et al.,  
24 2012).

25 The uncertainty around the annual growth rate based on the multiple stations data set ranges  
26 between 0.11 and 0.72 GtC yr<sup>-1</sup>, with a mean of 0.61 GtC yr<sup>-1</sup> for 1959-1979 and 0.19 GtC yr<sup>-1</sup> for  
27 1980-2016, when a larger set of stations were available (Dlugokencky and Tans, 2017). It is based  
28 on the number of available stations, and thus takes into account both the measurement errors  
29 and data gaps at each station. This uncertainty in decadal change is computed from the difference  
30 in concentration ten years apart based on a measurement error of 0.35 ppm. This error is based



1 on offsets between NOAA/ESRL measurements and those of the World Meteorological  
2 Organization World Data Centre for Greenhouse Gases (NOAA/ESRL, 2015) for the start and end  
3 points (the decadal change uncertainty is the  $\sqrt{(2(0.35\text{ppm})^2)}(10\text{ yr})^{-1}$  assuming that each  
4 yearly measurement error is independent).

5 We assign a high confidence to the annual estimates of  $G_{\text{ATM}}$  because they are based on direct  
6 measurements from multiple and consistent instruments and stations distributed around the  
7 world (Ballantyne et al., 2012).

8 In order to estimate the total carbon accumulated in the atmosphere since 1750 or 1870, we use  
9 an atmospheric  $\text{CO}_2$  concentration of  $277 \pm 3$  ppm or  $288 \pm 3$  ppm, respectively, based on a cubic  
10 spline fit to ice core data (Joos and Spahni, 2008). The uncertainty of  $\pm 3$  ppm (converted to  $\pm 1\sigma$ ) is  
11 taken directly from the IPCC's assessment (Ciais et al., 2013). Typical uncertainties in the growth  
12 rate in atmospheric  $\text{CO}_2$  concentration from ice core data are equivalent to  $\pm 0.1\text{--}0.15\text{ GtC yr}^{-1}$  as  
13 evaluated from the Law Dome data (Etheridge et al., 1996) for individual 20-year intervals over  
14 the period from 1870 to 1960 (Bruno and Joos, 1997).

### 15 **2.3.2 Growth rate projection**

16 We provide an assessment of  $G_{\text{ATM}}$  for 2017 based on the observed increase in atmospheric  $\text{CO}_2$   
17 concentration at the Mauna Loa station for January to September, and monthly forecasts for  
18 October to December updated from Betts et al. (2016). The forecast uses a statistical relationship  
19 between annual  $\text{CO}_2$  growth rate and sea surface temperatures (SSTs) in the Niño3.4 region. The  
20 forecast SSTs from the GLOSEA seasonal forecast model was then used to estimate monthly  $\text{CO}_2$   
21 concentrations at Mauna Loa throughout the following calendar year, assuming a stationary  
22 seasonal cycle. The forecast  $\text{CO}_2$  concentrations for January to August 2017 were close to the  
23 observations, so updating the 2017 forecast by simply averaging the observed and forecast values  
24 is considered justified. Growth at Mauna Loa is closely correlated with the global growth ( $r=0.95$ )  
25 and is used here as a proxy for global growth.

### 26 **2.4 Ocean $\text{CO}_2$ sink**

27 Estimates of the global ocean  $\text{CO}_2$  sink  $S_{\text{OCEAN}}$  are from an ensemble of global ocean  
28 biogeochemistry models (GOBM) that meet observational constraints over the 1990s (see below).  
29 We use observation-based estimates of  $S_{\text{OCEAN}}$  to provide a qualitative assessment of confidence in



1 the reported results, and to estimate the cumulative accumulation of  $S_{\text{OCEAN}}$  over the preindustrial  
2 period.

### 3 **2.4.1 Observation-based estimates**

4 We use the observational constraints assessed by IPCC of a mean ocean  $\text{CO}_2$  sink of  $2.2 \pm 0.4 \text{ GtC}$   
5  $\text{yr}^{-1}$  for the 1990s (Denman et al., 2007) to verify that the GOBMs provide a realistic assessment of  
6  $S_{\text{OCEAN}}$ . This is based on indirect observations and their spread: ocean/land  $\text{CO}_2$  sink partitioning  
7 from observed atmospheric  $\text{O}_2/\text{N}_2$  concentration trends (Manning and Keeling, 2006; updated in  
8 Keeling and Manning 2014), an oceanic inversion method constrained by ocean biogeochemistry  
9 data (Mikaloff Fletcher et al., 2006), and a method based on penetration time scale for CFCs  
10 (McNeil et al., 2003). This estimate is consistent with a range of methods (Wanninkhof et al.,  
11 2013). All GOBMs fall within 90% confidence of the observed range, or 1.6 to  $2.8 \text{ GtC yr}^{-1}$  for the  
12 1990s.

13 We use two estimates of the ocean  $\text{CO}_2$  sink and its variability based on interpolations of  
14 measurements of surface ocean fugacity of  $\text{CO}_2$  ( $p\text{CO}_2$  corrected for the non-ideal behaviour of  
15 the gas; Pfeil et al., 2013). We refer to these as  $p\text{CO}_2$ -based flux estimates. The measurements are  
16 from the Surface Ocean  $\text{CO}_2$  Atlas version 5, which is an update of version 3 (Bakker et al., 2016)  
17 and contains quality-controlled data to 2016 (see data attribution Table A2). The SOCAT v5 were  
18 mapped using a data-driven diagnostic method (Rödenbeck et al., 2013) and a combined self-  
19 organising map and feed-forward neural network (Landschützer et al., 2014). The global  $p\text{CO}_2$ -  
20 based flux estimates were adjusted to remove the preindustrial ocean source of  $\text{CO}_2$  to the  
21 atmosphere of  $0.45 \text{ GtC yr}^{-1}$  from river input to the ocean (Jacobson et al., 2007), per our  
22 definition of  $S_{\text{OCEAN}}$ . Several other flux products are available, but they show large discrepancies  
23 with observed variability that need to be resolved. Here we used the two  $p\text{CO}_2$ -based flux  
24 products that had the best fit to observations for their representation of tropical and global  
25 variability (Rödenbeck et al., 2015).

26 We further use results from two diagnostic ocean models of Khatiwala et al. (2013) and DeVries et  
27 al. (2014) to estimate the anthropogenic carbon accumulated in the ocean prior to 1959. The two  
28 approaches assume constant ocean circulation and biological fluxes over the preindustrial period,  
29 with  $S_{\text{OCEAN}}$  estimated as a response in the change in atmospheric  $\text{CO}_2$  concentration calibrated to  
30 observations. The uncertainty in cumulative uptake of  $\pm 20 \text{ GtC}$  (converted to  $\pm 1\sigma$ ) is taken directly



1 from the IPCC's review of the literature (Rhein et al., 2013), or about  $\pm 30\%$  for the annual values  
2 (Khatiwala et al., 2009).

### 3 **2.4.2 Global Ocean Biogeochemistry Models (GOBM)**

4 The ocean  $\text{CO}_2$  sink for 1959-2016 is estimated using eight GOBM (Table 4b) that meet  
5 observational constraints for the mean ocean sink in the 1990s. The GOBM represent the physical,  
6 chemical and biological processes that influence the surface ocean concentration of  $\text{CO}_2$  and thus  
7 the air-sea  $\text{CO}_2$  flux. The GOBM are forced by meteorological reanalysis and atmospheric  $\text{CO}_2$   
8 concentration data available for the entire time period, and mostly differ in the source of the  
9 atmospheric forcing data, spin up strategies, and in the resolution of the oceanic physical  
10 processes (Table 4b). GOBMs do not include the effects of anthropogenic changes in nutrient  
11 supply, which could lead to an increase of the ocean sink of up to about  $0.3 \text{ GtC yr}^{-1}$  over the  
12 industrial period (Duce et al., 2008). They also do not include the perturbation associated with  
13 changes in river organic carbon, which is discussed Sect. 2.7.

14 The ocean  $\text{CO}_2$  sink for each GOBM is no longer normalised to the observations as in previous  
15 global carbon budgets (e.g. Le Quéré et al. 2016). The normalisation was mostly intended to  
16 ensure  $S_{\text{LAND}}$  had a realistic mean value as it was previously estimated from the budget residual.  
17 With the introduction of the budget residual (Eq. 1) all terms can be estimated independently.  
18 Rather the oceanic observations are used in the selection of the GOBM, by using only the GOBM  
19 that produce an oceanic  $\text{CO}_2$  sink over the 1990s consistent with observations, as explained  
20 above.

### 21 **2.4.3 Uncertainty assessment for $S_{\text{OCEAN}}$**

22 The uncertainty around the mean ocean sink of anthropogenic  $\text{CO}_2$  was quantified by Denman et  
23 al. (2007) for the 1990s (see Sect. 2.4.1). To quantify the uncertainty around annual values, we  
24 examine the standard deviation of the GOBM ensemble, which averages between 0.2 and  $0.3 \text{ GtC}$   
25  $\text{yr}^{-1}$  during 1959-2017. We estimate that the uncertainty in the annual ocean  $\text{CO}_2$  sink is about  $\pm$   
26  $0.5 \text{ GtC yr}^{-1}$  from the combined uncertainty of the mean flux based on observations of  $\pm 0.4 \text{ GtC yr}^{-1}$   
27 <sup>1</sup> and the standard deviation across GOBMs of up to  $\pm 0.3 \text{ GtC yr}^{-1}$ , reflecting both the uncertainty  
28 in the mean sink from observations during the 1990's (Denman et al., 2007; Section 2.4.1) and in  
29 the interannual variability as assessed by GOBMs.



1 We examine the consistency between the variability of the model-based and the pCO<sub>2</sub>-based flux  
2 products to assess confidence in S<sub>OCEAN</sub>. The interannual variability of the ocean fluxes (quantified  
3 as the standard deviation) of the two pCO<sub>2</sub>-based products for 1986-2016 (where they overlap) is  
4 ± 0.35 GtC yr<sup>-1</sup> (Rödenbeck et al., 2014) and ± 0.36 GtC yr<sup>-1</sup> (Landschützer et al., 2015), compared  
5 to ± 0.27 GtC yr<sup>-1</sup> for the normalised GOBM ensemble. The standard deviation includes a  
6 component of trend and decadal variability in addition to interannual variability, and their relative  
7 influence differs across estimates. The estimates generally produce a higher ocean CO<sub>2</sub> sink during  
8 strong El Niño events. The annual pCO<sub>2</sub>-based flux products correlate with the ocean CO<sub>2</sub> sink  
9 estimated here with a correlation of  $r = 0.75$  (0.49 to 0.84 for individual GOBMs), and  $r = 0.78$   
10 (0.46 to 0.80) for the pCO<sub>2</sub>-based flux products of Rödenbeck et al. (2014) and Landschützer et al.  
11 (2015), respectively (simple linear regression), with their mutual correlation at 0.70. The  
12 agreement is better for decadal variability than for interannual variability. The use of annual data  
13 for the correlation may reduce the strength of the relationship because the dominant source of  
14 variability associated with El Niño events is less than one year. We assess a medium confidence  
15 level to the annual ocean CO<sub>2</sub> sink and its uncertainty because it is based on multiple lines of  
16 evidence, and the results are consistent in that the interannual variability in the GOBMs and data-  
17 based estimates are all generally small compared to the variability in the growth rate of  
18 atmospheric CO<sub>2</sub> concentration.

## 19 **2.5 Terrestrial CO<sub>2</sub> sink**

20 The terrestrial land sink (S<sub>LAND</sub>) is thought to be due to the combined effects of fertilisation by  
21 rising atmospheric CO<sub>2</sub> and N deposition on plant growth, as well as the effects of climate change  
22 such as the lengthening of the growing season in northern temperate and boreal areas. S<sub>LAND</sub> does  
23 not include gross land sinks directly resulting from land-use change (e.g. regrowth of vegetation)  
24 as these are part of the net land use flux (E<sub>LUC</sub>), although system boundaries make it difficult to  
25 attribute exactly CO<sub>2</sub> fluxes on land between S<sub>LAND</sub> and E<sub>LUC</sub> (Erb et al., 2013).

26 New to the 2017 Global Carbon Budget, S<sub>LAND</sub> is estimated from the multi-model mean of the  
27 DGVMs (Table 4a). As described in Sect. 2.2.3, DGVM simulations include all climate variability and  
28 CO<sub>2</sub> effects over land. The DGVMs do not include the perturbation associated with changes in  
29 river organic carbon, which is discussed Sect. 2.7. We apply three criteria for minimum DGVM  
30 realism by including only those DGVMs with (1) steady state after spin up, (2) where available, net  
31 land fluxes (S<sub>LAND</sub> – E<sub>LUC</sub>) that is a carbon sink over the 1990s between -0.3 and 2.3 GtC yr<sup>-1</sup>, within



1 90% confidence of constraints by global atmospheric and oceanic observations (Keeling and  
2 Manning, 2014; Wanninkhof et al., 2013), and (3) global  $E_{LUC}$  that is a carbon source over the  
3 1990s.

4 The standard deviation of the annual  $CO_2$  sink across the DGVMs averages to  $\pm 0.8 \text{ GtC yr}^{-1}$  for the  
5 period 1959 to 2016. We attach a medium confidence level to the annual land  $CO_2$  sink and its  
6 uncertainty because the estimates from the residual budget and averaged DGVMs match well  
7 within their respective uncertainties (Table 6).

## 8 **2.6 The atmospheric perspective**

9 The world-wide network of atmospheric measurements can be used with atmospheric inversion  
10 methods to constrain the location of the combined total surface  $CO_2$  fluxes from all sources,  
11 including fossil and land-use change emissions and land and ocean  $CO_2$  fluxes. The inversions  
12 assume  $E_{FF}$  to be well known, and they solve for the spatial and temporal distribution of land and  
13 ocean fluxes from the residual gradients of  $CO_2$  between stations that are not explained by  
14 emissions.

15 Three atmospheric inversions (Table 4c) used atmospheric  $CO_2$  data to the end of 2016 (including  
16 preliminary values in some cases) to infer the spatio-temporal  $CO_2$  flux field. We focus here on the  
17 largest and most consistent sources of information (namely the total  $CO_2$  flux over land regions,  
18 and the distribution of the total land and ocean  $CO_2$  fluxes for the mid-high latitude northern  
19 hemisphere (30°N-90°N), Tropics (30°S-30°N) and mid-high latitude region of the southern  
20 hemisphere (30°S-90°S)), and use these estimates to comment on the consistency across various  
21 data streams and process-based estimates.

### 22 **Atmospheric inversions**

23 The three inversion systems used in this release are the CarbonTracker Europe (CTE; van der Laan-  
24 Luijkx et al., 2017), the Jena CarboScope (Rödenbeck, 2005), and CAMS (Chevallier et al., 2005).  
25 See Table 4c for version numbers. The three inversions are based on the same Bayesian inversion  
26 principles that interpret the same, for the most part, observed time series (or subsets thereof),  
27 but use different methodologies (Table 4c). These differences mainly concern the selection of  
28 atmospheric  $CO_2$  data, the used prior fluxes, spatial breakdown (i.e. grid size), assumed  
29 correlation structures, and mathematical approach. The details of these approaches are



1 documented extensively in the references provided above. Each system uses a different transport  
2 model, which was demonstrated to be a driving factor behind differences in atmospheric-based  
3 flux estimates, and specifically their distribution across latitudinal bands (Stephens et al., 2007).

4 The three inversions use atmospheric CO<sub>2</sub> observations from various flask and in situ networks, as  
5 detailed in Table 4c. They prescribe global E<sub>FF</sub>, which is scaled to the present study for CAMS and  
6 CTE, while CarboScope uses CDIAC extended after 2013 using the emission growth rate of the  
7 present study. Inversion results for the sum of natural ocean and land fluxes (Fig. 8) are more  
8 constrained in the Northern hemisphere (NH) than in the Tropics, because of the higher  
9 measurement stations density in the NH. Results from atmospheric inversions, similar to the  
10 pCO<sub>2</sub>-based ocean flux products, need to be corrected for the river fluxes. The atmospheric  
11 inversions provide new information on the regional distribution of fluxes.

## 12 **2.7 Processes not included in the global carbon budget**

13 The contribution of anthropogenic CO and CH<sub>4</sub> to the global carbon budget has been partly  
14 neglected in Eq. 1 and is described in Sect. 2.7.1. The contribution of anthropogenic changes in  
15 river fluxes is conceptually included in Eq. 1 in S<sub>OCEAN</sub> and in S<sub>LAND</sub>, but it is not represented in the  
16 process models used to quantify these fluxes. This effect is discussed in Sect. 2.7.2. Similarly, the  
17 loss of additional sink capacity from reduced forest cover is missing in the combination of  
18 approaches used here to estimate both land fluxes (E<sub>LUC</sub> and S<sub>LAND</sub>) and its potential effect is  
19 discussed and quantified in Sect. 2.7.3.

### 20 **2.7.1 Contribution of anthropogenic CO and CH<sub>4</sub> to the global carbon budget**

21 Anthropogenic emissions of CO and CH<sub>4</sub> to the atmosphere are eventually oxidized to CO<sub>2</sub> and  
22 thus are part of the global carbon budget. These contributions are omitted in Eq. (1), but an  
23 attempt is made in this section to estimate their magnitude, and identify the sources of  
24 uncertainty. Anthropogenic CO emissions are from incomplete fossil fuel and biofuel burning and  
25 deforestation fires. The main anthropogenic emissions of fossil CH<sub>4</sub> that matter for the global  
26 carbon budget are the fugitive emissions of coal, oil and gas upstream sectors (see below). These  
27 emissions of CO and CH<sub>4</sub> contribute a net addition of fossil carbon to the atmosphere.

28 In our estimate of E<sub>FF</sub> we assumed (Sect. 2.1.1) that all the fuel burned is emitted as CO<sub>2</sub>, thus CO  
29 anthropogenic emissions and their atmospheric oxidation into CO<sub>2</sub> within a few months are



1 already counted implicitly in  $E_{FF}$  and should not be counted twice (same for  $E_{LUC}$  and  
2 anthropogenic CO emissions by deforestation fires). Anthropogenic emissions of fossil  $CH_4$  are not  
3 included in  $E_{FF}$ , because these fugitive emissions are not included in the fuel inventories. Yet they  
4 contribute to the annual  $CO_2$  growth rate after  $CH_4$  gets oxidized into  $CO_2$ . Anthropogenic  
5 emissions of fossil  $CH_4$  represent 15% of total  $CH_4$  emissions (Kirschke et al., 2013) that is  $0.061$   
6  $GtC\ yr^{-1}$  for the past decade. Assuming steady state, these emissions are all converted to  $CO_2$  by  
7 OH oxidation, and thus explain  $0.06\ GtC\ yr^{-1}$  of the global  $CO_2$  growth rate in the past decade, or  
8  $0.07\text{-}0.1\ GtC\ yr^{-1}$  using the higher  $CH_4$  emissions reported recently (Schwietzke et al., 2016).  
9 Other anthropogenic changes in the sources of CO and  $CH_4$  from wildfires, biomass, wetlands,  
10 ruminants or permafrost changes are similarly assumed to have a small effect on the  $CO_2$  growth  
11 rate.

## 12 **2.7.2 Anthropogenic carbon fluxes in the land to ocean aquatic continuum**

13 The approach used to determine the global carbon budget refers to the mean, variations, and  
14 trends in the perturbation of  $CO_2$  in the atmosphere, referenced to the preindustrial era. Carbon is  
15 continuously displaced from the land to the ocean through the land-ocean aquatic continuum  
16 (LOAC) comprising freshwaters, estuaries and coastal areas (Bauer et al., 2013; Regnier et al.,  
17 2013). A significant fraction of this lateral carbon flux is entirely ‘natural’ and is thus a steady state  
18 component of the preindustrial carbon cycle. We account for this preindustrial flux where  
19 appropriate in our study. However, changes in environmental conditions and land use change  
20 have caused an increase in the lateral transport of carbon into the LOAC – a perturbation that is  
21 relevant for the global carbon budget presented here.

22 The results of the analysis of Regnier et al. (2013) can be summarized in two points of relevance  
23 for the anthropogenic  $CO_2$  budget. First, the anthropogenic perturbation has increased the  
24 organic carbon export from terrestrial ecosystems to the hydrosphere at a rate of  $1.0 \pm 0.5\ GtC\ yr^{-1}$ ,  
25 mainly owing to enhanced carbon export from soils. Second, this exported anthropogenic  
26 carbon is partly respired through the LOAC, partly sequestered in sediments along the LOAC and  
27 to a lesser extent, transferred in the open ocean where it may accumulate. The increase in storage  
28 of land-derived organic carbon in the LOAC and open ocean combined is estimated by Regnier et  
29 al. (2013) at  $0.65 \pm 0.35\ GtC\ yr^{-1}$ . We do not attempt to incorporate the changes in LOAC in our  
30 study.



1 The inclusion of freshwater fluxes of anthropogenic CO<sub>2</sub> affects the estimates of, and partitioning  
2 between, S<sub>LAND</sub> and S<sub>OCEAN</sub> in Eq. (1) in complementary ways, but does not affect the other terms.  
3 This effect is not included in the GOBMs and DGVMs used in our global carbon budget analysis  
4 presented here.

### 5 **2.7.3 Loss of additional sink capacity**

6 The DGVM simulations now used to estimate S<sub>LAND</sub> are carried out with a fixed preindustrial land-  
7 cover. Hence, they overestimate the land sink by ignoring historical changes in vegetation cover  
8 due to land use and how this affected the global terrestrial biosphere's capacity to remove CO<sub>2</sub>  
9 from the atmosphere. Historical land-cover change was dominated by transitions from vegetation  
10 types that can provide a large sink per area unit (typically, forests) to others less efficient in  
11 removing CO<sub>2</sub> from the atmosphere (typically, croplands). The resultant decrease in land sink,  
12 called the 'loss of sink capacity', is calculated as the difference between the actual land sink under  
13 changing land-cover and the counter-factual land sink under preindustrial land-cover.

14 Here, we provide a quantitative estimate of this term to be used in the discussion. Our estimate  
15 uses the compact Earth system model OSCAR (Gasser et al., 2017) whose land carbon cycle  
16 component is designed to emulate the behaviour of TRENDY and CMIP5 complex models. We use  
17 OSCAR v2.2.1 (an update of v2.2 in which minor changes) in a probabilistic setup identical to the  
18 one of Arneeth et al. (2017) but with a Monte Carlo ensemble of 2000 simulations. For each, we  
19 calculate separately S<sub>LAND</sub> and the loss of additional sink capacity. We then constrain the ensemble  
20 by weighting each member to obtain a distribution of cumulative S<sub>LAND</sub> over 1850-2005 close to  
21 the DGVMs used here. From this ensemble, we estimate a loss of additional sink capacity of  $0.4 \pm$   
22  $0.3 \text{ GtC yr}^{-1}$  on average over 2005-2014, and by extrapolation of  $20 \pm 15 \text{ GtC}$  accumulated  
23 between 1870 and 2016.

## 24 **3 Results**

### 25 **3.1 Global carbon budget mean and variability for 1959 – 2016**

26 The global carbon budget averaged over the last half-century is shown in Fig. 3. For this time  
27 period, 82% of the total emissions (E<sub>FF</sub> + E<sub>LUC</sub>) were caused by fossil fuels and industry, and 18% by  
28 land-use change. The total emissions were partitioned among the atmosphere (45%), ocean (23%)  
29 and land (32%). All components except land-use change emissions have grown since 1959, with



1 important interannual variability in the growth rate in atmospheric CO<sub>2</sub> concentration and in the  
2 land CO<sub>2</sub> sink (Fig. 4), and some decadal variability in all terms (Table 7).

### 3 **3.1.1 CO<sub>2</sub> emissions**

4 Global CO<sub>2</sub> emissions from fossil fuels and industry have increased every decade from an average  
5 of  $3.1 \pm 0.2$  GtC yr<sup>-1</sup> in the 1960s to an average of  $9.4 \pm 0.5$  GtC yr<sup>-1</sup> during 2007-2016 (Table 7 and  
6 Fig. 5). The growth rate in these emissions decreased between the 1960s and the 1990s, from  
7  $4.5\%$  yr<sup>-1</sup> in the 1960s (1960-1969),  $2.8\%$  yr<sup>-1</sup> in the 1970s (1970-1979),  $1.9\%$  yr<sup>-1</sup> in the 1980s  
8 (1980-1989), and to  $1.1\%$  yr<sup>-1</sup> in the 1990s (1990-1999). After this period, the growth rate began  
9 increasing again in the 2000s at an average growth rate of  $3.3\%$  yr<sup>-1</sup>, decreasing to  $1.8\%$  yr<sup>-1</sup> for  
10 the last decade (2007-2016), and to  $+0.4\%$  yr<sup>-1</sup> during 2014-2016.

11 In contrast, CO<sub>2</sub> emissions from land-use change have remained relatively constant at around  $1.3$   
12  $\pm 0.7$  GtC yr<sup>-1</sup> over the past half-century, in agreement with the DGVM ensemble of models.  
13 However, there is no agreement on the trend over the full period, with two bookkeeping models  
14 suggesting opposite trends and no coherence among DGVMs (Fig. 6).

### 15 **3.1.2 Partitioning among the atmosphere, ocean and land**

16 The growth rate in atmospheric CO<sub>2</sub> level increased from  $1.7 \pm 0.1$  GtC yr<sup>-1</sup> in the 1960s to  $4.7 \pm$   
17  $0.1$  GtC yr<sup>-1</sup> during 2007-2016 with important decadal variations (Table 7). Both ocean and land  
18 CO<sub>2</sub> sinks increased roughly in line with the atmospheric increase, but with significant decadal  
19 variability on land (Table 7), and possibly in the ocean (Fig. 7).

20 The ocean CO<sub>2</sub> sink increased from  $1.0 \pm 0.5$  GtC yr<sup>-1</sup> in the 1960s to  $2.4 \pm 0.5$  GtC yr<sup>-1</sup> during 2007-  
21 2016, with interannual variations of the order of a few tenths of GtC yr<sup>-1</sup> generally showing an  
22 increased ocean sink during large El Niño events (i.e. 1997-1998) (Fig. 7; Rödenbeck et al., 2014).  
23 Note the lower ocean sink estimate compared to previous global carbon budget releases is due to  
24 the fact that ocean models are no longer normalised to observations. Although there is some  
25 coherence among the GOBMs and pCO<sub>2</sub>-based flux products regarding the mean, there is poor  
26 agreement for interannual variability and the ocean models underestimate decadal variability  
27 (Sect. 2.4.3 and Fig. 7, also see new data-based decadal estimate of DeVries et al. (2017)).

28 The terrestrial CO<sub>2</sub> sink increased from  $1.4 \pm 0.7$  GtC yr<sup>-1</sup> in the 1960s to  $3.0 \pm 0.8$  GtC yr<sup>-1</sup> during  
29 2007-2016, with important interannual variations of up to 2 GtC yr<sup>-1</sup> generally showing a



1 decreased land sink during El Niño events, overcompensating the increase in ocean sink and  
2 responsible for the enhanced growth rate in atmospheric CO<sub>2</sub> concentration during El Niño events  
3 (Fig. 6). The larger land CO<sub>2</sub> sink during 2007-2016 compared to the 1960s is reproduced by all the  
4 DGVMs in response to combined atmospheric CO<sub>2</sub> increase, climate and variability, consistent  
5 with constraints from the other budget terms (Table 6).

6 The total CO<sub>2</sub> fluxes on land ( $S_{\text{LAND}} - E_{\text{LUC}}$ ) constrained by the atmospheric inversions show in  
7 general very good agreement with the global budget estimate, as expected given the strong  
8 constrains of  $G_{\text{ATM}}$  and the small relative uncertainty assumed on  $S_{\text{OCEAN}}$  and  $E_{\text{FF}}$  by inversions. The  
9 total land flux is of similar magnitude for the decadal average, with estimates for 2007-2016 from  
10 the three inversions of 1.8, 1.4 and 2.3 GtC yr<sup>-1</sup> compared to  $1.7 \pm 0.7$  GtC yr<sup>-1</sup> from the DGVMs  
11 and  $2.3 \pm 0.7$  GtC yr<sup>-1</sup> for the total flux computed with the carbon budget constraints (Table 6).

### 12 **3.1.3 Budget imbalance**

13 The carbon budget imbalance ( $B_{\text{IM}}$ ; Eq. 1) quantifies the mismatch between the estimated total  
14 emissions and the estimated changes in the atmosphere, land and ocean reservoirs. The mean  
15 budget imbalance from 1959 to 2016 is very small ( $0.07$  GtC yr<sup>-1</sup>) and shows no trend over the full  
16 time series. Although the process models (GOBMs and DGVMs) have been selected to match  
17 observational constraints in the 1990s, they are independent of the estimated emissions from  
18 fossil fuels and industry, and therefore the near-zero mean and trend in the budget imbalance is  
19 an indirect evidence of a coherent community understanding of the emissions and their  
20 partitioning on those time scales (Fig. 4). However, the budget imbalance shows substantial  
21 variability of the order of  $\pm 1$  GtC yr<sup>-1</sup>, particularly over semi-decadal time scales, although most of  
22 the variability is within the uncertainty of the estimates. The imbalance during the 1960s, early  
23 1990s, and in the last decade, suggest that either the emissions were overestimated or the sinks  
24 were underestimated during these periods. The reverse is true for the 1970s and around 1995-  
25 2000 (Fig. 3).

26 We cannot attribute the cause of the variability in the budget imbalance with our analysis, only to  
27 note that the budget imbalance is unlikely to be explained by errors or biases in the emissions  
28 alone because of its large semi-decadal variability component, a variability that is untypical of  
29 emissions (Fig. 4). Errors in  $S_{\text{LAND}}$  and  $S_{\text{OCEAN}}$  are more likely to be the main cause for the budget  
30 imbalance. For example, underestimation of the  $S_{\text{LAND}}$  by DGVMs has been reported following the



1 eruption of Mount Pinatubo in 1991 possibly due to missing responses to changes in diffuse  
2 radiation (Mercado et al., 2009), and DGVMs are suspected to overestimate the land sink in  
3 response to the wet decade of the 1970s (Sitch et al., 2003). Decadal and semi-decadal variability  
4 in the ocean sink has been also reported recently (DeVries et al., 2017; Landschützer et al., 2015),  
5 with the pCO<sub>2</sub>-based ocean flux products suggesting a smaller than expected ocean CO<sub>2</sub> sink in the  
6 1990s and a larger than expected sink in the 2000s (Fig. 7), possibly caused by changes in ocean  
7 circulation (DeVries et al., 2017) not captured in coarse resolution GOBMs used here (Dufour et  
8 al., 2013).

#### 9 **3.1.4 Regional distribution**

10 Fig 8 shows the partitioning of the total surface fluxes excluding emissions from fossil fuels and  
11 industry ( $S_{\text{LAND}} + S_{\text{OCEAN}} - E_{\text{LUC}}$ ) according to the multi-model average of the process models in the  
12 ocean and on land (GOBMs and DGVMs), and to the three atmospheric inversions. The total  
13 surface fluxes provide information on the regional distribution of those fluxes by latitude bands  
14 (Fig. 8). The global mean CO<sub>2</sub> fluxes from process models for 2007-2016 is  $4.1 \pm 1.0 \text{ GtC yr}^{-1}$ . This is  
15 comparable to the fluxes of  $4.6 \pm 0.5 \text{ GtC yr}^{-1}$  inferred from the remainder of the carbon budget  
16 ( $E_{\text{FF}} - G_{\text{ATM}}$  in Equation 1; Table 7) within their respective uncertainties. The total CO<sub>2</sub> fluxes from  
17 the three inversions range between 4.1 and 5.0 GtC yr<sup>-1</sup>, consistent with the carbon budget as  
18 expected from the constraints on the inversions.

19 In the South (south of 30°S), the atmospheric inversions and process models all suggest a CO<sub>2</sub> sink  
20 for 2007-2016 around 1.3-1.4 GtC yr<sup>-1</sup> (Fig. 8), although interannual to decadal variability is not  
21 fully consistent across methods. The interannual variability in the South is low because of the  
22 dominance of ocean area with low variability compared to land areas.

23 In the Tropics (30°S-30°N), both the atmospheric inversions and process models suggest the  
24 carbon balance in this region is close to neutral on average over the past decade, with fluxes for  
25 2007-2016 ranging between -0.5 and +0.5 GtC yr<sup>-1</sup>. Both the process models and the inversions  
26 consistently allocate more year-to-year variability of CO<sub>2</sub> fluxes to the Tropics compared to the  
27 North (north of 30°N; Fig. 8), this variability being dominated by land fluxes.

28 In the North (north of 30°N), the inversions and process models are not in agreement on the  
29 magnitude of the CO<sub>2</sub> sink, with the ensemble mean of the process models suggesting a total



1 northern hemisphere sink for 2007-2016 of  $2.3 \pm 0.6$  GtC yr<sup>-1</sup>, below the estimates from the three  
2 inversions that estimate a sink of 2.7, 3.0 and 4.1 GtC yr<sup>-1</sup> (Fig. 8). The mean difference can only  
3 partly be explained by the influence of river fluxes, which is seen by the inversions but not  
4 included in the process models; this flux in the Northern Hemisphere would be less than 0.45 GtC  
5 yr<sup>-1</sup> because only the anthropogenic contribution to river fluxes needs to be accounted for. The  
6 CTE and Jena CarboScope inversions are within the one standard deviation of the process models  
7 for the mean sink during their overlap period, while the CAMS inversion gives a higher sink in the  
8 North than the process models, and a correspondingly higher source in the Tropics.

9 Differences between CTE, CAMS, and Jena CarboScope may be related e.g. to differences in  
10 interhemispheric mixing time of their transport models, and other inversion settings (Table 4c).  
11 Separate analysis has shown that the influence of the chosen prior land and ocean fluxes is minor  
12 compared to other aspects of each inversion. In comparison to the previous global carbon budget  
13 publication, the fossil fuel inputs for CarboScope changed to lower emissions in the North  
14 compared to CTE and CAMS, resulting in a smaller Northern sink for CarboScope compared to the  
15 previous estimate. Differences between the mean fluxes of CAMS in the North and the ensemble  
16 of process models cannot be simply explained. They could either reflect a bias in this inversion or  
17 missing processes or biases in the process models, such as the lack of adequate parameterizations  
18 for forest management in the North and for forest degradation emissions in Tropics for the  
19 DGVMs. The estimated contribution of the North and its uncertainty from process models is  
20 sensitive both to the ensemble of process models used and to the specifics of each inversion.

### 21 **3.2 Global carbon budget for the last decade (2007 – 2016)**

22 The global carbon budget averaged over the last decade (2007-2016) is shown in Fig. 2. For this  
23 time period, 88% of the total emissions ( $E_{FF} + E_{LUC}$ ) were from fossil fuels and industry ( $E_{FF}$ ), and  
24 12% from land-use change ( $E_{LUC}$ ). The total emissions were partitioned among the atmosphere  
25 (44%), ocean (22%) and land (28%), with a remaining unattributed budget imbalance (5%).

#### 26 **3.2.1 CO<sub>2</sub> emissions**

27 Global CO<sub>2</sub> emissions from fossil fuels and industry grew at a rate of 1.8% yr<sup>-1</sup> for the last decade  
28 (2007-2016), slowing down to +0.4% yr<sup>-1</sup> during 2014-2016. China's emissions increased by +3.8%  
29 yr<sup>-1</sup> on average (increasing by +1.7 GtC yr<sup>-1</sup> during the 10-year period) dominating the global



1 trends, followed by India's emissions increase by  $+5.8\% \text{ yr}^{-1}$  (increasing by  $+0.30 \text{ GtC yr}^{-1}$ ), while  
2 emissions decreased in EU28 by  $2.2\% \text{ yr}^{-1}$  (decreasing by  $-0.23 \text{ GtC yr}^{-1}$ ), and in the USA by  $1.0\% \text{ yr}^{-1}$   
3 <sup>1</sup> (decreasing by  $-0.19 \text{ GtC yr}^{-1}$ ). In the past decade, emissions from fossil fuels and industry  
4 decreased significantly (at the 95% level) in 26 countries. 22 of these countries had positive  
5 growth in GDP over the same time period, representing 20% of global emissions (Austria, Belgium,  
6 Bulgaria, Czech Republic, Denmark, France, Hungary, Ireland, Latvia, Lithuania, Luxembourg,  
7 Macedonia, Malta, Netherlands, Poland, Romania, Serbia, Slovakia, Sweden, Switzerland, United  
8 Kingdom, USA).

9 In contrast, there is no apparent trend in CO<sub>2</sub> emissions from land-use change (Fig. 6), though the  
10 data is very uncertain.

### 11 **3.2.2 Partitioning among the atmosphere, ocean and land**

12 The growth rate in atmospheric CO<sub>2</sub> concentration was initially constant and then increased  
13 during the later part of the decade 2007-2016, reflecting a similar constant level followed by a  
14 decrease in the land sink, albeit with large interannual variability (Fig. 4). During the same period,  
15 the ocean CO<sub>2</sub> sink appears to have intensified, an effect which is particularly apparent in the  
16 pCO<sub>2</sub>-based flux products (Fig. 7) and is thought to originate at least in part in the Southern Ocean  
17 (Landschützer et al., 2015).

### 18 **3.2.3 Budget imbalance**

19 The budget imbalance was  $0.6 \text{ GtC yr}^{-1}$  on average over 2007-2016. Although the uncertainties are  
20 large in each term, the sustained imbalance over a decade suggests an overestimation of the  
21 emissions and/or an underestimation of the sinks. Such a large imbalance is unlikely to originate  
22 from the emissions alone because it would indicate sustained bias in emissions over a 10-year  
23 period that is as large as the 1-sigma uncertainty. An origin in the land and/or ocean sink is more  
24 likely, given the large variability of the land sink and the suspected underestimation of decadal  
25 variability in the ocean sink.



### 1 **3.3 Global carbon budget for year 2016**

#### 2 **3.3.1 CO<sub>2</sub> emissions**

3 Preliminary global CO<sub>2</sub> emissions from fossil fuels and industry based on BP energy statistics are  
4 for emissions remaining nearly constant between 2015 and 2016 at  $9.9 \pm 0.5$  GtC in 2016 (Fig. 5),  
5 distributed among coal (40%), oil (34%), gas (19%), cement (5.6%) and gas flaring (0.7%).  
6 Compared to the previous year, emissions from coal decreased by  $-1.7\%$ , while emissions from  
7 oil, gas, and cement increased by  $1.5\%$ ,  $1.5\%$ , and  $1.0\%$ , respectively. All growth rates presented  
8 are adjusted for leap year, unless stated otherwise.

9 Emissions in 2016 were  $0.2\%$  higher than in 2015, continuing the low growth trends observed in  
10 2014 and 2015. This growth rate is as projected in Le Quéré et al. (2016) based on national  
11 emissions projections for China and the USA, and projections of gross domestic product corrected  
12 for I<sub>FF</sub> trends for the rest of the world. The specific projection for 2016 for China made last year of  
13  $-0.5\%$  (range of  $-3.8\%$  to  $+1.3\%$ ) is very close to the realised growth rate of  $-0.3\%$ . Similarly, the  
14 projected growth for the US of  $-1.7\%$  (range of  $-4.0\%$  to  $+0.6\%$ ) is very close to the realised  
15 growth rate of  $-2.1\%$ , and the projected growth for the rest of the world (ROW) of  $+1.0\%$  (range  
16 of  $-0.4\%$  to  $2.5\%$ ) matches the realised rate of  $1.3\%$ .

17 In 2016, the largest absolute contributions to global CO<sub>2</sub> emissions were from China (28%), the  
18 USA (15%), the EU (28 member states; 10%), and India (6.7%). The percentages are the fraction of  
19 the global emissions including bunker fuels (3.1%). These four regions account for 59% of global  
20 CO<sub>2</sub> emissions. Growth rates for these countries from 2015 to 2016 were  $-0.3\%$  (China),  $-2.1\%$   
21 (USA),  $-0.3\%$  (EU28), and  $+4.5\%$  (India). The per-capita CO<sub>2</sub> emissions in 2016 were  $1.1$  tC person<sup>-1</sup>  
22 yr<sup>-1</sup> for the globe, and were 4.5 (USA), 2.0 (China), 1.9 (EU28) and 0.5 (India) tC person<sup>-1</sup> yr<sup>-1</sup> for the  
23 four highest emitting countries (Fig. 5e).

24 Territorial emissions in Annex B countries (developed countries as per the Kyoto Protocol which  
25 initially had binding mitigation targets) decreased by  $-0.2\%$  yr<sup>-1</sup> on average during 1990-2015.  
26 Trends observed for consumption emissions were less monotonic, with  $0.7\%$  yr<sup>-1</sup> growth over  
27 1990-2007 and a  $-1.2\%$  yr<sup>-1</sup> decrease over 2007-2015 (Fig. 5c). In non-Annex B countries  
28 (emerging economies and less developed countries as per the Kyoto Protocol with no binding  
29 mitigation commitments) territorial emissions grew at  $4.6\%$  yr<sup>-1</sup> during 1990-2015, while  
30 consumption emissions grew at  $4.5\%$  yr<sup>-1</sup>. In 1990, 65% of global territorial emissions were



1 emitted in Annex B countries (32% in non-Annex B, and 2% in bunker fuels used for international  
2 shipping and aviation), while in 2015 this had reduced to 37% (60% in non-Annex B, and 3% in  
3 bunker fuels). For consumption emissions, this split was 68% in 1990 and 42% in 2015 (32% to  
4 58% in non-Annex B). The difference between territorial and consumption emissions (the net  
5 emission transfer via international trade) from non-Annex B to Annex B countries has increased  
6 from near zero in 1990 to  $0.3 \text{ GtC yr}^{-1}$  around 2005 and remained relatively stable afterwards until  
7 the last year available (2015; Fig. 5). The increase in net emission transfers of  $0.28 \text{ GtC yr}^{-1}$   
8 between 1990 and 2015 compares with the emission reduction of  $0.5 \text{ GtC yr}^{-1}$  in Annex B  
9 countries. These results show the importance of net emission transfer via international trade from  
10 non-Annex B to Annex B countries, and the stabilisation of emissions transfer when averaged over  
11 Annex B countries during the past decade. In 2015, the biggest emitters from a consumption  
12 perspective were China (23% of the global total), USA (16%), EU28 (12%), and India (6%).

13 The global  $\text{CO}_2$  emissions from land-use change are estimated as  $1.3 \pm 0.5 \text{ GtC}$  in 2016, as for the  
14 previous decade but with low confidence in the annual change.

### 15 **3.3.2 Partitioning among the atmosphere, ocean and land**

16 The growth rate in atmospheric  $\text{CO}_2$  concentration was  $6.1 \pm 0.2 \text{ GtC}$  in 2016 ( $2.89 \pm 0.09 \text{ ppm}$ ; Fig.  
17 4; Dlugokencky and Tans, 2017). This is well above the 2007-2016 average of  $4.7 \pm 0.1 \text{ GtC yr}^{-1}$  and  
18 reflects the large interannual variability in the growth rate of atmospheric  $\text{CO}_2$  concentration  
19 associated with El Niño and La Niña events.

20 The estimated ocean  $\text{CO}_2$  sink was  $2.6 \pm 0.5 \text{ GtC yr}^{-1}$  in 2016, only marginally above 2015 according  
21 to the average of the ocean models but with large differences among estimates (Fig. 7).

22 The terrestrial  $\text{CO}_2$  sink from the model ensemble was  $2.7 \pm 1.0 \text{ GtC}$  in 2016, near the decadal  
23 average (Fig. 4) and consistent with constraints from the rest of the budget (Table 6).

24 The budget imbalance was  $-0.3 \text{ GtC}$  in 2016, indicating a small overestimation of the emissions  
25 and/or underestimation of the sink for that year, with large uncertainties.

### 26 **3.4 Global carbon budget projection for year 2017**

#### 27 **3.4.1 $\text{CO}_2$ emissions**

28 Emissions from fossil fuels and industry ( $E_{\text{FF}}$ ) for 2017 are projected to increase by +2.0% (range of  
29 0.8% to +3.0%; Table 8; (Jackson et al., 2017; Peters et al., 2017)). Our method contains several



1 assumptions that could influence the estimate beyond the given range, and as such, it has an  
2 indicative value only. Within the given assumptions, global emissions would increase to  $10.0 \pm 0.5$   
3 GtC ( $36.8 \pm 1.8$  GtCO<sub>2</sub>) in 2017.

4 For China, the expected change based on available data as of 19 September 2017 (see Sect. 2.1.4)  
5 is for an increase in emissions of +3.5% (range of +0.7% to +5.4%) in 2017 compared to 2016. This  
6 is based on estimated growth in coal (+3%; the main fuel source in China), oil (+5.0%) and natural  
7 gas (+11.7%) consumption and a decline in cement production (−0.5%). The uncertainty range  
8 considers the spread between different data sources, and variances of typical revisions of Chinese  
9 data over time. The uncertainty in the growth rate of coal consumption also reflects uncertainty in  
10 the evolution of energy density and carbon content of coal.

11 For the USA, the EIA emissions projection for 2017 combined with cement data from USGS gives a  
12 decrease of −0.4 % (range of −2.7 to +1.9 %) compared to 2016.

13 For India, our projection for 2017 gives an increase of +2.0% (range of 0.2% to +3.8%) over 2016.

14 For the rest of the world (including EU28), the expected growth for 2017 is +1.9% (range of 0.3%  
15 to +3.4%). This is computed using the GDP projection for the world excluding China, USA, and  
16 India of 3.0% made by the IMF (IMF, 2017) and a decrease in  $I_{FF}$  of  $-1.1\% \text{ yr}^{-1}$  which is the average  
17 from 2007-2016. The uncertainty range is based on the standard deviation of the interannual  
18 variability in  $I_{FF}$  during 2007-2016 of  $\pm 1.0\% \text{ yr}^{-1}$  and our estimate of uncertainty in the IMF's GDP  
19 forecast of  $\pm 0.5\%$ . Applying the method to the EU28 individually would give a projection of −0.2%  
20 (range of −2.0% to +1.6%) for EU28 and +2.3% (range of +0.5% to +4.0%) for the remaining  
21 countries, though the uncertainties grow with the level of disaggregation.

22 Emissions from land-use change ( $E_{LUC}$ ) for 2017 are projected to remain in line with or slightly  
23 lower than their 2016 level of 1.3 GtC, based on active fire detections by October.

24 **3.4.2 Partitioning among the atmosphere, ocean and land**

25 The 2017 growth in atmospheric CO<sub>2</sub> concentration ( $G_{ATM}$ ) is projected to be 5.3 GtC with  
26 uncertainty around  $\pm 1$  GtC ( $2.5 \pm 0.5$  ppm). Combining projected  $E_{FF}$ ,  $E_{LUC}$  and  $G_{ATM}$  suggests a  
27 combined land and ocean sink ( $S_{LAND} + S_{OCEAN}$ ) of about 6 GtC for 2017. Although each term has  
28 large uncertainty, the oceanic sink  $S_{OCEAN}$  has generally low interannual variability and is likely to  
29 remain close to its 2016 value of around 2.6 GtC, leaving a rough estimated land sink  $S_{LAND}$  of



1 around 3.4 GtC, near its decadal average (Table 6). This behaviour of the sink is expected due to  
2 the El Niño-neutral conditions that prevailed during 2017, in stark contrast to the strong El Niño  
3 conditions in 2015 and 2016 that reduced the land sink.

#### 4 **3.5 Cumulative sources and sinks**

5 Cumulative historical sources and sinks have been revised compared to the previous global carbon  
6 budgets. This version of the global carbon budget uses two updated bookkeeping models instead  
7 of one bookkeeping model only, uses two ocean sink data-products instead of one data-product  
8 only, and uses multiple DGVMs for the land sink instead of deriving the land sink from the residual  
9 of the other terms. As a result of these methodological changes, the cumulative emissions and  
10 their partitioning is significantly larger (by about 50 GtC) than our previous estimates. This large  
11 difference highlights the uncertainty in reconstructing historical emission sources and sinks, and  
12 this is noted through the large uncertainty associated with each term.

13 Cumulative fossil fuel and industry emissions for 1870-2016 were  $420 \pm 20$  GtC for  $E_{FF}$  and, with  
14 the revised bookkeeping models,  $180 \pm 60$  GtC for  $E_{LUC}$  (Table 9), for a total of  $600 \pm 65$  GtC. The  
15 cumulative emissions from  $E_{LUC}$  are particularly uncertain, with large spread among individual  
16 estimates of 135 GtC (Houghton) and 225 GtC (BLUE) for the two bookkeeping models and a range  
17 of 70 to 230 GtC for the twelve DGVMs. These estimates are consistent with indirect constraints  
18 from biomass observations (Li et al., 2017), but given the large spread a best estimate is difficult  
19 to ascertain.

20 With the revised methodology, emissions were partitioned among the atmosphere ( $245 \pm 5$  GtC),  
21 ocean ( $145 \pm 20$  GtC), and the land ( $185 \pm 55$  GtC). The use of nearly independent estimates for  
22 the individual terms shows a cumulative budget imbalance of 20 GtC during 1870-2016, which, if  
23 correct, suggests emissions are too high by the same proportion or the land or ocean sinks are  
24 underestimated. The imbalance originates largely from the large  $E_{LUC}$  during the mid 1920s and  
25 the mid 1960s which is unmatched by a growth in atmospheric  $CO_2$  concentration as recorded in  
26 ice cores (Fig. 3). The known loss of additional sink capacity of about 15 GtC due to reduced forest  
27 cover has not been accounted in our method and further exacerbates the budget imbalance  
28 (Section 2.7.3).



1 Cumulative emissions through to year 2017 increase to  $610 \pm 65$  GtC ( $2235 \pm 240$  GtCO<sub>2</sub>), with  
2 about 70% contribution from  $E_{FF}$  and about 30% contribution from  $E_{LUC}$ . Cumulative emissions and  
3 their partitioning for different periods are provided in Table 9.

4 Given the large revision in cumulative emissions, and its persistent uncertainties, we suggest  
5 extreme caution is needed if using our updated cumulative emission estimate to determine the  
6 “remaining carbon budget” to stay below given temperature limit (Rogelj et al., 2016). We suggest  
7 estimating the remaining carbon budget by integrating scenario data from the current time to  
8 some time in the future as proposed recently (Millar et al., 2017).

#### 9 **4 Discussion**

10 Each year when the global carbon budget is published, each component for all previous years is  
11 updated to take into account corrections that are the result of further scrutiny and verification of  
12 the underlying data in the primary input data sets. The updates have generally been relatively  
13 small (Fig. 9). However this year, we introduced a major methodological change to assess both  
14  $S_{OCEAN}$  and  $S_{LAND}$  directly using multiple process models constrained by observations, and to keep  
15 track of the budget imbalance separately. We also use multiple bookkeeping estimates for  $E_{LUC}$ .  
16 Therefore, the update compared to previous years has led to more substantial revisions,  
17 particularly concerning the mean  $S_{OCEAN}$ , the variability of  $S_{LAND}$ , and the trends in  $E_{LUC}$  (Fig. 9).

18 The budget imbalance provides a measure of the limitations in observations, in understanding or  
19 full representation of processes in models, and/or in the integration of the carbon budget  
20 components. The mean global budget imbalance is close to zero and there is no trend over the  
21 entire time period (Fig. 4). However, the budget imbalance reaches as much as  $\pm 2$  GtC yr<sup>-1</sup> in  
22 individual years, and  $\pm 0.6$  GtC yr<sup>-1</sup> in individual decades (Table 7). Such large budget imbalance  
23 limits our ability to verify reported emissions and limits our confidence in the underlying  
24 processes regulating the carbon cycle feedbacks with climate change (Peters et al., 2017).

25 Another semi-independent way to evaluate the carbon budget results is provided through the use  
26 of atmospheric and oceanic CO<sub>2</sub> data in data-products (atmospheric inversions and pCO<sub>2</sub>-based  
27 ocean flux products). The comparison shows a first-order consistency between pCO<sub>2</sub>-based data-  
28 products and process models but with substantial discrepancies, particularly for the allocation of  
29 the mean surface fluxes between the tropics and the Northern hemisphere, and for highlighting  
30 underestimated decadal variability in  $S_{OCEAN}$ . Understanding the causes of these discrepancies and



1 further analysis of regional carbon budgets would provide additional information to quantify and  
2 improve our estimates, as has been shown by the project REgional Carbon Cycle Assessment and  
3 Processes (RECCAP; Canadell et al., 2012-2013).

4 To help improve the Global Carbon Budget components, we provide a list of the major known  
5 uncertainties for each component, defined as those uncertainties that have been a demonstrated  
6 effect of at least  $0.3 \text{ GtC yr}^{-1}$  (Table 10). We identified multiple sources of uncertainties for  $E_{LUC}$ ,  
7 including in the land-cover and land-use change statistics, representation of management  
8 processes, and methodologies. There are also multiple sources of uncertainties in  $S_{LAND}$ , mostly  
9 related to the understanding and representation of processes, and in  $S_{OCEAN}$ , particularly related to  
10 representing the effects of variable ocean circulation in models as highlighted by recent  
11 observations. Finally, the quality of the energy statistics and of the emissions factors are largest  
12 sources of uncertainties for  $E_{FF}$ . There are no demonstrated uncertainties in  $G_{ATM}$  larger than  $0.3$   
13  $\text{GtC yr}^{-1}$ , although the conversion of the growth rate into a global annual flux assuming  
14 instantaneous mixing throughout the atmosphere introduces additional errors that have not yet  
15 been quantified. Multiple other sources of uncertainties have been identified (i.e. in cement  
16 emissions) that could add up to significant contributions but are unlikely to be the main sources of  
17 the budget imbalance.

18 There are many more uncertainties affecting the annual estimates compared to the mean and  
19 trend, some of which could be improved with better data. Of the various terms in the global  
20 budget, only the emissions from fossil fuels and industry and the growth rate in atmospheric  $\text{CO}_2$   
21 concentration are based primarily on empirical inputs supporting annual estimates in this carbon  
22 budget.  $p\text{CO}_2$ -based flux products for the ocean  $\text{CO}_2$  sink provide new ways to evaluate the model  
23 results, but there are still large discrepancies among estimates. Given the growing reliance on  
24 process models and  $p\text{CO}_2$ -based flux products in our Global Carbon Budget, it is critical that data-  
25 based metrics are developed and used to inform the selection of models and the improvement of  
26 their process representation in the long term.

## 27 **5 Data availability**

28 The data presented here are made available in the belief that their wide dissemination will lead to  
29 greater understanding and new scientific insights of how the carbon cycle works, how humans are  
30 altering it, and how we can mitigate the resulting human-driven climate change. The free



1 availability of these data does not constitute permission for publication of the data. For research  
2 projects, if the data are essential to the work, or if an important result or conclusion depends on  
3 the data, co-authorship may need to be considered. Full contact details and information on how  
4 to cite the data are given at the top of each page in the accompanying database, and summarised  
5 in Table 2.

6 The accompanying database includes two Excel files organised in the following spreadsheets  
7 (accessible with the free viewer <http://www.microsoft.com/en-us/download/details.aspx?id=10>):

8 File `Global_Carbon_Budget_2017v1.0.xlsx` includes the following:

- 9 1. Summary
- 10 2. The global carbon budget (1959-2016);
- 11 3. Global CO<sub>2</sub> emissions from fossil fuels and cement production by fuel type, and the per-capita  
12 emissions (1959-2016);
- 13 4. CO<sub>2</sub> emissions from land-use change from the individual methods and models (1959-2016);
- 14 5. Ocean CO<sub>2</sub> sink from the individual ocean models and pCO<sub>2</sub>-based products (1959-2016);
- 15 6. Terrestrial CO<sub>2</sub> sink from the DGVMs (1959-2016);
- 16 7. Additional information on the carbon balance prior to 1959 (1750-2016).

17 File `National_Carbon_Emissions_2017v1.0.xlsx` includes the following:

- 18 1. Summary
- 19 2. Territorial country CO<sub>2</sub> emissions from fossil fuels and industry (1959-2016) from CDIAC,  
20 extended to 2016 using BP data;
- 21 3. Territorial country CO<sub>2</sub> emissions from fossil fuels and industry (1959-2016) from CDIAC with  
22 UNFCCC data overwritten where available, extended to 2016 using BP data;
- 23 4. Consumption country CO<sub>2</sub> emissions from fossil fuels and industry and emissions transfer  
24 from the international trade of goods and services (1990-2015) using CDIAC/UNFCCC data  
25 (worksheet 3 above) as reference;
- 26 5. Emissions transfers (Consumption minus territorial emissions; 1990-2015);
- 27 6. Country definitions;
- 28 7. Details of disaggregated countries;
- 29 8. Details of aggregated countries.

30 National emissions data are also available from the Global Carbon Atlas ([globalcarbonatlas.org](http://globalcarbonatlas.org)).



## 1 **6 Conclusions**

2 The estimation of global CO<sub>2</sub> emissions and sinks is a major effort by the carbon cycle research  
3 community that requires a combination of measurements and compilation of statistical estimates  
4 and results from models. The delivery of an annual carbon budget serves two purposes. First,  
5 there is a large demand for up-to-date information on the state of the anthropogenic perturbation  
6 of the climate system and its underpinning causes. A broad stakeholder community relies on the  
7 data sets associated with the annual carbon budget including scientists, policy makers, businesses,  
8 journalists, and the broader society increasingly engaged in adapting to and mitigating human-  
9 driven climate change. Second, over the last decade we have seen unprecedented changes in the  
10 human and biophysical environments (e.g. changes in the growth of fossil fuel emissions, ocean  
11 temperatures, and strength of the sink), which call for more frequent assessments of the state of  
12 the planet, and by implication, a better understanding of the future evolution of the carbon cycle.  
13 Both the ocean and the land surface presently remove a large fraction of anthropogenic  
14 emissions. Any significant change in the function of carbon sinks is of great importance to climate  
15 policymaking, as they affect the excess CO<sub>2</sub> remaining in the atmosphere and therefore the  
16 compatible emissions for any climate stabilisation target. Better constraints of carbon cycle  
17 models against contemporary data sets raise the capacity for the models to become more  
18 accurate at future projections. This all requires more frequent, robust, and transparent data sets  
19 and methods that can be scrutinized and replicated. This paper via 'living data' will help to keep  
20 track of new budget updates.

21 **Acknowledgments.** We thank all people and institutions who provided the data used in this  
22 carbon budget; C. Enright, W. Peters, and S. Shu for their involvement in the development, use  
23 and analysis of the models and data-products used here; F. Joos, S. Khatiwala and T. DeVries for  
24 providing historical data; and A. Kirk for technical support. We thank E. Dlugokencky who provided  
25 the atmospheric CO<sub>2</sub> measurements used here; C. Landa, C. Bernard and S. Jones of the Bjerknes  
26 Climate Data Centre and the ICOS Ocean Thematic Centre data management at the University of  
27 Bergen, who helped with gathering information from the SOCAT community, and all those  
28 involved in collecting and providing oceanographic CO<sub>2</sub> measurements used here, in particular for  
29 the new ocean data for year 2016 (see Table A2). This is NOAA-PMEL contribution number 4728.  
30 We thank the institutions and funding agencies responsible for the collection and quality control  
31 of the data included in SOCAT, and the support of the International Ocean Carbon Coordination



1 Project (IOCCP), the Surface Ocean Lower Atmosphere Study (SOLAS), and the Integrated Marine  
2 Biogeochemistry, Ecosystem Research (IMBER) programme. Long-term support for the CRU TS  
3 dataset is currently provided by the UK National Centre for Atmospheric Science (NCAS), a NERC  
4 collaborative centre.

5 Finally, we thank all funders who have supported the individual and joint contributions to this  
6 work (see Appendix Table A1), as well as M. Heimann, H. Dolman, and the many researchers who  
7 have provided feedback during the GCP community consultation held at the 10<sup>th</sup> International CO<sub>2</sub>  
8 Conference in Interlaken, Switzerland.

9



## 1 References

- 2 Andres, R., Boden, T., and Higdon, D.: A new evaluation of the uncertainty associated with CDIA  
3 estimates of fossil fuel carbon dioxide emission, *Tellus B*, 66, 23616, 2014.
- 4 Andres, R. J., Boden, T. A., Bréon, F.-M., Ciais, P., Davis, S., Erickson, D., Gregg, J. S., Jacobson, A., Marland,  
5 G., Miller, J., Oda, T., Olivier, J. G. J., Raupach, M. R., Rayner, P., and Treanton, K.: A synthesis of  
6 carbon dioxide emissions from fossil-fuel combustion, *Biogeosciences*, 9, 1845-1871, 2012.
- 7 Andrew, R. M.: Global CO<sub>2</sub> emissions from cement production, *Earth Syst. Sci. Data Discuss.*, 2017, 1-52,  
8 2017.
- 9 Andrew, R. M. and Peters, G. P.: A multi-region input-output table based on the Global Trade Analysis  
10 Project Database (GTAP-MRIO), *Economic Systems Research*, 25, 99-121, 2013.
- 11 Archer, D., Eby, M., Brovkin, V., Ridgwell, A., Cao, L., Mikolajewicz, U., Caldeira, K. M., K., Munhoven, G.,  
12 Montenegro, A., and Tokos, K.: Atmospheric Lifetime of Fossil Fuel Carbon Dioxide, *Annual Review of*  
13 *Earth and Planetary Sciences*, 37, 117-134, 2009.
- 14 Arneeth, A., Sitch, S., Pongratz, J., Stocker, B. D., Ciais, P., Poulter, B., Bayer, A. D., Bondeau, A., Calle, L.,  
15 Chini, L. P., Gasser, T., Fader, M., Friedlingstein, P., Kato, E., Li, W., Lindeskog, M., Nabel, J. E. M. S.,  
16 Pugh, T. A. M., Robertson, E., Viovy, N., Yue, C., and Zaehle, S.: Historical carbon dioxide emissions  
17 caused by land-use changes are possibly larger than assumed, *Nature Geosci*, 10, 79-84, 2017.
- 18 Arora, V. K., Boer, G. J., Christian, J. R., Curry, C. L., Denman, K. L., Zahariev, K., Flato, G. M., Scinocca, J. F.,  
19 Merryfield, W. J., and Lee, W. G.: The Effect of Terrestrial Photosynthesis Down Regulation on the  
20 Twentieth-Century Carbon Budget Simulated with the CCCma Earth System Model, *Journal of Climate*,  
21 22, 6066-6088, 2009.
- 22 Aumont, O. and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, *Global*  
23 *Biogeochemical Cycles*, 20, GB2017, 2006.
- 24 Bakker, D. C. E., Pfeil, B., Landa, C. S., Metzl, N., O'Brien, K. M., Olsen, A., Smith, K., Cosca, C., Harasawa, S.,  
25 Jones, S. D., Nakaoka, S. I., Nojiri, Y., Schuster, U., Steinhoff, T., Sweeney, C., Takahashi, T., Tilbrook, B.,  
26 Wada, C., Wanninkhof, R., Alin, S. R., Balestrini, C. F., Barbero, L., Bates, N. R., Bianchi, A. A., Bonou, F.,  
27 Boutin, J., Bozec, Y., Burger, E. F., Cai, W. J., Castle, R. D., Chen, L., Chierici, M., Currie, K., Evans, W.,  
28 Featherstone, C., Feely, R. A., Fransson, A., Goyet, C., Greenwood, N., Gregor, L., Hankin, S., Hardman-  
29 Mountford, N. J., Harlay, J., Hauck, J., Hoppema, M., Humphreys, M. P., Hunt, C. W., Huss, B., Ibáñez,  
30 J. S. P., Johannessen, T., Keeling, R., Kitidis, V., Körtzinger, A., Kozyr, A., Krasakopoulou, E., Kuwata, A.,  
31 Landschützer, P., Lauvset, S. K., Lefèvre, N., Lo Monaco, C., Manke, A., Mathis, J. T., Merlivat, L.,  
32 Millero, F. J., Monteiro, P. M. S., Munro, D. R., Murata, A., Newberger, T., Omar, A. M., Ono, T.,  
33 Paterson, K., Pearce, D., Pierrot, D., Robbins, L. L., Saito, S., Salisbury, J., Schlitzer, R., Schneider, B.,  
34 Schweitzer, R., Sieger, R., Skjelvan, I., Sullivan, K. F., Sutherland, S. C., Sutton, A. J., Tadokoro, K.,  
35 Telszewski, M., Tuma, M., van Heuven, S. M. A. C., Vandemark, D., Ward, B., Watson, A. J., and Xu, S.:  
36 A multi-decade record of high-quality fCO<sub>2</sub> data in version 3 of the Surface Ocean CO<sub>2</sub> Atlas (SOCAT),  
37 *Earth Syst. Sci. Data*, 8, 383-413, 2016.
- 38 Ballantyne, A. P., Alden, C. B., Miller, J. B., Tans, P. P., and White, J. W. C.: Increase in observed net carbon  
39 dioxide uptake by land and oceans during the last 50 years, *Nature*, 488, 70-72, 2012.
- 40 Ballantyne, A. P., Andres, R., Houghton, R., Stocker, B. D., Wanninkhof, R., Anderegg, W., Cooper, L. A.,  
41 DeGrandpre, M., Tans, P. P., Miller, J. B., Alden, C., and White, J. W. C.: Audit of the global carbon  
42 budget: estimate errors and their impact on uptake uncertainty, *Biogeosciences*, 12, 2565-2584, 2015.
- 43 Bauer, J. E., Cai, W.-J., Raymond, P. A., Bianchi, T. S., Hopkinson, C. S., and Regnier, P. A. G.: The changing  
44 carbon cycle of the coastal ocean, *Nature*, 504, 61-70, 2013.
- 45 Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. L. H., Ménard, C. B., Edwards, J. M., Hendry, M.  
46 A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S.  
47 B., and Harding, R. J.: The Joint UK Land Environment Simulator (JULES), model description - Part 1:  
48 Energy and water fluxes, *Geoscientific Model Development*, doi: 10.5194/gmd-4-677-2011, 2011. 677-  
49 699, 2011.
- 50 Betts, R. A., Jones, C. D., Knight, J. R., Keeling, R. F., and Kennedy, J. J.: El Niño and a record CO<sub>2</sub> rise,  
51 *Nature Clim. Change*, 6, 806-810, 2016.



- 1 Boden, T. A., Marland, G., and Andres, R. J.: Global, Regional, and National Fossil-Fuel CO<sub>2</sub> Emissions,  
2 available at: [http://cdiac.ornl.gov/trends/emis/overview\\_2014.html](http://cdiac.ornl.gov/trends/emis/overview_2014.html) (last access: July 2017), Oak Ridge  
3 National Laboratory, U.S. Department of Energy, Oak Ridge, Tenn., U.S.A. , 2017.
- 4 BP: BP Statistical Review of World Energy June 2017  
5 [https://www.bp.com/content/dam/bp/en/corporate/pdf/energy-economics/statistical-review-](https://www.bp.com/content/dam/bp/en/corporate/pdf/energy-economics/statistical-review-2017/bp-statistical-review-of-world-energy-2017-full-report.pdf)  
6 [2017/bp-statistical-review-of-world-energy-2017-full-report.pdf](https://www.bp.com/content/dam/bp/en/corporate/pdf/energy-economics/statistical-review-2017/bp-statistical-review-of-world-energy-2017-full-report.pdf) (last access: June 2017), 2017.
- 7 Bruno, M. and Joos, F.: Terrestrial carbon storage during the past 200 years: A monte carlo analysis of CO<sub>2</sub>  
8 data from ice core and atmospheric measurements, *Global Biogeochemical Cycles*, 11, 111-124, 1997.
- 9 Buitenhuis, E. T., Rivkin, R. B., Sailley, S., and Le Quéré, C.: Biogeochemical fluxes through  
10 microzooplankton, *Global Biogeochemical Cycles*, 24, 2010.
- 11 Canadell, J. G., Ciais, P., Sabine, C., and Joos, F.: REgional Carbon Cycle Assessment and Processes  
12 (RECCAP), available at: [http://www.biogeosciences.net/special\\_issue107.html](http://www.biogeosciences.net/special_issue107.html), *Biogeosciences Special*  
13 *Issue*, 2012-2013.
- 14 Canadell, J. G., Le Quéré, C., Raupach, M. R., Field, C. B., Buitenhuis, E. T., Ciais, P., Conway, T. J., Gillett, N.  
15 P., Houghton, R. A., and Marland, G.: Contributions to accelerating atmospheric CO<sub>2</sub> growth from  
16 economic activity, carbon intensity, and efficiency of natural sinks, *Proceedings of the National*  
17 *Academy of Sciences of the United States of America*, 104, 18866-18870, 2007.
- 18 Carbontracker Team: Compilation of near real time atmospheric carbon dioxide data provided by NOAA  
19 and EC; obspack\_co2\_1\_NRT\_v3.3\_2017-04-19; NOAA Earth System Research Laboratory, Global  
20 Monitoring Division. <http://doi.org/10.15138/G3G01J>, 2017.
- 21 CCIA: China Coal Industry Association (CCIA), 2017. Analysis of current economic trend of coal in China (当  
22 前全国煤炭经济运行走势分析).  
23 <http://www.coalchina.org.cn/detail/17/07/24/00000025/content.html>. (in Chinese; last access: 15  
24 September 2017), 2017.
- 25 CEA: Central Electricity Authority (CEA), 2017: Daily Coal – Archive. Central Electricity Authority.  
26 <http://www.cea.nic.in/dailyarchive.html> (Last Accessed: September 2017), 2017.
- 27 Chevallier, F.: On the statistical optimality of CO<sub>2</sub> atmospheric inversions assimilating CO<sub>2</sub> column  
28 retrievals, *Atmospheric Chemistry and Physics*, 15, 11133-11145, 2015.
- 29 Chevallier, F., M. Fisher, P. Peylin, S. Serrar, P. Bousquet, F.-M. Bréon, A. Chédin, and Ciais, P.: Inferring  
30 CO<sub>2</sub> sources and sinks from satellite observations: Method and application to TOVS data, *J. Geophys.*  
31 *Res.*, D24309, 2005.
- 32 Ciais, P., Sabine, C., Govindasamy, B., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway,  
33 J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R., Piao, S., and Thornton, P.: Chapter 6: Carbon and  
34 Other Biogeochemical Cycles. In: *Climate Change 2013 The Physical Science Basis*, Stocker, T., Qin, D.,  
35 and Plattner, G.-K. (Eds.), Cambridge University Press, Cambridge, 2013.
- 36 CIL: Coal India Limited, 2017: Production and Offtake Performance of CIL and Subsidiary Companies.  
37 <https://www.coalindia.in/en-us/performance/physical.aspx> . (Last Accessed: September 2017), 2017.
- 38 Clarke, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery,  
39 R. L. H., Blyth, E., Boucher, O., Cox, P. M., and Harding, R. J.: The Joint UK Land Environment Simulator  
40 (JULES), model description - Part 2: Carbon fluxes and vegetation dynamics. , *Geoscientific Model*  
41 *Development*, 4, 701-772, 2011.
- 42 Cox, P. M., Pearson, D., Booth, B. B., Friedlingstein, P., Huntingford, C., Jones, C. D., and Luke, C. M.:  
43 Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability, *Nature*, 494,  
44 341-344, 2013.
- 45 Davis, S. J. and Caldeira, K.: Consumption-based accounting of CO<sub>2</sub> emissions, *Proceedings of the National*  
46 *Academy of Sciences*, 107, 5687-5692, 2010.
- 47 Denman, K. L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze,  
48 C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., Leite da Silva Dias, P., Wofsy, S. C., and  
49 Zhang, X.: Couplings Between Changes in the Climate System and Biogeochemistry. In: *Climate Change*  
50 *2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of*  
51 *the Intergovernmental Panel on Climate Change*, Solomon, S., Qin D., Manning M., Marquis M., Averyt



- 1 K., Tignor M. M. B., Miller, H. L., and L., C. Z. (Eds.), Cambridge University Press, Cambridge, UK and  
2 New York, USA, 2007.
- 3 DeVries, T.: The oceanic anthropogenic CO<sub>2</sub> sink: Storage, air-sea fluxes, and transports over the industrial  
4 era, *Global Biogeochemical Cycles*, 28, 631-647, 2014.
- 5 DeVries, T., Holzer, M., and Primeau, F.: Recent increase in oceanic carbon uptake driven by weaker upper-  
6 ocean overturning, *Nature*, 542, 215-218, 2017.
- 7 Dlugokencky, E. and Tans, P.: Trends in atmospheric carbon dioxide, National Oceanic & Atmospheric  
8 Administration, Earth System Research Laboratory (NOAA/ESRL), available at  
9 <http://www.esrl.noaa.gov/gmd/ccgg/trends> (last access: 28 October 2016), 2016.
- 10 Dlugokencky, E. and Tans, P.: Trends in atmospheric carbon dioxide, National Oceanic & Atmospheric  
11 Administration, Earth System Research Laboratory (NOAA/ESRL), available at  
12 <http://www.esrl.noaa.gov/gmd/ccgg/trends/global.html> (last access: 7 September 2017), 2017.
- 13 Doney, S. C., Lima, I., Feely, R. A., Glover, D. M., Lindsay, K., Mahowald, N., Moore, J. K., and Wanninkhof,  
14 R.: Mechanisms governing interannual variability in upper-ocean inorganic carbon system and air-sea  
15 CO<sub>2</sub> fluxes: Physical climate and atmospheric dust, *Deep-Sea Res Pt II*, 56, 640-655, 2009.
- 16 Duce, R. A., LaRoche, J., Altieri, K., Arrigo, K. R., Baker, A. R., Capone, D. G., Cornell, S., Dentener, F.,  
17 Galloway, J., Ganeshram, R. S., Geider, R. J., Jickells, T., Kuypers, M. M., Langlois, R., Liss, P. S., Liu, S.  
18 M., Middelburg, J. J., Moore, C. M., Nickovic, S., Oschlies, A., Pedersen, T., Prospero, J., Schlitzer, R.,  
19 Seitzinger, S., Sorensen, L. L., Uematsu, M., Ulloa, O., Voss, M., Ward, B., and Zamora, L.: Impacts of  
20 atmospheric anthropogenic nitrogen on the open ocean, *Science*, 320, 893-897, 2008.
- 21 Dufour, C. O., Le Sommer, J., Gehlen, M., Orr, J. C., Molines, J. M., Simeon, J., and Barnier, B.: Eddy  
22 compensation and controls of the enhanced sea-to-air CO<sub>2</sub> flux during positive phases of the  
23 Southern Annular Mode, *Global Biogeochemical Cycles*, 27, 950-961, 2013.
- 24 Durant, A. J., Le Quéré, C., Hope, C., and Friend, A. D.: Economic value of improved quantification in global  
25 sources and sinks of carbon dioxide, *Phil. Trans. A*, 269, 1967-1979, 2011.
- 26 EIA: U.S. Energy Information Administration, Short-Term Energy and Winter Fuels Outlook,,  
27 <http://www.eia.gov/forecasts/steo/outlook.cfm>, (last access: September 2017), 2017.
- 28 Erb, K.-H., Kastner, T., Luyssaert, S., Houghton, R. A., Kuemmerle, T., Olofsson, P., and Haberl, H.: Bias in  
29 the attribution of forest carbon sinks, *Nature Climate Change*, 3, 854-856, 2013.
- 30 Etheridge, D. M., Steele, L. P., Langenfelds, R. L., and Francey, R. J.: Natural and anthropogenic changes in  
31 atmospheric CO<sub>2</sub> over the last 1000 years from air in Antarctic ice and firn, *Journal of Geophysical*  
32 *Research*, 101, 4115-4128, 1996.
- 33 FAO: Global Forest Resources Assessment 2015, Food and Agriculture Organization of the United Nations,  
34 Rome, Italy, 2015.
- 35 FAOSTAT: Food and Agriculture Organization Statistics Division, available at: <http://faostat.fao.org/> (last  
36 access: 2015), 2015.
- 37 Francey, R. J., Trudinger, C. M., van der Schoot, M., Law, R. M., Krummel, P. B., Langenfelds, R. L., Steele, L.  
38 P., Allison, C. E., Stavert, A. R., Andres, R. J., and Rodenbeck, C.: Reply to 'Anthropogenic CO<sub>2</sub>  
39 emissions', *Nature Clim. Change*, 3, 604-604, 2013.
- 40 Friedlingstein, P., Andrew, R. M., Rogelj, J., Peters, G. P., Canadell, J. G., Knutti, R., Luderer, G., Raupach, M.  
41 R., Schaeffer, M., van Vuuren, D. P., and Le Quéré, C.: Persistent growth of CO<sub>2</sub> emissions and  
42 implications for reaching climate targets, *Nature Geoscience*, 2014. 709-715, 2014.
- 43 Friedlingstein, P., Houghton, R. A., Marland, G., Hackler, J., Boden, T. A., Conway, T. J., Canadell, J. G.,  
44 Raupach, M. R., Ciais, P., and Le Quéré, C.: Update on CO<sub>2</sub> emissions, *Nature Geoscience*, 3, 811-812,  
45 2010.
- 46 Gasser, T., Ciais, P., Boucher, O., Quilcaille, Y., Tortora, M., Bopp, L., and Hauglustaine, D.: The compact  
47 Earth system model OSCAR v2.2: description and first results, *Geosci. Model Dev.*, 10, 271-319, 2017.
- 48 GCP: The Global Carbon Budget 2007, available at:  
49 <http://www.globalcarbonproject.org/carbonbudget/archive.htm>, last access: 7 November 2016.,  
50 2007.



- 1 Giglio, L., Descloitres, J., Justice, C. O., and Kaufman, Y. J.: An enhanced contextual fire detection algorithm  
2 for MODIS, *Remote Sensing of Environment*, 87, 273-282, 2003.
- 3 Gitz, V. and Ciais, P.: Amplifying effects of land-use change on future atmospheric CO<sub>2</sub> levels, *Global*  
4 *Biogeochemical Cycles*, 17, 1024, 2003.
- 5 GLOBALVIEW: Cooperative Global Atmospheric Data Integration Project; (2016): Multi-laboratory  
6 compilation of atmospheric carbon dioxide data for the period 1957-2015;  
7 obspack\_co2\_1\_GLOBALVIEWplus\_v2.1\_2016\_09\_02; NOAA Earth System Research Laboratory,  
8 Global Monitoring Division. <http://dx.doi.org/10.15138/G3059Z>, 2016.
- 9 Goll, D. S., V. Brovkin, J. Liski, T. Raddatz, T. Thum, and Todd-Brown, K. E. O.: Strong dependence of CO<sub>2</sub>  
10 emissions from anthropogenic land cover change on initial land cover and soil carbon parametrization,  
11 *Global Biogeochem. Cycles*, 29, 2015.
- 12 Gregg, J. S., Andres, R. J., and Marland, G.: China: Emissions pattern of the world leader in CO<sub>2</sub> emissions  
13 from fossil fuel consumption and cement production, *Geophysical Research Letters*, 35, L08806, 2008.
- 14 Guimberteau, M., Zhu, D., Maignan, F., Huang, Y., Yue, C., Dantec-Nédélec, S., Ottlé, C., Jornet-Puig, A.,  
15 Bastos, A., Laurent, P., Goll, D., Bowring, S., Chang, J., Guenet, B., Tifafi, M., Peng, S., Krinner, G.,  
16 Ducharne, A., Wang, F., Wang, T., Wang, X., Wang, Y., Yin, Z., Lauerwald, R., Joetzjer, E., Qiu, C., Kim,  
17 H., and Ciais, P.: ORCHIDEE-MICT (revision 4126), a land surface model for the high-latitudes: model  
18 description and validation, *Geosci. Model Dev. Discuss.*, 2017, 1-65, 2017.
- 19 Hansis, E., Davis, S. J., and Pongratz, J.: Relevance of methodological choices for accounting of land use  
20 change carbon fluxes, *Global Biogeochemical Cycles*, 29, 1230-1246, 2015.
- 21 Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H.: Updated high-resolution grids of monthly climatic  
22 observations – the CRU TS3.10 Dataset, *International Journal of Climatology*, 34, 623-642, 2014.
- 23 Hauck, J., Kohler, P., Wolf-Gladrow, D., and Volker, C.: Iron fertilisation and century-scale effects of open  
24 ocean dissolution of olivine in a simulated CO<sub>2</sub> removal experiment, *Environmental Research Letters*,  
25 11, 024007, 2016.
- 26 Haverd, V., Smith, B., Nieradzik, L., Briggs, P. R., Woodgate, W., Trudinger, C. M., and Canadell, J. G.: A new  
27 version of the CABLE land surface model, incorporating land use and land cover change, woody  
28 vegetation demography and a novel optimisation-based approach to plant coordination of electron  
29 transport and carboxylation capacity-limited photosynthesis., *Geoscientific Model Development*, doi:  
30 (submitted), 2017. 2017.
- 31 Hertwich, E. G. and Peters, G. P.: Carbon Footprint of Nations: A Global, Trade-Linked Analysis,  
32 *Environmental Science and Technology*, doi: 10.1021/es803496a, 2009. 6414-6420, 2009.
- 33 Hooijer, A., Page, S., Canadell, J. G., Silvius, M., Kwadijk, J., Wösten, H., and Jauhiainen, J.: Current and  
34 future CO<sub>2</sub> emissions from drained peatlands in Southeast Asia, *Biogeosciences*, 7, 1505-  
35 1514, 2010.
- 36 Houghton, R. A.: Revised estimates of the annual net flux of carbon to the atmosphere from changes in  
37 land use and land management 1850-2000, *Tellus Series B-Chemical and Physical Meteorology*, 55,  
38 378-390, 2003.
- 39 Houghton, R. A., House, J. I., Pongratz, J., van der Werf, G. R., DeFries, R. S., Hansen, M. C., Le Quééré, C.,  
40 and Ramankutty, N.: Carbon emissions from land use and land-cover change, *Biogeosciences*, 9, 5125-  
41 5142, 2012.
- 42 Houghton, R. A. and Nassikas, A. A.: Global and regional fluxes of carbon from land use and land cover  
43 change 1850-2015, *Global Biogeochemical Cycles*, 31, 456-472, 2017.
- 44 Hourdin, F., Musat, I., Bony, S., Braconnot, P., Codron, F., Dufresne, J.-L., Fairhead, L., Filiberti, M.-A.,  
45 Freidlingstein, P., Grandpeix, J.-Y., Krinner, G., LeVan, P., Li, Z.-X., and Lott, F.: The LMDZ4 general  
46 circulation model: climate performance and sensitivity to parametrized physics with emphasis on  
47 tropical convection, *Climate Dynamics*, 27, 787-813, 2006.
- 48 Hurrte, G., Chini, L., Sahajpal, R., and Froelking, S.: Harmonization of global land-use change and  
49 management for the period 850-2100, *Geoscientific Model Development*, in prep., in prep.
- 50 Hurrte, G. C., Chini, L. P., Froelking, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P., Hibbard, K., Houghton,  
51 R. A., Janetos, A., Jones, C. D., Kindermann, G., Kinoshita, T., Klein Goldewijk, K., Riahi, K., Shevliakova,



- 1 E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D. P., and Wang, Y. P.:  
2 Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual  
3 land-use transitions, wood harvest, and resulting secondary lands, *Climatic Change*, 109, 117-161,  
4 2011.
- 5 IEA/OECD: CO2 emissions from fuel combustion, International Energy Agency/Organisation for Economic  
6 Cooperation and Development, Paris, 2016.
- 7 Ilyina, T., Six, K., Segschneider, J., Maier-Reimer, E., Li, H., and Núñez-Riboni, I.: The global ocean  
8 biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-  
9 Earth System Model in different CMIP5 experimental realizations, *Journal of Advances in Modeling  
10 Earth Systems*, 5, 287-315, 2013.
- 11 IMF: World Economic Outlook of the International Monetary Fund, available at:  
12 <http://www.imf.org/external/ns/cs.aspx?id=29> (last access September 2017), 2017.
- 13 IPCC: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National  
14 Greenhouse Gas Inventories Programme, Institute for Global Environmental Strategies (IGES), Japan,  
15 2006.
- 16 Ito, A. and Inatomi, M.: Use of a process-based model for assessing the methane budgets of global  
17 terrestrial ecosystems and evaluation of uncertainty, *Biogeosciences*, 9, 759-773, 2012.
- 18 Jackson, R. B., Canadell, J. G., Le Quéré, C., Andrew, R. M., Korsbakken, J. I., Peters, G. P., and Nakicenovic,  
19 N.: Reaching peak emissions, *Nature Climate Change*, 6, 7-10, 2016.
- 20 Jackson, R. B., Le Quéré, C., Andrew, R. M., Canadell, J. G., Peters, G. P., Roy, J., and Wu, L.: Warning signs  
21 for stabilizing global CO2 emissions, *Environmental Research Letters*, doi: 10.1088/1748-9326/aa9662,  
22 2017. 2017.
- 23 Jacobson, A. R., Mikaloff Fletcher, S. E., Gruber, N., Sarmiento, J. L., and Gloor, M.: A joint atmosphere-  
24 ocean inversion for surface fluxes of carbon dioxide: 1. Methods and global-scale fluxes, *Global  
25 Biogeochemical Cycles*, 21, GB1019, 2007.
- 26 Jain, A. K., Meiyappan, P., Song, Y., and House, J. I.: CO2 Emissions from Land-Use Change Affected More  
27 by Nitrogen Cycle, than by the Choice of Land Cover Data, *Global Change Biology*, 9, 2893-2906, 2013.
- 28 Joos, F. and Spahni, R.: Rates of change in natural and anthropogenic radiative forcing over the past 20,000  
29 years, *Proceedings of the National Academy of Science*, 105, 1425-1430, 2008.
- 30 Kato, E., Kinoshita, T., Ito, A., Kawamiya, M., and Yamagata, Y.: Evaluation of spatially explicit emission  
31 scenario of land-use change and biomass burning using a process-based biogeochemical model,  
32 *Journal of Land Use Science*, 8, 104-122, 2013.
- 33 Keeling, C. D., Bacastow, R. B., Bainbridge, A. E., Ekdhal, C. A., Guenther, P. R., and Waterman, L. S.:  
34 Atmospheric carbon dioxide variations at Mauna Loa Observatory, Hawaii, *Tellus*, 28, 538-551, 1976.
- 35 Keeling, R. F. and Manning, A. C.: 5.15 - Studies of Recent Changes in Atmospheric O2 Content. In: *Treatise  
36 on Geochemistry: Second Edition*, Holland, H. D. and Turekian, K. K. (Eds.), Elsevier, Oxford, 2014.
- 37 Keller, K. M., Lienert, S., Bozbiyik, A., Stocker, T. F., Churakova, O. V., Frank, D. C., Klesse, S., Koven, C. D.,  
38 Leuenberger, M., Riley, W. J., Saurer, M., Siegwolf, R., Weigt, R. B., and Joos, F.: 20th century changes  
39 in carbon isotopes and water-use efficiency: tree-ring-based evaluation of the CLM4.5 and LPX-Bern  
40 models, *Biogeosciences*, 14, 2641-2673, 2017.
- 41 Khatiwala, S., Primeau, F., and Hall, T.: Reconstruction of the history of anthropogenic CO2 concentrations  
42 in the ocean, *Nature*, 462, 346-350, 2009.
- 43 Khatiwala, S., Tanhua, T., Mikaloff Fletcher, S. E., Gerber, M., Doney, S. C., Graven, H. D., Gruber, N.,  
44 McKinley, G. A., Murata, A., Rios, A. F., and Sabine, C. L.: Global ocean storage of anthropogenic  
45 carbon, *Biogeosciences*, 10, 2169-2191, 2013.
- 46 Kirschke, S., Bousquet, P., Ciais, P., Saunoy, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P.,  
47 Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser,  
48 A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Fraser, P. J., Krummel, P. B., Lamarque, J.,  
49 Langenfelds, R. L., Le Quéré, C., Naik, V., O'Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B.,  
50 Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R.,  
51 Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., van der Werf, G. R., Voulgarakis, A., van Weele, M.,



- 1 Weiss, R. F., Williams, J. E., and Zeng, G.: Three decades of global methane sources and sinks, *Nature*  
2 *Geoscience*, 6, 813-823, 2013.
- 3 Klein Goldewijk, K., Beusen, A., Doelman, J., and Stehfest, E.: Anthropogenic land use estimates for the  
4 Holocene; HYDE 3.2, *Earth Syst. Sci. Data*, doi: 10.5194/essd-2016-58, in press., in press.
- 5 Klein Goldewijk, K., Dekker, S. C., and van Zanden, J. L.: Per-capita estimations of long-term historical land  
6 use and the consequences for global change research, *Journal of Land Use Science*, 12, 313-337, 2017.
- 7 Korsbakken, J. I., Peters, G. P., and Andrew, R. M.: Uncertainties around reductions in China's coal use and  
8 CO<sub>2</sub> emissions, *Nature Climate Change*, 6, 687-690, 2016.
- 9 Krinner, G., Viovy, N., de Noblet, N., Ogée, J., Friedlingstein, P., Ciais, P., Sitch, S., Polcher, J., and Prentice,  
10 I. C.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system,  
11 *Global Biogeochemical Cycles*, 19, 1-33, 2005.
- 12 Landschützer, P., Gruber, N., and Bakker, D. C. E.: Decadal variations and trends of the global ocean carbon  
13 sink, *Global Biogeochemical Cycles*, 30, 1396-1417, 2016.
- 14 Landschützer, P., Gruber, N., Bakker, D. C. E., and Schuster, U.: Recent variability of the global ocean  
15 carbon sink, *Global Biogeochemical Cycles*, doi: 10.1002/2014GB004853, 2014. 2014.
- 16 Landschützer, P., Gruber, N., Haumann, A., Rödenbeck, C., Bakker, D. C. E., van Heuven, S., Hoppema, M.,  
17 Metzl, N., Sweeney, C., Takahashi, T., Tilbrook, B., and Wanninkhof, R.: The reinvigoration of the  
18 Southern Ocean carbon sink, *Science*, 349, 1221-1224, 2015.
- 19 Law, R. M., Ziehn, T., Matear, R. J., Lenton, A., Chamberlain, M. A., Stevens, L. E., Wang, Y. P., Sribnovsky,  
20 J., Bi, D., Yan, H., and Vohralik, P. F.: The carbon cycle in the Australian Community Climate and Earth  
21 System Simulator (ACCESS-ESM1) – Part 1: Model description and pre-industrial simulation, *Geosci.*  
22 *Model Dev.*, 10, 2567-2590, 2017.
- 23 Le Quéré, C.: Closing the global budget for CO<sub>2</sub> Global Change, 74, 28-31, 2009.
- 24 Le Quéré, C., Andres, R. J., Boden, T., Conway, T., Houghton, R. A., House, J. I., Marland, G., Peters, G. P.,  
25 van der Werf, G. R., Ahlström, A., Andrew, R. M., Bopp, L., Canadell, J. G., Ciais, P., Doney, S. C.,  
26 Enright, C., Friedlingstein, P., Huntingford, C., Jain, A. K., Jourdain, C., Kato, E., Keeling, R. F., Klein  
27 Goldewijk, K., Levis, S., Levy, P., Lomas, M., Poulter, B., Raupach, M. R., Schwinger, J., Sitch, S., Stocker,  
28 B. D., Viovy, N., Zaehle, S., and Zeng, N.: The global carbon budget 1959–2011, *Earth System Science*  
29 *Data*, 5, 165-185, 2013.
- 30 Le Quéré, C., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Peters, G. P., Manning, A. C., Boden,  
31 T. A., Tans, P. P., Houghton, R. A., Keeling, R. F., Alin, S., Andrews, O. D., Anthoni, P., Barbero, L., Bopp,  
32 L., Chevallier, F., Chini, L. P., Ciais, P., Currie, K., Delire, C., Doney, S. C., Friedlingstein, P., Gkritzalis, T.,  
33 Harris, I., Hauck, J., Haverd, V., Hoppema, M., Klein Goldewijk, K., Jain, A. K., Kato, E., Körtzinger, A.,  
34 Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D., Melton, J. R., Metzl, N., Millero,  
35 F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S. I., O'Brien, K., Olsen, A., Omar, A.  
36 M., Ono, T., Pierrot, D., Poulter, B., Rödenbeck, C., Salisbury, J., Schuster, U., Schwinger, J., Séférian,  
37 R., Skjelvan, I., Stocker, B. D., Sutton, A. J., Takahashi, T., Tian, H., Tilbrook, B., van der Laan-Luijckx, I. T.,  
38 van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget  
39 2016, *Earth Syst. Sci. Data*, 8, 605-649, 2016.
- 40 Le Quéré, C., Moriarty, R., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Friedlingstein, P.,  
41 Peters, G. P., Andres, R. J., Boden, T. A., Houghton, R. A., House, J. I., Keeling, R. F., Tans, P., Arneth, A.,  
42 Bakker, D. C. E., Barbero, L., Bopp, L., Chang, J., Chevallier, F., Chini, L. P., Ciais, P., Fader, M., Feely, R.  
43 A., Gkritzalis, T., Harris, I., Hauck, J., Ilyina, T., Jain, A. K., Kato, E., Kitidis, V., Klein Goldewijk, K., Koven,  
44 C., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lima, I. D., Metzl, N., Millero, F., Munro, D.  
45 R., Murata, A., Nabel, J. E. M. S., Nakaoka, S., Nojiri, Y., O'Brien, K., Olsen, A., Ono, T., Pérez, F. F., Pfeil,  
46 B., Pierrot, D., Poulter, B., Rehder, G., Rödenbeck, C., Saito, S., Schuster, U., Schwinger, J., Séférian, R.,  
47 Steinhoff, T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van der Laan-Luijckx, I. T., van der  
48 Werf, G. R., van Heuven, S., Vandemark, D., Viovy, N., Wiltshire, A., Zaehle, S., and Zeng, N.: Global  
49 Carbon Budget 2015, *Earth Syst. Sci. Data*, 7, 349-396, 2015a.
- 50 Le Quéré, C., Moriarty, R., Andrew, R. M., Peters, G. P., Ciais, P., Friedlingstein, P., Jones, S. D., Sitch, S.,  
51 Tans, P., Arneth, A., Boden, T. A., Bopp, L., Bozec, Y., Canadell, J. G., Chini, L. P., Chevallier, F., Cosca, C.



- 1 E., Harris, I., Hoppema, M., Houghton, R. A., House, J. I., Jain, A. K., Johannessen, T., Kato, E., Keeling,  
2 R. F., Kitidis, V., Goldewijk, K. K., Koven, C., Landa, C. S., Landschutzer, P., Lenton, A., Lima, I. D.,  
3 Marland, G., Mathis, J. T., Metz, N., Nojiri, Y., Olsen, A., Ono, T., Peng, S., Peters, W., Pfeil, B., Poulter,  
4 B., Raupach, M. R., Regnier, P., Rodenbeck, C., Saito, S., Salisbury, J. E., Schuster, U., Schwinger, J.,  
5 Seferian, R., Segschneider, J., Steinhoff, T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van  
6 der Werf, G. R., Viovy, N., Wang, Y. P., Wanninkhof, R., Wiltshire, A., and Zeng, N.: Global carbon  
7 budget 2014, *Earth System Science Data*, 7, 47-85, 2015b.
- 8 Le Quéré, C., Peters, G. P., Andres, R. J., Andrew, R. M., Boden, T. A., Ciais, P., Friedlingstein, P., Houghton,  
9 R. A., Marland, G., Moriarty, R., Sitch, S., Tans, P., Arneeth, A., Arvanitis, A., Bakker, D. C. E., Bopp, L.,  
10 Canadell, J. G., Chini, L. P., Doney, S. C., Harper, A., Harris, I., House, J. I., Jain, A. K., Jones, S. D., Kato,  
11 E., Keeling, R. F., Klein Goldewijk, K., Körtzinger, A., Koven, C., Lefèvre, N., Maignan, F., Omar, A., Ono,  
12 T., Park, G. H., Pfeil, B., Poulter, B., Raupach, M. R., Regnier, P., Rodenbeck, C., Saito, S., Schwinger, J.,  
13 Segschneider, J., Stocker, B. D., Takahashi, T., Tilbrook, B., van Heuven, S., Viovy, N., Wanninkhof, R.,  
14 Wiltshire, A., and Zaehle, S.: Global carbon budget 2013, *Earth Syst. Sci. Data*, 6, 235-263, 2014.
- 15 Le Quéré, C., Raupach, M. R., Canadell, J. G., Marland, G., Bopp, L., Ciais, P., Conway, T. J., Doney, S. C.,  
16 Feely, R. A., Foster, P., Friedlingstein, P., Gurney, K., Houghton, R. A., House, J. I., Huntingford, C., Levy,  
17 P. E., Lomas, M. R., Majkut, J., Metz, N., Ometto, J. P., Peters, G. P., Prentice, I. C., Randerson, J. T.,  
18 Running, S. W., Sarmiento, J. L., Schuster, U., Sitch, S., Takahashi, T., Viovy, N., van der Werf, G. R., and  
19 Woodward, F. I.: Trends in the sources and sinks of carbon dioxide, *Nature Geoscience*, 2, 831-836,  
20 2009.
- 21 Li, W., Ciais, P., Peng, S., Yue, C., Wang, Y., Thurner, M., Saatchi, S. S., Arneeth, A., Avitabile, V., Carvalhais,  
22 N., Harper, A. B., Kato, E., Koven, C., Liu, Y. Y., Nabel, J. E. M. S., Pan, Y., Pongratz, J., Poulter, B., Pugh,  
23 T. A. M., Santoro, M., Sitch, S., Stocker, B. D., Viovy, N., Wiltshire, A., Yousefpour, R., and Zaehle, S.:  
24 Land-use and land-cover change carbon emissions between 1901 and 2012 constrained by biomass  
25 observations, *Biogeosciences Discuss.*, 2017, 1-25, 2017.
- 26 Liu, Z., Guan, D., Wei, W., Davis, S. J., Ciais, P., Bai, J., Peng, S., Zhang, Q., Hubacek, K., Marland, G., Andres,  
27 R. J., Crawford-Brown, D., Lin, J., Zhao, H., Hong, C., Boden, T. A., Feng, K., Peters, G. P., Xi, F., Liu, J., Li,  
28 Y., Zhao, Y., Zeng, N., and He, K.: Reduced carbon emission estimates from fossil fuel combustion and  
29 cement production in China, *Nature*, 524, 335-338, 2015.
- 30 Manning, A. C. and Keeling, R. F.: Global oceanic and land biotic carbon sinks from the Scripps atmospheric  
31 oxygen flask sampling network, *Tellus Series B-Chemical and Physical Meteorology*, 58, 95-116, 2006.
- 32 Marland, G.: Uncertainties in accounting for CO<sub>2</sub> from fossil fuels, *Journal of Industrial Ecology*, 12, 136-  
33 139, 2008.
- 34 Marland, G., Hamal, K., and Jonas, M.: How Uncertain Are Estimates of CO<sub>2</sub> Emissions?, *Journal of*  
35 *Industrial Ecology*, 13, 4-7, 2009.
- 36 Marland, G. and Rotty, R. M.: Carbon-Dioxide Emissions from Fossil-Fuels - a Procedure for Estimation and  
37 Results for 1950-1982, *Tellus Series B-Chemical and Physical Meteorology*, 36, 232-261, 1984.
- 38 Masarie, K. A. and Tans, P. P.: Extension and integratino of atmospheric carbon dioxide data into a globally  
39 consistent measurement record, *Journal of Geophysical Research-Atmospheres*, 100, 11593-11610,  
40 1995.
- 41 MCI: Ministry of Commerce and Industry, 2017: Foreign Trade Data Dissemination Portal.  
42 <http://121.241.212.146/> (Last Accessed: September 2017), 2017.
- 43 McNeil, B. I., Matear, R. J., Key, R. M., Bullister, J. L., and Sarmiento, J. L.: Anthropogenic CO<sub>2</sub> uptake by the  
44 ocean based on the global chlorofluorocarbon data set, *Science*, 299, 235-239, 2003.
- 45 Melton, J. R. and Arora, V. K.: Competition between plant functional types in the Canadian Terrestrial  
46 Ecosystem Model (CTEM) v. 2.0, *Geosci. Model Dev.*, 9, 323-361, 2016.
- 47 Mercado, L. M., Bellouin, N., Sitch, S., Boucher, O., Huntingford, C., Wild, M., and Cox, P. M.: Impact of  
48 changes in diffuse radiation on the global land carbon sink, *Nature*, 458, 1014-1018, 2009.
- 49 Mikaloff Fletcher, S. E., Gruber, N., Jacobson, A. R., Doney, S. C., Dutkiewicz, S., Gerber, M., Follows, M.,  
50 Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., and Sarmiento, J. L.: Inverse estimates



- 1 of anthropogenic CO<sub>2</sub> uptake, transport, and storage by the oceans, *Global Biogeochemical Cycles*, 20,  
2 GB2002, 2006.
- 3 Millar, R. J., Fuglestedt, J. S., Friedlingstein, P., Rogelj, J., Grubb, M. J., Matthews, H. D., Skeie, R. B.,  
4 Forster, P. M., Frame, D. J., and Allen, A. R.: Emission budgets and pathways consistent with limiting  
5 warming to 1.5 degrees C, *Nature Geoscience*, 10, 741-+, 2017.
- 6 Ministry of Mines: Ministry of Mines, 2017: Mineral Production. <http://mines.nic.in/>. (Last Accessed:  
7 September 2017)
- 8 Myhre, G., Alterskjær, K., and Lowe, D.: A fast method for updating global fossil fuel carbon dioxide  
9 emissions, *Environmental Research Letters*, 4, 034012, 2009.
- 10 Narayanan, B., Aguiar, A., and McDougall, R.:  
11 <https://www.gtap.agecon.purdue.edu/databases/v9/default.asp>, last access: September 2015.
- 12 NBS: National Bureau of Statistics of China (NBS), 2017. "Value added in Industrial enterprises above the  
13 reporting limit grew 7.6% in August 2017" ("2017年6月份规模以上工业增加值增长7.6%").  
14 [http://www.stats.gov.cn/tjsj/zxfb/201707/t20170717\\_1513524.html](http://www.stats.gov.cn/tjsj/zxfb/201707/t20170717_1513524.html). (last access: 15 September  
15 2016), 2017.
- 16 NEA: National Energy Administration of China (NEA), 2017. News conference on the energy situation in the  
17 first half of 2017. [http://mp.weixin.qq.com/s/DIMA4Zod2y\\_nG8pely\\_iQw](http://mp.weixin.qq.com/s/DIMA4Zod2y_nG8pely_iQw). (Last Accessed: 15  
18 September 2017), 2017.
- 19 NOAA/ESRL: [http://www.esrl.noaa.gov/gmd/ccgg/about/global\\_means.html](http://www.esrl.noaa.gov/gmd/ccgg/about/global_means.html), last access: 7 October 2015.
- 20 OEA: Office of the Economic Advisor (OEA), 2017: Index of Eight Core Industries. Office of the Economic  
21 Advisor. <http://eaindustry.nic.in/home.asp>, 2017.
- 22 Oleson, K., Lawrence, D., Bonan, G., Drewniak, B., Huang, M., Koven, C., Levis, S., Li, F., Riley, W., Subin, Z.,  
23 Swenson, S., Thornton, P., Bozbiyik, A., Fisher, R., Heald, C., Kluzek, E., Lamarque, J., Lawrence, P.,  
24 Leung, L., Lipscomb, W., Muszala, S., Ricciuto, D., Sacks, W., Tang, J., and Yang, Z.: Technical  
25 Description of version 4.5 of the Community Land Model (CLM), NCAR, 2013.
- 26 Paulsen, H., Ilyina, T., Six, K. D., and Stemmler, I.: Incorporating a prognostic representation of marine  
27 nitrogen fixers into the global ocean biogeochemical model HAMOCC, *Journal of Advances in  
28 Modeling Earth Systems*, 9, 438-464, 2017.
- 29 Peters, G. P., Andrew, R., and Lennox, J.: Constructing a multi-regional input-output table using the GTAP  
30 database, *Economic Systems Research*, 23, 131-152, 2011a.
- 31 Peters, G. P., Andrew, R. M., Boden, T., Canadell, J. G., Ciais, P., Le Quéré, C., Marland, G., Raupach, M. R.,  
32 and Wilson, C.: The challenge to keep global warming below 2°C, *Nature Climate Change*, 3, 4-6, 2013.
- 33 Peters, G. P., Davis, S. J., and Andrew, R.: A synthesis of carbon in international trade, *Biogeosciences*, 9,  
34 3247-3276, 2012a.
- 35 Peters, G. P., Le Quéré, C., Andrew, R. M., Canadell, J. G., Friedlingstein, P., Ilyina, T., Jackson, R. B.,  
36 Korsbakken, J. I., McKinley, G., Sitch, S., and Tans, P.: Towards real-time verification of carbon dioxide  
37 emissions, *Nature Climate Change*, doi: 10.1038/s41558-017-0013-9, 2017. 2017.
- 38 Peters, G. P., Marland, G., Le Quéré, C., Boden, T. A., Canadell, J. G., and Raupach, M. R.: Correspondence:  
39 Rapid growth in CO<sub>2</sub> emissions after the 2008-2009 global financial crisis, *Nature Climate Change*, 2, 2-  
40 4, 2012b.
- 41 Peters, G. P., Minx, J. C., Weber, C. L., and Edenhofer, O.: Growth in emission transfers via international  
42 trade from 1990 to 2008, *Proceedings of the National Academy of Sciences of the United States of  
43 America*, 108, 8903-8908, 2011b.
- 44 Pfeil, B., Olsen, A., Bakker, D. C. E., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., Metzl, N.,  
45 Sabine, C. L., Akl, J., Alin, S. R., Bates, N., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J.,  
46 Chavez, F. P., Chen, A., Cosca, C., Fassbender, A. J., Feely, R. A., González-Dávila, M., Goyet, C., Hales,  
47 B., Hardman-Mountford, N., Heinze, C., Hood, M., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M.,  
48 Johannessen, T., Jones, S. D., Key, R. M., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N.,  
49 Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate,  
50 A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K., Perez, F. F.,  
51 Pierrot, D., Poisson, A., Ríos, A. F., Santana-Casiano, J. M., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R.,



- 1 Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K.,  
2 Telszewski, M., Thomas, H., Tilbrook, B., Tjiputra, J., Vandemark, D., Veness, T., Wanninkhof, R.,  
3 Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: A uniform, quality controlled Surface  
4 Ocean CO<sub>2</sub> Atlas (SOCAT) A uniform, quality controlled Surface Ocean CO<sub>2</sub> Atlas (SOCAT), Earth Syst.  
5 Sci. Data, 5, 125-143, 2013.
- 6 Pongratz, J., Reick, C. H., Houghton, R. A., and House, J. I.: Terminology as a key uncertainty in net land use  
7 and land cover change carbon flux estimates, Earth System Dynamics, 5, 177-195, 2014.
- 8 PPAC: PPAC, 2017: Natural Gas. Petroleum Planning and Analysis Cell, Ministry of Petroleum and Natural  
9 Gas. <http://eaindstry.nic.in/home.asp> (Last Accessed: September 2017), 2017a.
- 10 PPAC: PPAC, 2017: Petroleum. Petroleum Planning and Analysis Cell, Ministry of Petroleum and Natural  
11 Gas. <http://eaindstry.nic.in/home.asp> (Last Accessed: September 2017), 2017b.
- 12 Prentice, I. C., Farquhar, G. D., Fasham, M. J. R., Goulden, M. L., Heimann, M., Jaramillo, V. J., Khesghi, H.  
13 S., Le Quéré, C., Scholes, R. J., and Wallace, D. W. R.: The Carbon Cycle and Atmospheric Carbon  
14 Dioxide. In: Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third  
15 Assessment Report of the Intergovernmental Panel on Climate Change, Houghton, J. T., Ding, Y.,  
16 Griggs, D. J., Noguer, M., van der Linden, P. J., Dai, X., Maskell, K., and Johnson, C. A. (Eds.), Cambridge  
17 University Press, Cambridge, United Kingdom and New York, NY, USA., 2001.
- 18 Raupach, M. R., Marland, G., Ciais, P., Le Quéré, C., Canadell, J. G., Klepper, G., and Field, C. B.: Global and  
19 regional drivers of accelerating CO<sub>2</sub> emissions, Proceedings of the National Academy of Sciences of the  
20 United States of America, 104, 10288-10293, 2007.
- 21 Regnier, P., Friedlingstein, P., Ciais, P., Mackenzie, F. T., Gruber, N., Janssens, I. A., Laruelle, G. G.,  
22 Lauerwald, R., Luyssaert, S., Andersson, A. J., Arndt, S., Arnosti, C., Borges, A. V., Dale, A. W., Gallego-  
23 Sala, A., Goddérís, Y., Goossens, N., Hartmann, J., Heinze, C., Ilyina, T., Joos, F., La Rowe, D. E., Leifeld,  
24 J., Meysman, F. J. R., Munhoven, G., Raymond, P. A., Spahni, R., Suntharalingam, P., and Thullner M.:  
25 Anthropogenic perturbation of the carbon fluxes from land to ocean, Nature Geoscience, 6, 597-607,  
26 2013.
- 27 Reick, C. H., T. Raddatz, V. Brovkin, and Gayler, V.: The representation of natural and anthropogenic land  
28 cover change in MPI-ESM, Journal of Advances in Modeling Earth Systems, 5, 459-482, 2013.
- 29 Rhein, M., Rintoul, S. R., Aoki, S., Campos, E., Chambers, D., Feely, R. A., Gulev, S., Johnson, G. C., Josey, S.  
30 A., Kostianoy, A., Mauritzen, C., Roemmich, D., Talley, L. D., and Wang, F.: Chapter 3: Observations:  
31 Ocean. In: Climate Change 2013 The Physical Science Basis, Cambridge University Press, 2013.
- 32 Rödenbeck, C.: Estimating CO<sub>2</sub> sources and sinks from atmospheric mixing ratio measurements using a  
33 global inversion of atmospheric transport, Max Plank Institute, MPI-BGC, 2005.
- 34 Rödenbeck, C., Bakker, D. C. E., Gruber, N., Iida, Y., Jacobson, A. R., Jones, S., Landschützer, P., Metzl, N.,  
35 Nakaoka, S., Olsen, A., Park, G. H., Peylin, P., Rodgers, K. B., Sasse, T. P., Schuster, U., Shutler, J. D.,  
36 Valsala, V., Wanninkhof, R., and Zeng, J.: Data-based estimates of the ocean carbon sink variability –  
37 first results of the Surface Ocean CO<sub>2</sub> Mapping intercomparison (SOCOM),  
38 Biogeosciences, 12, 7251-7278, 2015.
- 39 Rödenbeck, C., Bakker, D. C. E., Metzl, N., Olsen, A., Sabine, C., Cassar, N., Reum, F., Keeling, R. F., and  
40 Heimann, M.: Interannual sea–air CO<sub>2</sub> flux variability from an observation-driven ocean  
41 mixed-layer scheme, Biogeosciences, 11, 4599-4613, 2014.
- 42 Rödenbeck, C., Keeling, R. F., Bakker, D. C. E., Metzl, N., Olsen, A., Sabine, C., and Heimann, M.: Global  
43 surface-ocean pCO<sub>2</sub> and sea–air CO<sub>2</sub> flux variability from an observation-driven ocean mixed-layer  
44 scheme, Ocean Science, doi: 10.5194/os-9-193-2013, 2013.
- 45 Rödenbeck, C., S. Houweling, M. Gloor, and M. Heimann: CO<sub>2</sub> flux history 1982–2001 inferred from  
46 atmospheric data using a global inversion of atmospheric transport, Atmos. Chem. Phys. Discuss., 3,  
47 1919-1964, 2003.
- 48 Rogelj, J., Schaeffer, M., Friedlingstein, P., Gillett, N. P., van Vuuren, D. P., Riahi, K., Allen, M., and Knutti,  
49 R.: Differences between carbon budget estimates unravelled, Nature Climate Change, 6, 245-252,  
50 2016.



- 1 Rypdal, K., Paciomik, N., Eggleston, S., Goodwin, J., Irving, W., Penman, J., and Woodfield, M.: Chapter 1  
2 Introduction to the 2006 Guidelines. In: 2006 IPCC Guidelines for National Greenhouse Gas  
3 Inventories, Eggleston, S., Buendia, L., Miwa, K., Ngara, T., and Tanabe, K. (Eds.), Institute for Global  
4 Environmental Strategies (IGES), Hayama, Kanagawa, Japan, 2006.
- 5 SCCL: Singareni Collieries Company Limited (SCCL), 2017: Provisional Production and Dispatches  
6 Performance. Singareni Collieries Company Limited .  
7 [https://sclmines.com/sclnew/performance\\_production.asp](https://sclmines.com/sclnew/performance_production.asp) . (Last Accessed: September 2017),  
8 2017.
- 9 Schimel, D., Alves, D., Enting, I., Heimann, M., Joos, F., Raynaud, D., Wigley, T., Prater, M., Derwent, R.,  
10 Ehhalt, D., Fraser, P., Sanhueza, E., Zhou, X., Jonas, P., Charlson, R., Rodhe, H., Sadasivan, S., Shine, K.  
11 P., Fouquart, Y., Ramaswamy, V., Solomon, S., Srinivasan, J., Albritton, D., Derwent, R., Isaksen, I., Lal,  
12 M., and Wuebbles, D.: Radiative Forcing of Climate Change. In: Climate Change 1995 The Science of  
13 Climate Change. Contribution of Working Group I to the Second Assessment Report of the  
14 Intergovernmental Panel on Climate Change, Houghton, J. T., Meira Rilho, L. G., Callander, B. A.,  
15 Harris, N., Kattenberg, A., and Maskell, K. (Eds.), Cambridge University Press, Cambridge, United  
16 Kingdom and New York, NY, USA., 1995.
- 17 Schwietzke, S., Sherwood, O. A., Bruhwiler, L. M. P., Miller, J. B., Etiope, G., Dlugokencky, E. J., Michel, S. E.,  
18 Arling, V. A., Vaughn, B. H., White, J. W. C., and Tans, P. P.: Upward revision of global fossil fuel  
19 methane emissions based on isotope database, *Nature*, 538, 88-91, 2016.
- 20 Schwinger, J., Goris, N., Tjiputra, J. F., Kriest, I., Bentsen, M., Bethke, I., Ilicak, M., Assmann, K. M., and  
21 Heinze, C.: Evaluation of NorESM-OC (versions 1 and 1.2), the ocean carbon-cycle stand-alone  
22 configuration of the Norwegian Earth System Model (NorESM1), *Geosci. Model Dev.*, 9, 2589-2622,  
23 2016.
- 24 Séférian, R., Bopp, L., Gehlen, M., Orr, J., Ethé, C., Cadule, P., Aumont, O., Salas y Mélia, D., Voltaire, A.  
25 and Madec, G.: Skill assessment of three earth system models with common marine biogeochemistry,  
26 *Climate Dynamics*, 40, 2549–2573, 2013.
- 27 Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W.,  
28 Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and  
29 terrestrial carbon cycling in the LPJ dynamic global vegetation model *Global Change Biology*, 9, 161-  
30 185, 2003.
- 31 Smith, B., Warlind, D., Arneth, A., Hickler, T., Leadley, P., Siltberg, J., and Zaehle, S.: Implications of  
32 incorporating N cycling and N limitations on primary production in an individual-based dynamic  
33 vegetation model, *Biogeosciences*, 11, 2027-2054, 2014.
- 34 Stephens, B. B., Gurney, K. R., Tans, P. P., Sweeney, C., Peters, W., Bruhwiler, L., Ciais, P., Ramonet, M.,  
35 Bousquet, P., Nakazawa, T., Aoki, S., Machida, T., Inoue, G., Vinnichenko, N., Lloyd, J., Jordan, A.,  
36 Heimann, M., Shibistova, O., Langenfelds, R. L., Steele, L. P., Francey, R. J., and Denning, A. S.: Weak  
37 northern and strong tropical land carbon uptake from vertical profiles of atmospheric CO<sub>2</sub>, *Science*,  
38 316, 1732-1735, 2007.
- 39 Stocker, T., Qin, D., and Plattner, G.-K.: *Climate Change 2013 The Physical Science Basis*, Cambridge  
40 University Press, 2013.
- 41 Swart, N. C., Fyfe, J. C., Saenko, O. A., and Eby, M.: Wind-driven changes in the ocean carbon sink,  
42 *Biogeosciences*, 11, 6107-6117, 2014.
- 43 Tian, H. Q., Chen, G. S., Lu, C. Q., Xu, X. F., Hayes, D. J., Ren, W., Pan, S. F., Huntzinger, D. N., and Wofsy, S.  
44 C.: North American terrestrial CO<sub>2</sub> uptake largely offset by CH<sub>4</sub> and N<sub>2</sub>O emissions: toward a full  
45 accounting of the greenhouse gas budget, *Climatic Change*, 129, 413-426, 2015.
- 46 UN: United Nations Statistics Division: Energy Statistics, available at: <http://unstats.un.org/unsd/energy/>  
47 (last access: June 2017), 2017.
- 48 UN: United Nations Statistics Division: National Accounts Main Aggregates Database, available at:  
49 <http://unstats.un.org/unsd/snaama/Introduction.asp> (last access February 2017), 2016.



- 1 UNFCCC: National Inventory Submissions, available at:  
2 [http://unfccc.int/national\\_reports/annex\\_i\\_ghg\\_inventories/national\\_inventories\\_submissions/items](http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/10116.php)  
3 [/10116.php](http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/10116.php) (last access: June 2017), 2017.
- 4 USGS: 2014 Minerals Yearbook - Cement, US Geological Survey, Reston, Virginia, 2017.
- 5 van der Laan-Luijckx, I. T., van der Velde, I. R., van der Veen, E., Tsuruta, A., Stanislawski, K.,  
6 Babenhauserheide, A., Zhang, H. F., Liu, Y., He, W., Chen, H. L., Masarie, K. A., Krol, M. C., and Peters,  
7 W.: The CarbonTracker Data Assimilation Shell (CTDAS) v1.0: implementation and global carbon  
8 balance 2001-2015, *Geoscientific Model Development*, 10, 2785-2800, 2017.
- 9 van der Velde, I. R., Miller, J. B., Schaefer, K., van der Werf, G. R., Krol, M. C., and Peters, W.: Terrestrial  
10 cycling of  $\text{CO}_2$  by photosynthesis, respiration, and biomass burning in  
11 SiBCASA, *Biogeosciences*, 11, 6553-6571, 2014.
- 12 van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P., Morton, D. C., DeFries,  
13 R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation,  
14 savanna, forest, agricultural, and peat fires (1997–2009), *Atmospheric Chemistry and Physics*, 10,  
15 11707-11735, 2010.
- 16 van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van  
17 Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions  
18 estimates during 1997–2015, *Earth Syst. Sci. Data Discuss.*, 2017, 1-43, 2017.
- 19 Viovy, N.: CRUNCEP data set, available at:  
20 [ftp://nacp.ornl.gov/synthesis/2009/frescati/temp/land\\_use\\_change/original/readme.htm](ftp://nacp.ornl.gov/synthesis/2009/frescati/temp/land_use_change/original/readme.htm) (last  
21 access: June 2016), 2016.
- 22 Walker, A. P., Quaipe, T., van Bodegom, P. M., De Kauwe, M. G., Keenan, T. F., Joiner, J., Lomas, M. R.,  
23 MacBean, N., Xu, C. G., Yang, X. J., and Woodward, F. I.: The impact of alternative trait-scaling  
24 hypotheses for the maximum photosynthetic carboxylation rate ( $V_{\text{cmax}}$ ) on global gross primary  
25 production, *New Phytologist*, 215, 1370-1386, 2017.
- 26 Wanninkhof, R., Park, G.-H., Takahashi, T., Sweeney, C., Feely, R. A., Nojiri, Y., Gruber, N., Doney, S. C.,  
27 McKinley, G. A., Lenton, A., Le Quéré, C., Heinze, C., Schwinger, J., Graven, H. D., and Khatiwala, S.:  
28 Global ocean carbon uptake: magnitude, variability and trends, *Biogeosciences*, 10, 1983-2000, 2013.
- 29 Watson, R. T., Rodhe, H., Oeschger, H., and Siegenthaler, U.: Greenhouse Gases and Aerosols. In: *Climate*  
30 *Change: The IPCC Scientific Assessment*. Intergovernmental Panel on Climate Change (IPCC),  
31 Houghton, J. T., Jenkins, G. J., and Ephraums, J. J. (Eds.), Cambridge University Press, Cambridge, 1990.
- 32 Wenzel, S., Cox, P. M., Eyring, V., and Friedlingstein, P.: Projected land photosynthesis constrained by  
33 changes in the seasonal cycle of atmospheric  $\text{CO}_2$ , *Nature*, 538, 499-501, 2016.
- 34 Wilkenskjeld, S., Kloster, S., Pongratz, J., Raddatz, T., and Reick, C. H.: Comparing the influence of net and  
35 gross anthropogenic land-use and land-cover changes on the carbon cycle in the MPI-ESM,  
36 *Biogeosciences*, 11, 4817-4828, 2014.
- 37 Woodward, F. I. and Lomas, M. R.: Vegetation dynamics - simulating responses to climatic change, *Biol.*  
38 *Rev.*, 79, 643-670, 2004.
- 39 Woodward, F. I., Smith, T. M., and Emanuel, W. R.: A global land primary productivity and phytogeography  
40 model, *Global Biogeochemical Cycles*, 9, 471-490, 1995.
- 41 Xi, F., Davis, S. J., Ciais, P., Crawford-Brown, D., Guan, D., Pade, C., Shi, T., Syddall, M., Lv, J., Ji, L., Bing, L.,  
42 Wang, J., Wei, W., Yang, K.-H., Lagerblad, B., Galan, I., Andrade, C., Zhang, Y., and Liu, Z.: Substantial  
43 global carbon uptake by cement carbonation, *Nature Geosci*, 9, 880-883, 2016.
- 44 Zaehle, S., Ciais, P., Friend, A. D., and Prieur, V.: Carbon benefits of anthropogenic reactive nitrogen offset  
45 by nitrous oxide emissions, *Nature Geosci*, 4, 601-605, 2011.
- 46 Zaehle, S. and Friend, A. D.: Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model  
47 description, site-scale evaluation, and sensitivity to parameter estimates, *Global Biogeochemical*  
48 *Cycles*, 24, GB1005, 2010.
- 49 Zscheischler, J., Mahecha, M. D., Avitabile, V., Calle, L., Carvalhais, N., Ciais, P., Gans, F., Gruber, N.,  
50 Hartmann, J., Herold, M., Ichii, K., Jung, M., Landschützer, P., Laruelle, G. G., Lauerwald, R., Papale, D.,  
51 Peylin, P., Poulter, B., Ray, D., Regnier, P., Rödenbeck, C., Roman-Cuesta, R. M., Schwalm, C.,



1 Tramontana, G., Tyukavina, A., Valentini, R., van der Werf, G., West, T. O., Wolf, J. E., and Reichstein,  
2 M.: Reviews and syntheses: An empirical spatiotemporal description of the global surface–  
3 atmosphere carbon fluxes: opportunities and data limitations, *Biogeosciences*, 14, 3685-3703, 2017.

4

5

6 **Tables**

7 **Table 1.** Factors used to convert carbon in various units (by convention, Unit 1 = Unit 2  
8 conversion).

Unit 1	Unit 2	Conversion	Source
GtC (gigatonnes of carbon)	ppm (parts per million) <sup>a</sup>	2.12 <sup>b</sup>	Ballantyne et al. (2012)
GtC (gigatonnes of carbon)	PgC (petagrams of carbon)	1	SI unit conversion
GtCO <sub>2</sub> (gigatonnes of carbon dioxide)	GtC (gigatonnes of carbon)	3.664	44.01/12.011 in mass equivalent
GtC (gigatonnes of carbon)	MtC (megatonnes of carbon)	1000	SI unit conversion

9 <sup>a</sup>Measurements of atmospheric CO<sub>2</sub> concentration have units of dry-air mole fraction. ‘ppm’ is an  
10 abbreviation for micromole/mol, dry air.

11 <sup>b</sup>The use of a factor of 2.12 assumes that all the atmosphere is well mixed within one year. In reality, only  
12 the troposphere is well mixed and the growth rate of CO<sub>2</sub> concentration in the less well-mixed stratosphere  
13 is not measured by sites from the NOAA network. Using a factor of 2.12 makes the approximation that the  
14 growth rate of CO<sub>2</sub> concentration in the stratosphere equals that of the troposphere on a yearly basis.  
15



1 **Table 2.** How to cite the individual components of the global carbon budget presented here.

Component	Primary reference
Global emissions from fossil fuels and industry ( $E_{FF}$ ), total and by fuel type	Boden et al., (2017)
National territorial emissions from fossil fuels and industry ( $E_{FF}$ )	CDIAC source: Boden et al., (2017) UNFCCC (2017)
National consumption-based emissions from fossil fuels and industry ( $E_{FF}$ ) by country (consumption)	Peters et al. (2011b) updated as described in this paper
Land-use change emissions ( $E_{LUC}$ )	average from Houghton and Nassikas (2017) and Hansis et al., (2015), both updated as described in this paper
Growth rate in atmospheric $CO_2$ concentration ( $G_{ATM}$ )	Dlugokencky and Tans (2017)
Ocean and land $CO_2$ sinks ( $S_{OCEAN}$ and $S_{LAND}$ )	This paper for $S_{OCEAN}$ and $S_{LAND}$ and references in Table 5 for individual models.



**Table 3.** Main methodological changes in the global carbon budget since first publication. Unless specified below, the methodology was identical to that described in the current paper. Furthermore, methodological changes introduced in one year are kept for the following years unless noted. Empty cells mean there were no methodological changes introduced that year.

Publication year <sup>a</sup>	Global	Fossil fuel emissions Country (territorial)	Country (consumption)	LUC emissions	Atmosphere	Reservoirs Ocean	Land	Uncertainty & other changes
2006		Split in regions						
Raupach et al. (2007)				$E_{luc}$ based on FAO-FRA 2005; constant $E_{luc}$ for 2006	1959-1979 data from Mauna Loa; data after 1980 from global average	Based on one ocean model tuned to reproduced observed 1990s sink		±1σ provided for all components
Canadell et al. (2007)				Constant $E_{luc}$ for 2007				
2008 (online)				Fire-based emission anomalies used for 2006-2008				
2009		Split between Annex B and non-Annex B	Results from an independent study discussed	$E_{luc}$ updated with FAO-FRA 2010		Based on four ocean models normalised to observations with constant delta	First use of five DGVMs to compare with budget residual	
2010 Friedlingstein et al. (2010)		Emissions for top emitters						
2011			Split between Annex B and non-Annex B					
Peters et al. (2012b)		129 countries from 1959	129 countries and regions from 1990-2010 based on GTPA8.0	$E_{luc}$ for 1997-2011 includes interannual anomalies from fire-based emissions	All years from global average	Based on 5 ocean models normalised to observations with ratio	Ten DGVMs available for $S_{land}$ ; First use of four models to compare with $E_{luc}$	Confidence levels; cumulative emissions; budget from 1750
2012				$E_{luc}$ for 2012 estimated 1990-2011 based on GTPA8.1, with detailed estimates for years 1997, 2001, 2004, and 2007		Based on six models compared with two data-sets to year 2011	Coordinated DGVM experiments for $S_{land}$ and $E_{luc}$	
Peters et al. (2013)				Extended to 2012 with updated GDP data				
2013		250 countries <sup>b</sup>	134 countries and regions from 1990-2011 based on GTPA8.1, with detailed estimates for years 1997, 2001, 2004, and 2007	$E_{luc}$ for 1997-2013 includes interannual anomalies from fire-based emissions				
Le Quéré et al. (2014)								
2014		Three years of BP data	Three years of BP data					
Le Quéré et al. (2015b)								
2015		projection for current year based on Jan-Aug data	National emissions from UNFCCC extended to 2014 also provided	Detailed estimates introduced for 2011 based on GTPA9		Based on eight models	Based on ten models with assessment of minimum realism	Inclusion of breakdown of the sinks in three latitude bands and comparison with three atmospheric inversions
Le Quéré et al. (2015a)								
Jackson et al. (2016)		Two years of BP data	Added three small countries; CHN emissions from 1990 from BP data (this release only)	Preliminary $E_{luc}$ using FRA-2015 shown for comparison; use of five DGVMs		Based on seven models	Based on fourteen models	The decadal uncertainty for the DGVM ensemble mean now uses ±1σ of the decadal spread across models
Le Quéré et al. (2016)								Discussion of projection for full budget for current year
2017 (this study)		projection includes India-specific data		Average of two bookkeeping models; use of twelve DGVMs		Based on eight models that match the observed sink for the 1990s; no longer normalised	Based on fifteen models that meet three criteria (see Sect. 2.5)	Land multi-model average now used in main carbon budget; with the carbon imbalance presented separately; new table of key uncertainties



- 1 <sup>a</sup>The naming convention of the budgets has changed. Up to and including 2010, the budget year (Carbon Budget 2010) represented the latest year of the data. From 2012,
- 2 the budget year (Carbon Budget 2012) refers to the initial publication year.
- 3 <sup>b</sup>The CDIAC database has about 250 countries, but we show data for 219 countries since we aggregate and disaggregate some countries to be consistent with current
- 4 country definitions (see Sect. 2.1.1 for more details).



1 **Table 4a.** Comparison of the processes included (Y) or not (N) in the bookkeeping and Dynamic  
 2 Global Vegetation Models for their estimates of  $E_{LUC}$  and  $S_{LAND}$ . See Table 5 for model references.  
 3 All models include deforestation and forest regrowth after abandonment of agriculture (or from  
 4 afforestation activities on agricultural land).

	bookkeeping models		DGVMs														
	H&N2007	BLUE	CABLE	CLASS-CTEM	CLM4.5(BGC)	DLEM	ISAM	JSBACH <sup>f</sup>	JULES	LPJ-GUESS <sup>i</sup>	LPJ	LPX-Bern	OCN	ORCHIDEE	Orchidee-MICT	SDGVM	VISIT <sup>j</sup>
Processes relevant for $E_{LUC}$																	
Wood harvest and forest degradation <sup>a</sup>	Y	Y	Y	N	Y	Y	Y		N	N	N <sup>d</sup>	Y	Y	N	N		
Shifting cultivation / subgrid scale transitions	N <sup>b</sup>	Y	Y	N	Y	N	N		N	N	N <sup>d</sup>	N	N	N	N		
Cropland harvest	Y <sup>f</sup>	Y <sup>f</sup>	N	L	N	Y	Y		N	Y	Y	Y	Y	Y	Y	Y	Y
Peat fires	Y	Y	N	N	Y	N	N		N	N	N	N	N	N	N	N	N
Fire as a management tool	Y <sup>f</sup>	Y <sup>f</sup>	N	N	N	N	N		N	N	N	N	N	N	N	N	N
N fertilization	Y <sup>f</sup>	Y <sup>f</sup>	N	N	N	Y	Y		N	N	Y	Y	N	N	N	N	N
Tillage	Y <sup>f</sup>	Y <sup>f</sup>	N	Y <sup>f</sup>	N	N	N		N	N	N	N	Y <sup>h</sup>	Y <sup>h</sup>	N	N	N
Irrigation	Y <sup>f</sup>	Y <sup>f</sup>	N	N	N	Y	Y		N	N	N	N	N	N	N	N	N
Wetland drainage	Y <sup>f</sup>	Y <sup>f</sup>	N	N	N	N	N		N	N	N	N	N	N	N	N	N
Erosion	Y <sup>f</sup>	Y <sup>f</sup>	N	N	N	N	N		N	N	N	N	N	N	N	N	N
South East Asia peat drainage	Y	Y	N	N	N	N	N		N	N	N	N	N	N	N	N	N
Grazing and mowing harvest	Y <sup>f</sup>	Y <sup>f</sup>	N	N	N	N	Y		N	Y	N	N	N	N	N	N	N
Processes relevant also for $S_{LAND}$																	
Fire simulation	US only	N	N	Y	Y	Y	N	Y	N	Y	Y	Y	N	N	Y	Y	Y
Climate and variability	N	N	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
CO <sub>2</sub> fertilisation	N <sup>e</sup>	N <sup>e</sup>	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Carbon-nitrogen interactions, including N deposition	N <sup>i</sup>	N <sup>i</sup>	Y	N <sup>e</sup>	Y	Y	Y	N	N	Y	N	Y	Y	N <sup>e</sup>	N	Y <sup>c</sup>	N

5 <sup>a</sup> Refers to the routine harvest of established managed forests rather than pools of harvested products.  
 6 <sup>b</sup> No back- and forth-transitions between vegetation types at the country-level, but if forest loss based on FRA  
 7 exceeded agricultural expansion based on FAO, then this amount of area  
 8 <sup>c</sup> Limited. Nitrogen uptake is simulated as a function of soil C, and  $V_{cmax}$  is an empirical function of canopy N. Does  
 9 not consider N deposition.  
 10 <sup>d</sup> Available but not active for comparability between the two LU forcings.  
 11 <sup>e</sup> Although C-N cycle interactions are not represented, the model includes a parameterization of down-regulation of  
 12 photosynthesis as CO<sub>2</sub> increases to emulate nutrient constraints (Arora et al., 2009)  
 13 <sup>f</sup> Tillage is represented over croplands by increased soil carbon decomposition rate and reduced humification of litter  
 14 to soil carbon.  
 15 <sup>g</sup> Bookkeeping models include effect of CO<sub>2</sub>-fertilization as captured by observed carbon densities, but not as an effect  
 16 transient in time.  
 17 <sup>h</sup> 20% reduction of active SOC pool turnover time for C3 crop and 40% reduction for C4 crops  
 18 <sup>i</sup> Process captured implicitly by use of observed carbon densities.  
 19 <sup>j</sup> Three DGVMs were excluded from the  $E_{LUC}$  estimate due to an initial peak of  $E_{LUC}$  emissions caused by a cold start of  
 20 shifting cultivation in 1860.  
 21  
 22



1 **Table 4b.** Comparison of the processes included in the Global Ocean Biogeochemistry Models for  
 2 their estimates of  $S_{OCEAN}$ . See Table 5 for model references.  
 3

	CCSM-BEC	CSIRO	NorESM-OC	MITgcm-REcoM2	MPIOM-HAMOC	NEMO-PISCES (CNRM)	NEMO-PISCES (IPSL)	NEMO-PlankTOM5
Atmospheric forcing	NCEP	JRA55	CORE-I (spin up) / NCEP with CORE-II corrections	JRA55	ERA-20C	NCEP	NCEP	NCEP
Initialisation of carbon chemistry	GLODAP	GLODAP + spin up 1000+ years	GLODAP v1 + spin up 1000 years	GLODAP, then spin-up 116 years (2 cycles with >1000 yrs JRA55)	from previous model runs with >1000 yrs spinup	spin up 3000 years offline + 300 years online	GLODAP from 1948 onwards	GLODAP + spin up 30 years
Physical ocean model	POP Version 1.4.3	MOM5	MICOM	MITgcm 65n	MPIOM	NEMOv2.4-ORCA1L42	NEMOv3.2-ORCA2L31	NEMOv2.3-ORCA2
Resolution	3.6° lon, 0.8 to 1.8° lat	1° x 1° with enhanced resolution at the tropics and high lat S. Ocean; 50 levels	1° lon, 0.17 to 0.25 lat; 51 isopycnal layers + 2 bulk mixed layer	2° lon, 0.38-2° lat, 30 levels	1.5°; 40 levels	2° lon, 0.3 to 1° lat	2° lon, 0.3 to 1.5° lat; 31 levels	2° lon, 0.3 to 1.5° lat; 31 levels

4

5



1 **Table 4c.** Comparison of the inversion set up and input fields for the atmospheric inversions. See  
2 Table 5 for references.

	CarbonTracker Europe (CTE)	Jena CarboScope	CAMS	3
Version number	CTE2017-FT	s85oc_v4.1s	v16r1	
Observations				
Atmospheric observations	Hourly resolution (well-mixed conditions) OBSPACK GLOBALVIEWplus v2.1 & NRTv3.3 <sup>a</sup>	Flasks and hourly (outliers removed by 2-sigma criterion)	Daily averages of well-mixed conditions - OBSPACK GLOBALVIEWplus v2.1 <sup>a</sup> & NRT v3.2.3, WDCGG, RAMCES and ICOS ATC	
Prior fluxes				
Biosphere and fires	SIBCASA-GFED4s <sup>b</sup>	Zero	ORCHIDEE (climatological), GFEDv4 & GFAS	
Ocean	Ocean inversion by Jacobson et al. (2007)	pCO <sub>2</sub> -based ocean flux product oc_v1.5 (update of Rödenbeck et al., 2014)	Landschützer et al. (2015)	
Fossil fuels	EDGAR+IER, scaled to CDIAC	CDIAC (extended after 2013 with GCP totals)	EDGAR scaled to CDIAC	
Transport and optimization				
Transport model	TM5	TM3	LMZ v5A	
Weather forcing	ECMWF	NCEP	ECMWF	
Resolution (degrees)	Global: 3° x 2°, Europe: 1° x 1°, North America: 1° x 1°	Global: 4° x 5°	Global: 3.75° x 1.875°	
Optimization	Ensemble Kalman filter	Conjugate gradient (re-ortho-normalization)	Variational	

4 <sup>a</sup>(Carbontracker Team, 2017; GLOBALVIEW, 2016)

5 <sup>b</sup>(van der Velde et al., 2014)

6



1 **Table 5.** References for the process models, pCO<sub>2</sub>-based ocean flux products, and atmospheric  
 2 inversions included in Figs. 6-8. All models and products are updated with new data to end of year  
 3 2016.

Model/data name	Reference	Change from Le Quéré et al. (2016)
<i>Bookkeeping models for land-use change emissions</i>		
BLUE	Hansis et al. (2015)	Not applicable (not used in previous carbon budgets)
H&N	Houghton and Nassikas (2017)	updated from Houghton et al. (2012); key differences include Revised land-use change data to FAO2015, revised vegetation carbon densities, Indonesian and Malaysian peat burning and drainage added, removal of shifting cultivation
<i>Dynamic global vegetation models</i>		
CABLE	Haverd et al., (2017)	Optimisation of plant investment in Rubisco- vs electron transport-limited photosynthesis; temperature-dependent onset of spring recovery in evergreen needle-leaves
CLASS-CTEM	Melton and Arora (2016)	A soil colour index is now used to determine soil albedo as opposed to soil texture. Soil albedo still gets modulated by other factors including soil moisture.
CLM4.5(BGC)	Oleson et al. (2013)	No change
DLEM	Tian et al. (2015)	Consideration of the expansion of cropland and pasture, compared with no pasture expansion in previous version.
ISAM	Jain et al. (2013)	No change
JSBACH	Reick et al. (2013) <sup>a</sup>	Adapted the pre-processing of the LUH data; scaling crop and pasture states and transitions with the desert fractions in jsbach in order to maintain as much of the prescribed agricultural areas as possible.
JULES <sup>b</sup>	Clarke et al. (2011) <sup>c</sup>	No Change
LPJ-GUESS	Smith et al. (2014) <sup>d</sup>	LUH2 with land use aggregated to LPJ-GUESS land cover inputs, shifting cultivation based on LUH2 gross transitions matrix, and wood harvest based on LUH2 area fractions of wood harvest; $\alpha_a$ reduction by 15%
LPJ <sup>e</sup>	Sitch et al. (2003) <sup>f</sup>	No change
LPX-Bern	Keller et al., (2017)	Updated model parameter values (Keller et. al. 2017) due to assimilation of observational data.
OCN	Zaehle and Friend (2010) <sup>g</sup>	uses r293, including minor bugfixes; use of the CMIP6 N deposition data set (Hegglin et al. in prep)
ORCHIDEE	Krinner et al. (2005) <sup>h</sup>	improved water stress, new soil albedo, improved snow scheme
ORCHIDEE-MICT	Guimberteau et al. (2017)	new version of ORCHIDEE including fires, permafrost regions coupling between soil thermics and carbon dynamics, managed grasslands
SDGVM	Woodward et al (1995) <sup>i</sup>	Uses Kattge et al. (2009) Vcmax~leaf N relationships (with oxisol relationship for evergreen broadleaves)
VISIT	Kato et al. (2013) <sup>j</sup>	LUH2 is applied for land-use, wood harvest, and land-use change. Sensitivity of soil decomposition parameters from Lloyd and Taylor (1994) are modified.



<i>Global ocean biogeochemistry models</i>		
CCSM-BEC	Doney et al. (2009)	Change in atmospheric CO <sub>2</sub> concentration <sup>k</sup>
CSIRO	Law et al. (2017)	Physical model change from MOM4 to MOM5 and atmospheric forcing from JRA-55
MITgcm-REcoM2	Hauck et al. (2016)	1% iron solubility and atmospheric forcing from JRA-55
MPIOM-HAMOCC <sup>l</sup>	Ilyina et al. (2013)	Cyanobacteria added to HAMOCC (Paulsen et al., 2017)
NEMO-PISCES (CNRM)	Séférian et al. (2013)	No change
NEMO-PISCES (IPSL)	Aumont and Bopp (2006)	No change
NEMO-PlankTOM5	Buitenhuis et al. (2010) <sup>m</sup>	No change
NorESM-OC	Schwinger et al. (2016)	No change
<i>pCO<sub>2</sub>-based flux ocean products</i>		
Landschützer	Landschützer et al. (2016)	No change
Jena CarboScope	Rödenbeck et al. (2014)	Updated to version oc_1.5
<i>Atmospheric inversions</i>		
CarbonTracker Europe (CTE)	van der Laan-Luijkx et al. (2017)	Minor changes in the inversion set up
Jena CarboScope	Rödenbeck et al. (2003)	Prior fluxes, outlier removal, changes in atmospheric observations station suite
CAMS <sup>n</sup>	Chevallier et al. (2005)	Change from half-hourly observations to daily averages of well-mixed conditions

1 <sup>a</sup> See also Goll et al. (2015).2 <sup>b</sup> Joint UK Land Environment Simulator.3 <sup>c</sup> See also Best et al. (2011).4 <sup>d</sup> To account for the differences between the derivation of SWRAD from CRU cloudiness and SWRAD from CRU-NCEP, the photosynthesis scaling parameter  $\alpha_3$  was modified (-15%) to yield similar results.6 <sup>e</sup> Lund-Potsdam-Jena.7 <sup>f</sup> Compared to published version, decreased LPJ wood harvest efficiency so that 50% of biomass was removed off-site compared to 85% used in the 2012 budget. Residue management of managed grasslands increased so that 100% of harvested grass enters the litter pool.10 <sup>g</sup> See also Zaehle et al. (2011).11 <sup>h</sup> Compared to published version, revised parameters values for photosynthetic capacity for boreal forests (following assimilation of FLUXNET data), updated parameters values for stem allocation, maintenance respiration and biomass export for tropical forests (based on literature) and, CO<sub>2</sub> down-regulation process added to photosynthesis.14 <sup>i</sup> See also Woodward & Lomas (2004) and Walker et al. (2017). Changes from publications include sub-daily light downscaling for calculation of photosynthesis and other adjustment.16 <sup>j</sup> See also Ito and Inatomi (2012).17 <sup>k</sup> Previous simulations used atmospheric CO<sub>2</sub> concentration from the IPCC IS92a scenario. This has been re-run using observed atmospheric CO<sub>2</sub> concentration consistent with the protocol used here.19 <sup>l</sup> Last included in Le Quéré et al. (2015)20 <sup>m</sup> With no nutrient restoring below the mixed layer depth.21 <sup>n</sup> See also Supplementary Material (Chevallier, 2015; Hourdin et al., 2006).

22

23

24



**Table 6.** Comparison of results from the bookkeeping method and budget residuals with results from the DGVMs and inverse estimates for different periods, last decade and last year available. All values are in GtC yr<sup>-1</sup>. The DGVM uncertainties represent ±1σ of the decadal or annual (for 2016 only) estimates from the individual DGVMs, for the inverse models all three results are given where available.

	Mean (GtC yr <sup>-1</sup> )						2016
	1960-1969	1970-1979	1980-1989	1990-1999	2000-2009	2007-2016	
<i>Land-use change emissions (E<sub>LUC</sub>)</i>							
Bookkeeping methods	1.4 ± 0.7	1.1 ± 0.7	1.2 ± 0.7	1.3 ± 0.7	1.2 ± 0.7	1.3 ± 0.7	1.3 ± 0.7
DGVMs	1.3 ± 0.5	1.2 ± 0.5	1.2 ± 0.4	1.2 ± 0.3	1.2 ± 0.4	1.3 ± 0.4	1.4 ± 0.8
<i>Terrestrial sink (S<sub>LAND</sub>)</i>							
Residual sink from global budget (E <sub>FF-E<sub>LUC</sub>-G<sub>ATM</sub>-S<sub>OCEAN</sub>)</sub>	1.8 ± 0.9	1.8 ± 0.9	1.5 ± 0.9	2.6 ± 0.9	3.0 ± 0.9	3.6 ± 1.0	2.4 ± 1.0
DGVMs <sup>a</sup>	1.4 ± 0.7	2.4 ± 0.6	2.0 ± 0.6	2.5 ± 0.5	2.9 ± 0.8	3.0 ± 0.8	2.7 ± 1.0
<i>Total land fluxes (S<sub>LAND</sub> - E<sub>LUC</sub>)</i>							
Budget constraint (E <sub>FF-G<sub>ATM</sub>-S<sub>OCEAN</sub>)</sub>	0.4 ± 0.5	0.7 ± 0.6	0.4 ± 0.6	1.3 ± 0.6	1.7 ± 0.6	2.3 ± 0.7	1.1 ± 0.7
DGVMs	0.1 ± 0.9	1.2 ± 0.8	0.7 ± 0.7	1.2 ± 0.5	1.7 ± 0.8	1.7 ± 0.7	1.3 ± 1.0
Inversions (CTE/Jena CarboScope/CAMS)*	-/-/-	-/-/-	-/-/0.2	-/0.6/1.3	1.4/1.1/1.9	1.8/1.4/2.3	0.0/0.0/2.2

<sup>a</sup> Estimates are corrected for the preindustrial influence of river fluxes (Sect. 2.7.2). See Tables 4c & 5 for references.



**Table 7.** Decadal mean in the five components of the anthropogenic CO<sub>2</sub> budget for different periods, and last year available. All values are in GtC yr<sup>-1</sup>, and uncertainties are reported as ±1σ. Unlike previous versions of the Global Carbon Budget, the terrestrial sink (S<sub>LAND</sub>) is now estimated independently from the mean of DGVM models. Therefore the table also shows the budget imbalance (B<sub>IM</sub>), which provides a measure of the discrepancies among the nearly independent estimates and has an uncertainty exceeding ± 1 GtC yr<sup>-1</sup>. A positive imbalance means the emissions are overestimated and/or the sinks are too small.

	Mean (GtC yr <sup>-1</sup> )						
	1960-1969	1970-1979	1980-1989	1990-1999	2000-2009	2007-2016	2016
<i>Emissions</i>							
Fossil fuels and industry (E <sub>FF</sub> )	3.1 ± 0.2	4.7 ± 0.2	5.5 ± 0.3	6.3 ± 0.3	7.8 ± 0.4	9.4 ± 0.5	9.9 ± 0.5
Land-use change emissions (E <sub>LUC</sub> )	1.4 ± 0.7	1.1 ± 0.7	1.2 ± 0.7	1.3 ± 0.7	1.2 ± 0.7	1.3 ± 0.7	1.3 ± 0.7
<i>Partitioning</i>							
Growth rate in atmospheric CO <sub>2</sub> concentration (G <sub>ATM</sub> )	1.7 ± 0.1	2.8 ± 0.1	3.4 ± 0.1	3.1 ± 0.1	4.0 ± 0.1	4.7 ± 0.1	6.1 ± 0.2
Ocean sink (S <sub>OCEAN</sub> )	1.0 ± 0.5	1.3 ± 0.5	1.7 ± 0.5	1.9 ± 0.5	2.1 ± 0.5	2.4 ± 0.5	2.6 ± 0.5
Terrestrial sink (S <sub>LAND</sub> )	1.4 ± 0.7	2.4 ± 0.6	2.0 ± 0.6	2.5 ± 0.5	2.9 ± 0.8	3.0 ± 0.8	2.7 ± 1.0
<i>Budget imbalance</i>							
B <sub>IM</sub> = E <sub>FF</sub> +E <sub>LUC</sub> - (G <sub>ATM</sub> +S <sub>OCEAN</sub> +S <sub>LAND</sub> )	(0.4)	(-0.6)	(-0.4)	(0.1)	(0.0)	(0.6)	(-0.3)



1 **Table 8.** Comparison of the projection with realised emissions from fossil fuels and industry ( $E_{FF}$ ).  
 2 The ‘Actual’ values are first estimate available using actual data, and the ‘Projected’ values refers  
 3 to estimate made before the end of the year for each publication. Projections based on a different  
 4 method from that described here during 2008-2014 are available in Le Quéré et al., (2016). All  
 5 values are adjusted for leap years.  
 6

	World		China		USA		India		Rest of World	
	Projected	Actual								
2015 <sup>a</sup>	-0.6% (-1.6 to 0.5)	0.06%	-3.9% (-4.6 to -1.1)	-0.7%	-1.5% (-5.5 to 0.3)	-2.5%	-	-	1.2% (-0.2 to 2.6)	0.7%
2016 <sup>b</sup>	-0.2% (-1.0 to +1.8)	+0.18%	-0.5% (-3.8 to +1.3)	-0.3%	-1.7% (-4.0 to +0.6)	-2.1%	-	-	+1.0% (-0.4 to +2.5)	0.6%
2017 <sup>c</sup>	+2.0% (+0.8 to +3.0)	-	+3.5 (+0.7 to +5.4)	-	-0.4% (-2.7 to +1.0)	-	+2.0% (+0.2 to +3.8)	-	+1.9% (0.3 to +3.4)	-

7 <sup>a</sup>Jackson et al. (2016) and Le Quéré et al. (2015a). <sup>b</sup>Le Quéré et al., (2016). <sup>c</sup>This study.  
 8



1

2 **Table 9.** Cumulative CO<sub>2</sub> emissions for different time periods in gigatonnes of carbon (GtC). All  
 3 uncertainties are reported as  $\pm 1\sigma$ .  $E_{LUC}$  and  $S_{OCEAN}$  have been revised to incorporate multiple  
 4 estimates (Section 3.5), and unlike previous versions of the Global Carbon Budget, the terrestrial  
 5 sink ( $S_{LAND}$ ) is now estimated independently from the mean of the DGVM. Therefore the table  
 6 also shows the budget imbalance, which provides a measure of the discrepancies among the  
 7 nearly independent estimates. Its uncertainty exceeds  $\pm 60$  GtC. The method used here does not  
 8 capture the loss of additional sink capacity from reduced forest cover, which is about 15 GtC and  
 9 would exacerbate the budget imbalance (see Section 2.7.3). All values are rounded to the  
 10 nearest 5 GtC and therefore columns do not necessarily add to zero.

	Units of GtC	1750-2016	1850-2005	1959-2016	1870-2016	1870-2017 <sup>a</sup>
<i>Emissions</i>						
Fossil fuels and industry ( $E_{FF}$ )		420 $\pm$ 20	320 $\pm$ 15	345 $\pm$ 15	420 $\pm$ 20	430 $\pm$ 20
Land-use change emissions ( $E_{LUC}$ )		225 $\pm$ 75	180 $\pm$ 60	75 $\pm$ 40	180 $\pm$ 60	180 $\pm$ 60
Total emissions		645 $\pm$ 80	500 $\pm$ 60	415 $\pm$ 45	600 $\pm$ 65	610 $\pm$ 65
<i>Partitioning</i>						
Growth rate in atmospheric CO <sub>2</sub> concentration ( $G_{ATM}$ ) <sup>b</sup>		270 $\pm$ 5	200 $\pm$ 5	185 $\pm$ 5	245 $\pm$ 5	250 $\pm$ 5
Ocean sink ( $S_{OCEAN}$ )		160 $\pm$ 20	145 $\pm$ 20	95 $\pm$ 20	145 $\pm$ 20	150 $\pm$ 20
Terrestrial sink ( $S_{LAND}$ ) <sup>c</sup>		205 $\pm$ 55	155 $\pm$ 45	135 $\pm$ 35	190 $\pm$ 45	190 $\pm$ 55
<i>Budget imbalance</i>						
$B_{IM} = E_{FF} + E_{LUC} - (G_{ATM} + S_{OCEAN} + S_{LAND})$		(15)	(0)	(0)	(20)	(20)

11 <sup>a</sup>Using projections for year 2017 (Sect. 3.3).

12 <sup>b</sup>A small change was introduced from Le Quéré et al. (2016) to be consistent with the annual analysis, whereby the  
 13 growth in atmospheric CO<sub>2</sub> concentration is calculated from the difference between concentrations at the end of the  
 14 year (deseasonalised), rather than averaged over the year.

15 <sup>c</sup>Assuming  $S_{LAND}$  increases proportionally to  $G_{ATM}$  prior to 1860 when the DGVM estimates start.

16



1 **Table 10.** Major known sources of uncertainties in each component of the Global Carbon Budget,  
2 defined as input data or processes that have a demonstrated effect of at least 0.3 GtC yr<sup>-1</sup>.

3

Source of uncertainty	Time scale (years)	Location	Status	Evidence
<b>Emissions from fossil fuels and industry (<math>E_{FF}</math>; Section 2.1)</b>				
energy statistics	annual to decadal	mainly China	see Sect. 2.1	(Korsbakken et al., 2016)
carbon content of coal	decadal	mainly China	see Sect. 2.1	(Liu et al., 2015)
<b>Emissions from land-use change (<math>E_{LUC}</math>; section 2.2)</b>				
land-cover and land-use change statistics	continuous	global	see Sect. 2.2	(Houghton et al., 2012)
sub-grid-scale transitions	annual to decadal	global; in particular tropics	see Table 5	(Wilkenskjeld et al., 2014)
vegetation biomass	annual to decadal	global; in particular tropics	see Table 5	(Houghton et al., 2012)
wood and crop harvest	annual to decadal	global	see Table 5	(Arneeth et al., 2017)
peat burning <sup>a</sup>	multi-decadal trend	global; SE Asia	see Table 5	(van der Werf et al., 2010)
loss of additional sink capacity	multi-decadal trend	global	not included; Section 2.7.3	(Gitz and Ciais, 2003)
Atmospheric growth rate ( $G_{ATM}$ ) → no demonstrated uncertainties larger than $\pm 0.3$ GtC yr <sup>-1, b</sup>				
<b>Ocean sink (<math>S_{OCEAN}</math>)</b>				
variability in oceanic circulation <sup>c</sup>	semi-decadal to decadal	global; in particular Southern Ocean	see Sect. 2.4.2	(DeVries et al., 2017)
anthropogenic changes in nutrient supply	multi-decadal trend	global	not included	(Duce et al., 2008)
<b>Land sink (<math>S_{LAND}</math>)</b>				
strength of CO <sub>2</sub> fertilisation	multi-decadal trend	global	see Sect. 2.5	(Wenzel et al., 2016)
response to variability in temperature and rainfall	annual to decadal	global; in particular tropics	see Sect. 2.5	(Cox et al., 2013)
nutrient limitation and supply	multi-decadal trend	global	see Sect. 2.5	(Zaehle et al., 2011)
response to diffuse radiation	annual	global	see Sect. 2.5	(Mercado et al., 2009)


4 <sup>a</sup>As result of interactions between land-use and climate

5 <sup>b</sup>The uncertainties in  $G_{ATM}$  have been estimated as  $\pm 0.2$  GtC yr<sup>-1</sup>, although the conversion of the growth rate into a  
6 global annual flux assuming instantaneous mixing throughout the atmosphere introduces additional errors that have  
7 not yet been quantified.

8 <sup>c</sup>Could in part be due to uncertainties in atmospheric forcing (Swart et al., 2014)




1 **Figure Captions**



2

3 **Figure 1.** Surface average atmospheric CO<sub>2</sub> concentration, deseasonalised (ppm). The 1980-2017  
 4 monthly data are from NOAA/ESRL (Dlugokencky and Tans, 2017) and are based on an average of  
 5 direct atmospheric CO<sub>2</sub> measurements from multiple stations in the marine boundary layer  
 6 (Masarie and Tans, 1995). The 1958-1979 monthly data are from the Scripps Institution of  
 7 Oceanography, based on an average of direct atmospheric CO<sub>2</sub> measurements from the Mauna  
 8 Loa and South Pole stations (Keeling et al., 1976). To take into account the difference of mean CO<sub>2</sub>  
 9 between the NOAA/ESRL and the Scripps station networks used here, the Scripps surface average  
 10 (from two stations) was harmonised to match the NOAA/ESRL surface average (from multiple  
 11 stations) by adding the mean difference of 0.542 ppm, calculated here from overlapping data  
 12 during 1980-2012. The mean seasonal cycle is also shown from 1980 (in pink).

13




1  
 2 **Figure 2.** Schematic representation of the overall perturbation of the global carbon cycle caused  
 3 by anthropogenic activities, averaged globally for the decade 2007-2016. The arrows represent  
 4 emission from fossil fuels and industry ( $E_{FF}$ ); emissions from deforestation and other land-use  
 5 change ( $E_{LUC}$ ); the growth rate in atmospheric  $CO_2$  concentration ( $G_{ATM}$ ) and the uptake of carbon  
 6 by the ‘sinks’ in the ocean ( $S_{OCEAN}$ ) and land ( $S_{LAND}$ ) reservoirs. The budget imbalance ( $B_{IM}$ ) is also  
 7 shown. All fluxes are in units of  $GtC\ yr^{-1}$ , with uncertainties reported as  $\pm 1\sigma$  (68% confidence that  
 8 the real value lies within the given interval) as described in the text. This figure is an update of one  
 9 prepared by the International Geosphere Biosphere Programme for the GCP, using diagrams  
 10 created with symbols from the Integration and Application Network, University of Maryland  
 11 Center for Environmental Science ([ian.umces.edu/symbols/](http://ian.umces.edu/symbols/)), first presented in Le Quéré (2009).

12




1



2


3 **Figure 3.** Combined components of the global carbon budget illustrated in Fig. 2 as a function of  
 4 time, for emissions from fossil fuels and industry ( $E_{FF}$ ; grey) and emissions from land-use change  
 5 ( $E_{LUC}$ ; brown), as well as their partitioning among the atmosphere ( $G_{ATM}$ ; purple), land ( $S_{LAND}$ ;  
 6 green) and oceans ( $S_{OCEAN}$ ; dark blue). The partitioning is based on nearly independent estimates  
 7 from observations (for  $G_{ATM}$ ) and from process model ensembles constrained by data (for  $S_{OCEAN}$   
 8 and  $S_{LAND}$ ), and does not exactly add up to the sum of the emissions, resulting in a budget  
 9 imbalance which is reflected in the difference between the bottom red line and the sum of the  
 10 ocean, land and atmosphere. All time series are in  $\text{GtC yr}^{-1}$ .  $G_{ATM}$  and  $S_{OCEAN}$  prior to 1959 are  
 11 based on different methods.  $E_{FF}$  are primarily from Boden et al. (2017), with uncertainty of about  
 12  $\pm 5\%$  ( $\pm 1\sigma$ );  $E_{LUC}$  are from two bookkeeping models (Table 2) with uncertainties of about  $\pm 50\%$ ;  
 13  $G_{ATM}$  prior to 1959 is from Joos and Spahni (2008) with uncertainties equivalent to about  $\pm 0.1$ - $0.15$   
 14  $\text{GtC yr}^{-1}$ , and from Dlugokencky and Tans (2017) from 1959 with uncertainties of about  $\pm 0.2$   $\text{GtC}$   
 15  $\text{yr}^{-1}$ ;  $S_{OCEAN}$  prior to 1959 is averaged from Khatiwala et al. (2013) and DeVries (2014) with  
 16 uncertainty of about  $\pm 30\%$ , and from a multi-model mean (Table 5) from 1959 with uncertainties  
 17 of about  $\pm 0.5$   $\text{GtC yr}^{-1}$ ;  $S_{LAND}$  is a multi-model mean (Table 5) with uncertainties of about  $\pm 0.9$   $\text{GtC}$   
 18  $\text{yr}^{-1}$ . See the text for more details of each component and their uncertainties.



1

2 **Figure 4.** Components of the global carbon budget and their uncertainties as a function of time,  
 3 presented individually for **(a)** emissions from fossil fuels and industry ( $E_{FF}$ ), **(b)** emissions from  
 4 land-use change ( $E_{LUC}$ ), **(c)** the budget imbalance that is not accounted for by the other terms, **(d)**  
 5 growth rate in atmospheric  $CO_2$  concentration ( $G_{ATM}$ ), and **(e)** the land  $CO_2$  sink ( $S_{LAND}$ , positive  
 6 indicates a flux from the atmosphere to the land), **(f)** the ocean  $CO_2$  sink ( $S_{OCEAN}$ , positive indicates  
 7 a flux from the atmosphere to the ocean). All time series are in  $GtC\ yr^{-1}$  with the uncertainty  
 8 bounds representing  $\pm 1\sigma$  in shaded colour. Data sources are as in Fig. 3. The black dots in **(a)**  
 9 values for 2015 and 2016 that originate from a different data set to the remainder of the data (see  
 10 text). The dashed line in **(b)** identifies the pre-satellite period before the inclusion of peatland  
 11 burning.

12




1  
 2 **Figure 5.** CO<sub>2</sub> emissions from fossil fuels and industry for **(a)** the globe, including an uncertainty of  
 3 ± 5% (grey shading), the emissions extrapolated using BP energy statistics (black dots) and the  
 4 emissions projection for year 2017 based on GDP projection (red dot), **(b)** global emissions by fuel  
 5 type, including coal (salmon), oil (olive), gas (turquoise), and cement (purple), and excluding gas  
 6 flaring which is small (0.6% in 2013), **(c)** territorial (solid line) and consumption (dashed line)  
 7 emissions for the countries listed in Annex B of the Kyoto Protocol (salmon lines; mostly advanced  
 8 economies with emissions limitations) versus non-Annex B countries (green lines); also shown are  
 9 the emissions transfer from non-Annex B to Annex B countries (light blue line) **(d)** territorial CO<sub>2</sub>  
 10 emissions for the top three country emitters (USA - olive; China - salmon; India - purple) and for



1 the European Union (EU; turquoise for the 28 member states of the EU as of 2012), and **(e)** per-  
2 capita emissions for the top three country emitters and the EU (all colours as in panel **(d)**) and the  
3 world (black). In **(b-e)**, the dots show the data that were extrapolated from BP energy statistics  
4 for 2014 and 2015. All time series are in  $\text{GtC yr}^{-1}$  except the per-capita emissions **(e)**, which are in  
5 tonnes of carbon per person per year ( $\text{tC person}^{-1} \text{yr}^{-1}$ ). Territorial emissions are primarily from  
6 Boden et al. (2017) except national data for the USA and EU28 for 1990-2014, which are reported  
7 by the countries to the UNFCCC as detailed in the text; consumption-based emissions are updated  
8 from Peters et al. (2011a). See Sect. 2.1.1 for details of the calculations and data sources.

9




1  
2 **Figure 6.** CO<sub>2</sub> exchanges between the atmosphere and the terrestrial biosphere as used in the  
3 global carbon budget (black with  $\pm 1\sigma$  uncertainty in grey shading), for **(a)** CO<sub>2</sub> emissions from  
4 land-use change ( $E_{LUC}$ ), showing also individually the two bookkeeping models (two blue lines) and  
5 the DGVM model results (green) and their multi-model mean (olive). The dashed line identifies  
6 the pre-satellite period before the inclusion of peatland burning; **(b)** Land CO<sub>2</sub> sink ( $S_{LAND}$ ) with  
7 individual DGVMs (green); **(c)** Total land CO<sub>2</sub> fluxes (**b minus a**) with individual DGVMs (green) and  
8 their multi-model mean (olive), and atmospheric inversions (CAMS in purple, Jena CarboScope in  
9 violet, CTE in salmon; see details in Table 5). In **(c)** the inversions were corrected for the  
10 preindustrial land sink of CO<sub>2</sub> from river input, by removing a sink of  $0.45 \text{ GtC yr}^{-1}$  (Jacobson et al.,  
11 2007), but not for the anthropogenic contribution to river fluxes (see Sect. 2.7.2).

12



1




2

3 **Figure 7.** Comparison of the anthropogenic atmosphere-ocean CO<sub>2</sub> flux showing the budget values  
 4 of  $S_{\text{OCEAN}}$  (black; with  $\pm 1\sigma$  uncertainty in grey shading), individual ocean models (blue), and the two  
 5 ocean pCO<sub>2</sub>-based flux products (Rödenbeck et al. (2014) in salmon and Landschützer et al. (2015)  
 6 in purple; see Table 5). Both pCO<sub>2</sub>-based flux products were adjusted for the preindustrial ocean  
 7 source of CO<sub>2</sub> from river input to the ocean, which is not present in the ocean models, by adding a  
 8 sink of 0.45 GtC yr<sup>-1</sup> (Jacobson et al., 2007), to make them comparable to  $S_{\text{OCEAN}}$ . This adjustment  
 9 does not take into account the anthropogenic contribution to river fluxes (see Sect. 2.7.2).




1



2

3 **Figure 8.** CO<sub>2</sub> fluxes between the atmosphere and the surface ( $S_{\text{OCEAN}} + S_{\text{LAND}} - E_{\text{LUC}}$ ) by latitude  
4 bands for the **(a)** North (north of 30°N), **(b)** Tropics (30°S-30°N), and **(c)** South (south of 30°S).  
5 Estimates from the combination of the process models for the land and oceans are shown  
6 (turquoise) with  $\pm 1\sigma$  of the model ensemble (in grey). Results from the three atmospheric  
7 inversions are also shown (CAM5 in purple, Jena CarboScope in violet, CTE in salmon; references  
8 and version number in Table 5). Where available the uncertainty in the inversions are also shown.  
9 Positive values indicate a flux from the atmosphere to the land and/or ocean.

10



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13

**Figure 9.** Comparison of global carbon budget components released annually by GCP since 2006. CO<sub>2</sub> emissions from **(a)** fossil fuels and industry ( $E_{FF}$ ), and **(b)** land-use change ( $E_{LUC}$ ), as well as their partitioning among **(c)** the atmosphere ( $G_{ATM}$ ), **(d)** the land ( $S_{LAND}$ ), and **(e)** the ocean ( $S_{OCEAN}$ ). See legend for the corresponding years, and Table 3 for references. The budget year corresponds to the year when the budget was first released. All values are in GtC yr<sup>-1</sup>. Grey shading shows the uncertainty bounds representing  $\pm 1\sigma$  of the current global carbon budget.



1 **Table A1.** Funding supporting the production of the various components of the global carbon  
2 budget (see also acknowledgements).

<b>Funder and grant number (where relevant)</b>	<b>author initials</b>
Australia, Integrated Marine Observing System (IMOS)	BT
Australian National Environment Science Program (NESP)	JGC, VH
EC H2020 European Research Council (ERC) (QUINCY; grant no. 647204).	SZ
EC H2020 ERC Synergy grant (IMBALANCE-P; grant no. ERC-2013-SyG-610028)	DZ
EC H2020 project CRESCENDO (grant no. 641816)	PF, RS
EC FP7 project HELIX (grant no. 603864)	PF, RAB, SS
EU FP7 project LUC4C (grant no. 603542)	PF, MK, SS
French Institut National des Sciences de l'Univers (INSU) and Institut Paul Emile Victor (IPEV), Sorbonne Universités (UPMC, Univ Paris 06)	NM
German federal Ministry for Education and Research (BMBF)	GR, AK, SVH
German Federal Ministry of Transport and Digital Infrastructure (BMVI)	AK, SVH
German Research Foundation's Emmy Noether Programme (grant no. PO1751/1-1)	JEMSN, JP
IRD, RI Integrated Carbon Observation System (ICOS)	NL
Japan National Institute for Environmental Studies (NIES), Ministry of Environment (MOE)	SK, YN
NASA LCLUC programme (grant no. NASA NNX14AD94G)	AJ
Netherlands Organization for Scientific Research (NWO) Veni grant (016.Veni.171.095)	IvdLL
New Zealand National Institute of Water and Atmospheric Research (NIWA) Core Funding	KC
Norwegian Research Council, Norwegian Environmental Agency	IS
Norwegian Research Council (ICOS 245927)	BP, MB
Norwegian Research Council (grant no. 229771)	JS
South Africa Council for Scientific and Industrial Research, Department of Science and Technology (DST)	PMSM
RI Integrated Carbon Observation System (ICOS)	AW, GR, AK, SVH, IS, BP, MB
Swiss National Science Foundation (grant no.200020_172476)	SL
UK BEIS/Defra Met Office Hadley Centre Climate Programme (grant no. GA01101)	RAB
UK Natural Environment Research Council (SONATA: grant no. NE/P021417/1)	CLQ, OA
UK NERC, EU FP7, EU Horizon2020	AW
USA Department of Energy, Office of Science and BER prg. (grant no. DE-SC000 0016323)	ATJ
USA National Oceanographic and Atmospheric Administration (NOAA) Ocean Acidification Program (OAP) NA16NOS0120023	CWH
USA National Science Foundation (grant no. OPP 1543457)	DRM
USA National Science Foundation (grant no. AGS 12-43071)	AKJ
<b>Computing resources</b>	
Grand Équipement National de Calcul Intensif (allocation x2016016328), France	NV
Météo-France/DSI supercomputing centre	RS
Netherlands Organization for Scientific Research (NWO) (SH-312-14)	IvdL-L
Norwegian Metacenter for Computational Science (NOTUR, project nn2980k) and the Norwegian Storage Infrastructure (NorStore, project ns2980k)	JS





1 **Table A2** Attribution of fCO<sub>2</sub> measurements for the year 2016 included in SOCAT v5 (Bakker et al., 2016) to inform ocean pCO<sub>2</sub>-based flux  
2 products.  
3

Vessel	Regions	No. of samples	Principal investigators	Number of data sets
Allure of the Seas	North Atlantic, Tropical Atlantic	71744	Wanninkhof, R.; Pierrot, D.	36
Atlantic Cartier	North Atlantic	44302	Steinhoff, T.; Körtinger, A.; Becker, M.; Wallace, D.	12
Aurora Australis	Southern Ocean	43885	Tilbrook, B.	2
Benguela Stream	North Atlantic, Tropical Atlantic	137902	Schuster, U.; Watson, A.J.	21
Cap Blanche	North Pacific, Tropical Pacific	17913	Casca, C.; Alin, S.; Feely, R.; Herndon, J.	3
Cap San Lorenzo	North Atlantic, Tropical Atlantic	9126	Lefèvre, N.	3
Colibri	North Atlantic, Tropical Atlantic	27780	Lefèvre, N.	6
Equinox	North Atlantic, Tropical Atlantic	97106	Wanninkhof, R.; Pierrot, D.	35
F.G. Walton Smith	North Atlantic, Tropical Atlantic	43222	Millero, F.; Wanninkhof, R.	16
Finnmaid	North Atlantic	34303	Reider, G.; Glockzin, M.	3
G.O. Sars	Arctic, North Atlantic	109125	Skjelvan, I.	13
GAKOA (149W 60N)	North Pacific	488	Cross, J.; Mathis, J.; Monaco, N.; Musielewicz, S.; Maenner, S.; Osborne, J.	1
Gordon Gunter	North Atlantic, Tropical Atlantic	59310	Wanninkhof, R.; Pierrot, D.	13
Henry B. Bigelow	North Atlantic	61021	Wanninkhof, R.	13
Investigator	Southern Ocean, Tropical Pacific	108721	Tilbrook, B.	3
Laurence M. Gould	Southern Ocean	26150	Sweeney, C.; Takahashi, T.; Newberger, T.; Sutherland, S.C.; Munro, D.	5
Marion Dufresne	Southern Ocean	3214	Metz, N.; Lo Monaco, C.	1
New Century 2	North Atlantic, North Pacific, Tropical Pacific	25222	Nakaka, S.	15
Nuka Arctica	North Atlantic	47392	Becker, M.; Olsen, A.; Omar, A.; Johannessen, T.	12
Polarstern	Arctic, North Atlantic, Southern Ocean, Tropical Atlantic	164407	van Heuven, S.; Hoppema, M.	5



Roger Revelle	Indian Ocean, Southern Ocean, Tropical Pacific	93689	Wanninkhof, R.; Pierrot, D.	8
Ronald H. Brown	North Pacific, Tropical Pacific	52267	Wanninkhof, R.; Pierrot, D.	8
S.A. Aguilhas II	Southern Ocean	27851	Monteiro, P.M.S.; Joubert, W.R.	
Samiendo de Gamboa	North Atlantic, Southern Ocean, Tropical Atlantic	16122	Padin, X.A.	2
Savannah	North Atlantic	2803	Gai, W.-J.; Reimer, J.J.	1
SE Alaska (56N 134W)	North Pacific	271	Cross, J.; Mathis, J.; Monacchi, N.; Musielewicz, S.; Maenner, S.; Osborne, J.	1
Skogafoss	North Atlantic	22541	Wanninkhof, R.; Pierrot, D.	4
Tangaroa	Southern Ocean	118997	Currie, K.	7
Thomas G. Thompson	North Pacific, Tropical Pacific	14656	Alin, S.; Cosca, C.; Henson, J.; Feely, R.	1
Trans Future 5	North Pacific, Tropical Pacific, Southern Ocean	23087	Nakakka, S.; Nojiri, Y.	21
UNH Gulf Challenger	North Atlantic	2984	Hunt, C.W.	3

1

2