

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

A comparison of the chemical sinks of atmospheric organics in the gas and aqueous phase

S. A. Epstein and S. A. Nizkorodov

Department of Chemistry, University of California, Irvine, CA, USA

Received: 24 March 2012 – Accepted: 1 April 2012 – Published: 19 April 2012

Correspondence to: S. A. Nizkorodov (nizkorod@uci.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Photochemical reactions represent the main pathway for the removal of non-methane volatile organic compounds (VOCs) in the atmosphere. VOCs may react with hydroxyl radical (OH), the most important atmospheric oxidant, or they can be photolyzed by actinic radiation. In the presence of clouds and fog, VOCs may partition into the aqueous phase where they can undergo aqueous photolysis and/or reaction with dissolved OH. The significance of direct aqueous photolysis is largely uncertain due to the lack of published absorption cross sections and photolysis quantum yields. In light of this, we strive to identify atmospherically relevant VOCs where removal by aqueous photolysis may be a significant sink. The relative importance of different photochemical sinks is assessed by calculating the ratios of the removal rates inside air parcels containing cloud and fog droplets. This relative approach provides useful information in spite of the limited aqueous photolysis data. Results of this work should help guide researchers in identifying molecules that are the most likely to undergo aqueous OH oxidation and photolysis. We find that out of the 27 atmospherically relevant species investigated, the removal of glyceraldehyde and pyruvic acid by aqueous photolysis is potentially an important sink. We also determine the relative magnitudes of these four chemical sinks for the set of relevant organic compounds.

1 Introduction

Organic compounds play a key role in the chemistry of the atmosphere. Global emissions of non-methane volatile organic compounds (VOCs) total approximately $1300 \text{ Tg C yr}^{-1}$ (Atkinson and Arey, 2003; Goldstein and Galbally, 2007). It is estimated that $130\text{--}270 \text{ Tg C yr}^{-1}$ of emitted VOCs are removed by dry and wet deposition, while the remaining $1030\text{--}1170 \text{ Tg C yr}^{-1}$ are transformed by chemical reactions. Oxidation by hydroxyl radical is the most important atmospheric sink for many VOCs (Andreae and Crutzen, 1997). Gas phase photolysis can also be important for certain classes of

ACPD

12, 10015–10058, 2012

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



compounds including carbonyls (Moortgat, 2001), peroxides (Lee et al., 2000), and organic nitrates (Atkinson, 1990). In the presence of clouds and fog, VOCs may partition into the aqueous phase to various extents depending on their solubility. As in the gas phase, OH is the main oxidant for the transformation of aqueous organics in cloud and fog droplets (Ervens et al., 2011). Direct photolysis in the aqueous phase can also be an important sink depending on the chemical functionality of the species (Vione et al., 2006; Graedel and Goldberg, 1983).

Given the complexity of possible VOC photo-oxidation mechanisms and products, we strive to classify compounds by the rate that they are removed by gaseous OH oxidation, gaseous photolysis, aqueous OH oxidation, and direct aqueous photolysis. We develop a framework to compare these four rates of removal for a subset of compounds identified as likely contributors to the water-soluble fraction of atmospheric particles (Saxena and Hildemann, 1996). Figure 1 is a schematic of our analysis showing the four chemical sinks under investigation and the dimensionless parameters we derive to model the behavior of the selected VOCs. Direct liquid photolysis is parameterized into a photolysis rate constant that depends on the actinic flux of solar radiation, the quantum yield of photolysis, and the absorption cross section of the absorbing species. In the absence of aqueous absorption cross section and quantum yield data for most organics, we aim to identify systems where direct aqueous photolysis is potentially a significant sink. While photolysis of certain organic compounds may generate OH radical in the gaseous or aqueous phase, we investigate direct photolysis and OH oxidation independently. With published OH rate constants for oxidation in the gas and aerosol phase, along with absorption cross section and quantum yield data for photolysis in the gas phase, we can estimate the significance of these four chemical sinks to better understand the chemical fate of water soluble atmospheric organics. We stress that this is not a modeling study – our goal is to help guide researchers in identifying molecules that are the most likely to undergo aqueous OH oxidation and photolysis.

We ignore several other VOC chemical sinks to simplify our analysis and focus on the competition between oxidation by OH and photolysis. Alkenes are susceptible to

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

oxidation by ozone (Bailey, 1958) and NO₃ (Atkinson, 1991) but we are not investigating any olefinic compounds. On a global scale, organics may react with Cl atoms, however this is thought to be a minor VOC loss process (Atkinson and Arey, 2003). For specific compounds under certain conditions, aqueous reactions with dissolved sulfur dioxide in cloud and fog droplets may be significant (Finlayson-Pitts and Pitts, 2000). In addition, inorganic ammonium ions can catalyze reactions of carbonyl compounds in the liquid phase (Noziere et al., 2008). Biological activity may also lead to the destruction of dissolved organics in cloud droplets. Microbes can drive the oxidation of carbonaceous compounds during periods with low dissolved OH concentrations in cloud droplets (Vaitilingom et al., 2010; Vaitilingom et al., 2011). We also neglect potential photosensitization processes from contaminants dissolved in atmospheric waters (Zepp et al., 1981).

2 Methods

Out of the oxygenated organic compounds identified by Saxena and Hildemann, (Saxena and Hildemann, 1996) we choose a subset of 27 compounds for which there is the required experimental data published in the literature. These selected compounds contain a range of functionalities: monocarboxylic acids, dicarboxylic acids, aldehydes, ketones, keto-carboxylic acids, ethers, and peroxides. While a similar analysis for a series of nitrogen-containing organic compounds is beyond the scope of this manuscript, we also investigate methyl nitrate, the simplest organic nitrate. With the exception of tartaric acid and malic acid, all compounds are either intermediate-volatility organic compounds (IVOCs) or volatile organic compounds (VOCs); the majority of their mass partitions in the vapor phase as opposed to the condensed phase. Tartaric acid and malic acid are low-volatility organic compounds (LVOCs, $C^* \approx 1.6 \times 10^{-2} \mu\text{g m}^{-3}$ and $C^* \approx 2.4 \mu\text{g m}^{-3}$, respectively (Pankow and Asher, 2008)) indicating that the majority of their mass will partition to the condensed phase in almost all atmospheric conditions (Donahue et al., 2009). The oxygen to carbon ratio of the selected compounds ranges

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from 0 to 2. A full list of the compounds used and their corresponding references is presented in the Supplement. All of the IVOCs and VOCs are delivered to cloud and fog droplets by gaseous diffusion. However, tartaric acid and malic acid, as components of particulate matter, may also be delivered to the aqueous phase by particle scavenging, known to be very efficient in clouds (Limbeck and Puxbaum, 2000). For each of the compounds studied, we assume that the partitioning between the gaseous and aqueous phase is in a local state of equilibrium. Because of the variability of transport processes and competing reactions, equilibrium may not be completely established in all cases (Winiwarter et al., 1994; Audiffren et al., 1998). For example, complete equilibrium is unlikely when water droplets of different sizes and pHs are present or in the case of an air mass with a variable liquid water content. In polluted environments, organics may enter cloud or fog droplets through the dissolution of organic aerosol particles potentially resulting in aqueous concentrations that exceed equilibrium values. However, for atmospheric gases at typical concentrations, Henry's law reasonably approximates equilibrium with cloud or fog droplets (Seinfeld and Pandis, 1998), and our equilibrium-based calculations should therefore provide a good first-order estimate for the magnitude of aqueous phase processes. In the presence of high ion concentrations, Henry's law constants can be modified with Setchenow coefficients (De Bruyn et al., 1995; Kolb et al., 1997). However, in order to streamline the analysis and maintain its generality, we assume that each of the compounds studied do not interact with each other or other inorganic compounds typically found in cloud or fog droplets.

2.1 Comparison of aqueous and gaseous photolysis

For atmospheric organics, the implications of direct aqueous photolysis (direct UV-irradiation, Bateman et al., 2011; Schwarzenbach et al., 2005) are elusive due to the lack of literature data. However, the direct gaseous photolysis of atmospheric organics is well-studied (Atkinson et al., 2006; Sander et al., 2011). We strive to exploit the likely similarities in these processes in order to constrain the magnitude of aqueous phase photolysis by comparison with gas phase photolysis. The photolysis rate constant, J ,

is calculated with the following integral:

$$J = \int F_A(\lambda) \cdot \Phi(\lambda) \cdot \sigma(\lambda) \cdot d\lambda \quad (1)$$

where F_A is the actinic flux, Φ is the quantum yield of photolysis, and σ is the absorption cross section, all functions of wavelength, λ . In the absence of aqueous quantum yields and/or absorption cross sections, we can constrain the ratio of the gas phase and aqueous phase rate constants for a particular compound using reasonable assumptions on how its photochemistry is affected by dissolution in water. The actinic flux can be enhanced slightly inside a cloud droplet compared to the interstitial space (Finlayson-Pitts and Pitts, 2000). This enhancement has been calculated to be up to a factor of 1.6 by Madronich (Madronich, 1987) or possibly more than a factor of two (Ruggaber et al., 1997). The solvent has the potential to change the absorption cross sections upon dissolution, known as solvatochromism (Marini et al., 2010), however, this effect is likely insignificant for the set of compounds that we are investigating. Xu et al. (1993) finds that the absorption cross section shifts by about 10 nm to the blue and the intensity increases by 15–35 % for a series of small ketones. For hydrogen peroxide (Graedel and Weschler, 1981) and methyl peroxide (Epstein et al., 2012), the absorption cross sections of the gas and aqueous phases are similar. The magnitude of the change in quantum yield is far less certain. For direct aqueous photolysis, the quantum yield must be between 0 and the corresponding gas-phase value because energy dissipation due to collisions with the solvent molecules (cage effect) usually depresses the quantum yield in solution. Very few experimental measurements of aqueous quantum yields exist, making attempts to constrain the change in quantum yield upon dissolution difficult. To obtain an upper estimate of aqueous phase photolysis, we will assume that the quantum yield in solution and in the gas-phase is the same.

To constrain the ratio of the rate of removal due to gaseous and aqueous photolysis we consider a system with an air parcel containing liquid water droplets and a pollutant x , which is in equilibrium between the gaseous and aqueous phase. The parcel has a

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



liquid water content (LWC) defined in units of mass of liquid water per volume of air. The ratio of the rate of gaseous photolysis and aqueous photolysis for species x is defined by the parameter Z :

$$Z = \frac{\frac{dn_{x+h\nu}^{\text{gas}}}{dt}}{\frac{dn_{x+h\nu}^{\text{aq}}}{dt}} = (R \cdot T \cdot \text{LWC}_v \cdot k_{hx})^{-1} \left(\frac{J_{x+h\nu}^{\text{gas}}}{J_{x+h\nu}^{\text{aq}}} \right) \quad (2)$$

5 where $n_{x+h\nu}$ is the moles of species x undergoing photolysis in the gas or aqueous phase in a given volume of air, t represents time, R is the gas constant, T is temperature, LWC_v is the volume based liquid water content in units of volume of liquid water per volume of air, k_{Hx} is the effective Henry's law constant for species x in units of molar concentration per pressure, J^{gas} and J^{aq} are the photolysis rate constants of species
 10 x in the gaseous and aqueous phase, respectively. The ratio of the rates does not depend on the concentration of species x because we assume that the gaseous and aqueous forms are in equilibrium. Derivations of all significant equations are presented in Appendix A.

Equation (2) is a model for systems where the species of interest does not hydrate or
 15 dissociate in the aqueous phase. However, photolabile carbonyls hydrate in the aqueous phase to form the corresponding gem-diol (Bell and McDougall, 1960). Using a generic mono-aldehyde as an example, we can write:



where K_{hyd} is the hydration equilibrium constant. We define the variable α_N as the
 20 fraction of the species that is present in form N upon dissolution. For a monocarbonyl, there may be two forms in solution: the free molecule and the hydrated molecule. For a dicarbonyl, there may be three forms in solution: the free molecule, the partially hydrated molecule, and the fully hydrated molecule. For a mono-carbonyl:

$$\alpha_{\text{free}} = (K_{\text{hyd}} + 1)^{-1} \quad \text{and} \quad \alpha_{\text{hyd}} = ((K_{\text{hyd}})^{-1} + 1)^{-1} \quad (4)$$

10021

Comparison of the chemical sinks

S. A. Epstein and
 S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where α_{free} is the fraction of the free molecule in solution and α_{hyd} is the fraction of the hydrated molecule in solution. Aldehydes hydrate to a larger extent than ketones. For most ketones, only a small fraction of molecules exist in the gem-diol form (Bell and Gold, 1966). This has significant implications for photolysis as gem-diols do not absorb actinic UV radiation (Xu et al., 1993), and therefore are not photolabile in the atmosphere. We make a simplifying assumption that if we normalize the photolysis rate constant of the free form of species x to unity, the photolysis rate constant of the fully hydrated form will be null, and in the special case of a dicarbonyl, the normalized photolysis rate constant of the partially hydrated form will be half of the free form because one of the photoactive carbonyls has been converted to non-photoactive gem-diol form. We define this normalized photolysis rate constant as β_N :

$$\beta_{\text{free}} = 1, \beta_{\text{hyd}} = 0, \text{ in the case of a dicarbonyl } \beta_{\text{part.hyd}} = \frac{1}{2} \quad (5)$$

Therefore, in the case of a hydrating carbonyl, x , we can penalize the aqueous rate constant by the following factor, γ_x :

$$\gamma_x = \sum_N \alpha_N \cdot \beta_N \quad (6)$$

where $\gamma_x \leq 1$. Using this factor in Eq. (2) allows us to continue using the effective Henry's law constant of species x despite the fact that x may hydrate into multiple forms in solution:

$$Z = \frac{\frac{dn_{x+hv}^{\text{gas}}}{dt}}{\frac{dn_{x+hv}^{\text{aq}}}{dt}} = (R \cdot T \cdot \text{LWC}_v \cdot k_{\text{H}x})^{-1} \left(\frac{j_{x+hv}^{\text{gas}}}{\gamma_x \cdot j_{x+hv}^{\text{aq}}} \right) \quad (7)$$

Because γ_x is always less than unity, the maximum effective photolysis rate is reduced in the aqueous phase for all carbonyls.

Finally, Henry's Law constants are strongly dependent on temperature. We perform calculations of Z at 298 K where most constants are measured to ensure a straightforward analysis. However, in Appendix B, we describe a way of correcting these equations for the temperature effects.

5 2.2 Comparison of aqueous and gaseous OH oxidation

The majority of atmospheric organics are vulnerable to OH oxidation in both the gaseous and aqueous phase. We aim to elucidate the relative significance of each of these processes for our selected set of compounds by calculating the ratio of the rate of OH oxidation in the gas and aqueous phase. The most significant difference between this analysis and the previous discussion is that the rate constants due to OH oxidation in the aqueous (Ervens et al., 2003b; Buxton et al., 1988; Monod et al., 2005) and gaseous phases (Sander et al., 2011; Atkinson et al., 2006) have been experimentally determined for many atmospheric organics. Even if gas phase OH oxidation rate constants have not been measured, structure activity relationships are well developed, and provide accurate rate constant estimates (Atkinson, 1987, 1988; Kwok and Atkinson, 1995). We define W as the ratio of the rate of gas phase OH oxidation of x and the rate of aqueous phase OH oxidation of x . For non-ionizable species:

$$W = \frac{\frac{dn_{x+\text{OH}}^{\text{gas}}}{dt}}{\frac{dn_{x+\text{OH}}^{\text{aq}}}{dt}} = (\text{LWC}_V \cdot k_H \cdot R \cdot T)^{-1} \left(\frac{k_{x+\text{OH}}^{\text{gas}}}{k_{x+\text{OH}}^{\text{aq}}} \right) \left(\frac{C_{\text{OH}}^{\text{gas}}}{C_{\text{OH}}^{\text{aq}}} \right) \quad (8)$$

Where $n_{x+\text{OH}}$ represents the moles of x undergoing oxidation by OH in the gas or aqueous phase, $k_{x+\text{OH}}$ is the rate constant for x oxidation by OH in the aqueous or gas phase, $C_{\text{OH}}^{\text{gas}}$ is the gas phase concentration of OH (moles of OH per volume of gas), and $C_{\text{OH}}^{\text{aq}}$ is the aqueous phase concentration of OH (moles of OH per volume of water). When using this equation, the units of $k_{x+\text{OH}}$ and C_{OH} must be consistent so that they cancel out upon division.

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Equation (8) can be used for species that hydrate because hydroxyl radical rate constants are typically tabulated for a solution containing all aqueous forms of the parent molecule (Ervens et al., 2003b; Monod et al., 2005). However, for species that ionize, we must use a modified version of Eq. (8) to account for the dissimilar aqueous OH rate constants of the ionized and native form. With measured pK_a values, we can calculate the fraction of the acid HA that is present in the ionized form (f_{A^-}) and the native form (f_{HA}) at a given solution pH.

$$f_{A^-} = \left(1 + 10^{pK_a - pH}\right)^{-1} \quad f_{HA} = \left(1 + 10^{pH - pK_a}\right)^{-1} \quad (9)$$

For a given pH and with OH rate constants for reaction with the acid, k_{HA+OH} , and its anion, k_{A^-+OH} , we can write W as:

$$W = \frac{\frac{dn_{HA+OH}^{gas}}{dt}}{\frac{dn_{HA+A^-+OH}^{aq}}{dt}} = (R \cdot T \cdot LWC_v \cdot k_{Hx})^{-1} \left(\frac{k_{HA+OH}^{gas}}{f_{HA} \cdot k_{HA+OH}^{aq} + f_{A^-} \cdot k_{A^-+OH}^{aq}} \right) \left(\frac{C_{OH}^{gas}}{C_{OH}^{aq}} \right) \quad (10)$$

In this case, both the Henry's law constant and the rate constants are temperature dependent, but we perform calculations at 298 K where most experimental data is measured to keep our analysis as straightforward as possible. Appendix B describes a way for incorporating temperature effects in this equation.

2.3 Comparison of gaseous OH oxidation and photolysis

The comparison of gas phase OH oxidation and photolysis is also based on the ratio of the rates of each of these processes to cancel out the dependence on species concentration. We define Y as the ratio of the rate of gas phase OH oxidation of species x and the rate of gas phase photolysis of species x . At a constant temperature and

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



pressure, Y is a function of the solar zenith angle, SZA:

$$Y(\text{SZA}) = \frac{\frac{dn_{x+\text{OH}}^{\text{gas}}}{dt}}{\frac{dn_{x+h\nu}^{\text{gas}}}{dt}} = \frac{k_{x+\text{OH}}^{\text{gas}} \cdot C_{\text{OH}}(\text{SZA})}{J_{x+h\nu}^{\text{gas}}(\text{SZA})} \quad (11)$$

where $C_{\text{OH}}(\text{SZA})$ is the gas phase OH concentration as a function of the solar zenith angle in units that correspond to the units of $k_{x+\text{OH}}$. This concentration is described by the empirical power-law function relating the OH concentration to the photolysis rate of $\text{O}_3 \rightarrow \text{O}(^1\text{D})$ described in (Rohrer and Berresheim, 2006).

For species with measured absorption cross sections as a function of temperature and quantum yields as a function of temperature and/or pressure, we can go one step further and examine how Y varies with atmospheric height and solar zenith angle.

$$Y(\text{SZA}, z) = \frac{\frac{dn_{x+\text{OH}}^{\text{gas}}}{dt}}{\frac{dn_{x+h\nu}^{\text{gas}}}{dt}} = \frac{k_{x+\text{OH}}^{\text{gas}}(T) \cdot C_{\text{OH}}(\text{SZA}, T)}{J_{x+h\nu}^{\text{gas}}(\text{SZA}, z, T)} \quad (12)$$

where the change in temperature at a given height in the atmosphere, z , is given by:

$$-\frac{dT}{dz} = \Gamma \quad (13)$$

The parameter, Γ represents the lapse rate, for which we use the average lapse rate in the troposphere, 6.5 K km^{-1} (Wallace and Hobbs, 2006). The actinic flux as a function of solar zenith angle and wavelength is predicted with the Tropospheric Ultraviolet and Visible (TUV) Radiation Model (Madronich).

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results

3.1 Comparison of aqueous and gaseous photolysis

Our comparison of aqueous and gaseous photolysis with Eqs. (2) and (7) provide an upper estimate to the extent of in-cloud photolysis for the following reasons. We perform this analysis using an air parcel at 298 K with a liquid water content of 3 g m^{-3} , typically the largest value found in the atmosphere (Seinfeld and Pandis, 1998). Aqueous processes are more significant in wetter air parcels as a larger fraction of each organic resides in the aqueous phase. The assumption of Henry's law equilibrium is also an upper bound of the aqueous phase concentration under most conditions (Winiwarter et al., 1994; Audiffren et al., 1998). In addition, aqueous quantum yields may be significantly depressed by solvent effects, thus slowing down the rate of aqueous photolysis relative to gaseous photolysis.

Figure 2 shows the results of this analysis for a series of atmospherically relevant, non-hydrating species, specifically peroxides and methyl nitrate. The ratio of gas and aqueous rate constants are shown on the axis of abscissas and the effective Henry's law constants are shown on the axis of ordinates. Colors and diagonal isopleths show the magnitude of Z . Molecules are labeled in order of increasing Z . References for the Henry's law parameters are presented in the Supplement. In cases where measurements exist from multiple researchers and an agreed value was not established, we used the mean of the experimental values. The x-axis intentionally spans a large range to provide an indication of the sensitivity of this analysis. Actual values should lie somewhere to the right ($J_{\text{aqueous}} < J_{\text{gas}}$) of our upper bound estimations.

Of the four molecules included in Fig. 2, only hydrogen peroxide has the potential for an appreciable aqueous photolysis removal rate relative to gas phase photolysis. With actual experimental data on the absorption cross sections and quantum yields of gaseous and aqueous hydrogen peroxide from (Sander et al., 2011; Chu and Anastasio, 2005), we can pinpoint the location of this molecule on the x-axis more precisely. The two inward arrows adjacent to the hydrogen peroxide estimation point (1) on Fig. 2

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

detailing the experimentally measured range in $J_{\text{gas}}/J_{\text{aqueous}}$ from a solar zenith angle of 0–90° shows that the actual value is moderately below our upper-bound estimate. Gelencser and Varga (Gelencser and Varga, 2005) estimate that multi-phase reactions become important relative to gas-phase photo-oxidation processes when the effective Henry's law constant is greater than 10^3 M atm^{-1} . We find a similar threshold value in our analysis of the photolysis of non-hydrating species.

The series of atmospherically relevant, hydrating species is presented in Fig. 3. When experimental measurements of the Henry's Law constants were unavailable (malonic acid, succinic acid, malic acid, lactic acid, tartaric acid, and glyceraldehyde), group contribution estimations were used (Saxena and Hildemann, 1996). Aqueous photolytic removal is potentially important in several of these species due to their large effective Henry's law parameters. Upper-bound estimates are not all centered about the $J_{\text{gas}}/J_{\text{aqueous}} = 1$ line due to hydration into non-photolabile gem-diols in the aqueous phase. For example, at low concentrations, formaldehyde is almost exclusively in diol form (Dong and Dasgupta, 1986; Le Botlan et al., 1983). Therefore, formaldehyde (6) is represented by the arrow pointing towards $J_{\text{gas}}/J_{\text{aqueous}} = \infty$. Aqueous photolytic removal of glyceraldehyde ($Z \leq 1.7 \times 10^{-5}$) is potentially very significant. Aqueous removal of pyruvic acid ($Z \leq 4.6 \times 10^{-2}$) is likely significant and aqueous removal of gluteraldehyde ($Z \leq 1$), methyl glyoxal ($Z \leq 2.6$), glyoxal ($Z \leq 3.3$), glycolaldehyde ($Z \leq 3.7$), and glyoxylic acid ($Z \leq 4.8$) has the potential to be somewhat important. Ethers and carboxylic acids were not analyzed because they are not photolabile under typical tropospheric conditions (Chebbi and Carlier, 1996; Atkinson and Arey, 2003; Finlayson-Pitts and Pitts, 2000).

3.2 Comparison of aqueous and gaseous OH oxidation

The ratio of the hydroxyl radical concentration in the gas and aqueous phase is a key uncertainty in the evaluation of W (the ratio of the rate of gas phase OH oxidation to aqueous phase OH oxidation). We use this OH ratio as one of our free parameters to

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

allow for a sensitivity analysis, but present an educated guess to better constrain our results. For the gaseous OH concentration, we use the upper end of the measured and theoretical estimates suggested in (Seinfeld and Pandis, 1998), $C_{\text{OH}}^{\text{daytime}} = 1 \times 10^7$ molecules cm^{-3} and $C_{\text{OH}}^{\text{nighttime}} = 2 \times 10^5$ molecules cm^{-3} . For the aqueous OH concentration in a cloud or fog droplet, we use the recommended values in the review by (Ervens et al., 2011) that were determined with theoretical models (Ervens and Volkamer, 2010; Ervens et al., 2003a): $C_{\text{OH}}^{\text{daytime}} = 10^{-13}$ M and $C_{\text{OH}}^{\text{nighttime}} = 10^{-14}$ M. The OH concentrations that we use are similar to the values assumed in (Gelencser and Varga, 2005).

Figure 4 shows a graphical representation of Eq. (8) for a series of peroxides and ethers in an air parcel with a LWC of 3 g m^{-3} at 298 K. The ratio of the gas and aqueous phase OH concentrations (moles of OH per volume of air/moles of OH per volume of liquid), which we will refer to as the “OH ratio”, are shown on the x-axis. The product of the ratio of the gas and aqueous OH rate constants and the inverse Henry’s law constant (atm M^{-1}) is represented on the y-axis. Colors along with the diagonal isopleths indicate the magnitude of W . Each species is represented by a labeled horizontal line bound by our daytime and nighttime estimates in the OH ratio. Species are labeled in order of increasing W . A line showing the OH ratio in Henry’s Law equilibrium (rarely realized in practice because of the high OH reactivity) is also presented for perspective. All of the rate constants are referenced in the Supplement. Based on this analysis, we find that OH oxidation of hydrogen peroxide is important in the aqueous phase. Aqueous OH oxidation of ethyl peroxide and methyl peroxide may be important depending on the OH ratio. Gas OH oxidation strongly dominates the removal of methyl tert-butyl ether, dimethyl ether, and ethyl ether.

Figure 5 is a continuation of this evaluation for a series of carbonyls under the same conditions. Aqueous oxidation dominates the removal of glyoxal compared to gas phase oxidation under the conditions of high LWC. Also, aqueous reaction with OH is likely a significant sink in the removal of methyl glyoxal and formaldehyde compared to gas phase oxidation. Removal of acetone by aqueous oxidation may be more

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

important depending on the OH ratio. Gaseous oxidation dominates the removal of 2-butanone, acetaldehyde, butanal, and propanal compared to aqueous oxidation by OH.

Figure 6 shows a similar analysis for a series of acids using Eq. (10) without color contours for simplicity. Molecules are represented by labeled rectangles instead of horizontal lines due to the dependence of aqueous oxidation rates on pH. Labeled rectangles are horizontally bound by estimates of the OH ratio in the daytime and nighttime. Rectangles are vertically bound by typical cloud or fog droplet pH from 2–6 (Finlayson-Pitts and Pitts, 2000). Species are labeled according to increasing W at pH 6. Aqueous removal by OH is an important sink compared to gaseous removal for all of the acids investigated here because of their large water solubilities. We expect aqueous removal to dominate for all acids except acetic acid, 3-methyl butanoic acid, and 2,2-dimethyl 1-propanoic acid.

The vertical coordinate on Figs. 4–6 can be simplified because gas and aqueous OH rate constants are related for an individual species. Most organic rate constants lie within a factor of 3 of a line relating $\log(k_{\text{gas}}^{\text{OH}})$ and $\log(k_{\text{aqueous}}^{\text{OH}})$ (Haag and Yao, 1992). In light of this relationship, the dimensionless number, DN , introduced in this work, is a reasonable predictor for the magnitude of W for our range of assumed OH ratio where:

$$DN = (\text{LWC}_v \cdot R \cdot T \cdot k_H)^{-1} \quad (14)$$

Figure 7 shows the predictive power of DN for the series of compounds investigated during daytime OH concentrations in an air parcel with a $\text{LWC} = 3 \text{ g m}^{-3}$ at 298 K. When $DN \gg 10$, gaseous OH oxidation is more important than gas phase OH oxidation. Aqueous phase oxidation is more important when $DN \ll 10$. Both reactions are potentially significant when DN is on the order of 10. This relationship holds with air parcels containing different liquid water contents or parcels with nighttime OH concentrations; a multi-panel plot showing W as a function of DN under different conditions is presented in the Supplement.

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

3.3 Comparison of gaseous OH oxidation and photolysis

For a given species, the ratio of the rate of gas phase OH oxidation to the rate of gaseous photolysis, Y , is dependent on the solar zenith angle, temperature, and pressure (see Eq. 12). We first investigate a series of 12 compounds with published quantum yield and absorption cross section measurements at 298 K and 1 atm using Eq. (11). The references for these values are presented in the Supplement. Figure 8 shows how Y varies with SZA for each of the selected compounds at sea level and 298 K. The width of the red swathe is due to the uncertainty in the empirical function from (Rohrer and Berresheim, 2006) that correlates OH radical concentration with the photolysis rate of O_3 . The actinic flux at each wavelength as a function of SZA was predicted with the TUV Radiation Model (Madronich) with a surface albedo of 0.154, the globally averaged value from (Hummel and Reck, 1979), and an ozone column of 300 Dobson units. In general, the selected molecules exhibit suppressed rates of photolysis at SZA > 80 degrees. For acetaldehyde, glycolaldehyde, acetone, 2-butanone, butanal, methyl peroxide, and propanal, gas phase oxidation is more important than photolysis at all SZA. Removal of methyl nitrate by photolysis is more significant than removal by OH oxidation. For formaldehyde, glyoxal, methyl glyoxal, and hydrogen peroxide, the relative importance of each of these removal processes is dependent on the SZA and the local OH concentration.

We also investigated this ratio as a function of SZA and altitude assuming a standard tropospheric lapse rate of 6.5 K km^{-1} (Wallace and Hobbs, 2006) and a scale height of 7.5 km for four compounds with published quantum yields and absorption cross section measurements as a function of temperature and pressure and OH rate constants as a function of temperature. See the Supplement for these references. In this case, the actinic flux was predicted as a function of wavelength, SZA, and altitude with a surface albedo of 0.154 and an ozone column of 300 Dobson units. Figure 9 shows a complicated dependence of Y on altitude and SZA for each of the four species investigated.

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This analysis at the bounds of the OH concentration uncertainty is presented in the Supplement.

4 Atmospheric implications

With knowledge of the three ratios, W , Z , and Y , which characterize the significance of each of the four chemical sinks investigated, we can calculate the relative contribution of each of the sinks for a specific compound. The results of this analysis are presented in Fig. 10 for compounds with a full set of requisite data in an air mass with a $LWC = 0.05 \text{ g m}^{-3}$ (characteristic of light clouds and fog) and a $LWC = 3 \text{ g m}^{-3}$ (characteristic of very dense clouds) at 298 K with $SZA = 20^\circ$. The length of each bar shows the best-guess estimation of the chemical sink corresponding to its color. In the case of aqueous photolysis, the cyan bars show an upper bound estimate of its contribution to the removal of each specific species at 298 K. One of the major conclusions of this analysis is that even in conditions where aqueous processes are most favorable, $LWC = 3 \text{ g m}^{-3}$, removal by aqueous photolysis remains a minor sink of the species investigated in Fig. 10. The only compounds with an appreciable aqueous photolysis contribution are hydrogen peroxide and glyoxal. (However, aqueous photolysis may be important for more structurally complicated compounds that do not appear in Fig. 10. Published OH rate constants, gaseous absorption cross sections, and quantum yields are available only for relatively simple compounds.) At lower temperatures, the increased solubility of cloud and fog droplets slightly enhances this contribution, but even in an air mass with a $LWC = 3 \text{ g m}^{-3}$ at 273 K, liquid photolysis of hydrogen peroxide accounts for less than 20 % of its removal. However, photolysis of aqueous H_2O_2 is a significant source of aqueous OH; its rate of production is on the order of the rate of droplet scavenging from the gas phase (Zellner et al., 1990). A complete analysis detailing the magnitude of each of the chemical sinks at 273 K for species with published Henry's Law temperature dependences is presented in Appendix B.

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Our analysis of the ratio of gas phase photolysis and aqueous phase photolysis can help guide the selection of compounds for future experiments. Aqueous photolysis absorption cross sections and quantum yields are not well established in the literature, but may be important parameters for specific compounds. Glyceraldehyde and pyruvic acid were not included in Fig. 10 due to the lack of published gaseous absorption cross sections and quantum yields. However, based on the results of Fig. 3, we expect aqueous photolysis to be a significant sink for these compounds. For compounds that were not analyzed in this manuscript, the potential significance of aqueous photolysis can be determined with knowledge of the Henry's law constant and the hydration equilibrium constants if hydration occurs in the aqueous phase.

For most acids in wet clouds, aqueous phase OH oxidation is more important than gas phase OH oxidation. For atmospherically relevant compounds without published aqueous rate constants, experiments designed to measure these values may prove to be fruitful. Since the ratio of the gas and aqueous phase OH removal rates are a strong function of LWC and the Henry's Law constant, one can determine additional species for which experimental measurements are warranted.

We have chosen to treat aqueous phase photolysis and oxidation separately, but in certain cases, photolytic cleavage may liberate OH and other free radicals that will react with the organic precursor. This will effectively increase the quantum yield of photolysis for the removal of the precursor beyond what would be measured if the liberated OH was being quenched by a competing reaction. However, we expect this potential enhancement to be small (at most a factor of 2 or 3) as in the case of methyl peroxide (Epstein et al., 2012).

As indicated in Fig. 9, the ratio of gas phase oxidation to gas phase photolysis is a complicated function of altitude and SZA. Gas phase absorption cross section and quantum yield measurements at 298 K and 1 atm may not provide enough information to describe the atmospheric behavior of certain organics especially at higher SZA. For example, photolysis of methyl glyoxal, and hydrogen peroxide relative to oxidation is enhanced at high latitudes ($SZA > 60^\circ$) and altitudes compared to the ratio at 1 atm.

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In the case of methyl glyoxal at solar zenith angles $<60^\circ$, photolysis relative to oxidation is suppressed at higher altitudes. In light of these complicated dependencies, we recommend caution when extrapolating room temperature and 1 atm measurements of photolysis quantum yields and absorption cross sections.

We have assumed that the organic molecules in our analysis do not interact with each other. While this assumption is needed to constrain the complexity of interacting molecules and draw conclusions, it may not be appropriate in all conditions. For example, in a polluted environment, organics may compete for aqueous phase OH radicals in a cloud or fog droplet. A more complicated analysis is required to capture these effects.

Appendix A

Equation derivations

A1 Derivation of Z

We start with the rate laws for photolysis in the gas and aqueous phase:

$$\frac{dC_{x+h\nu}^{\text{gas}}}{dt} = -J_{x+h\nu}^{\text{gas}} C_x^{\text{gas}} \quad (\text{A1})$$

$$\frac{dC_{x+h\nu}^{\text{aq}}}{dt} = -J_{x+h\nu}^{\text{aq}} C_x^{\text{aq}} \quad (\text{A2})$$

where C_x^{gas} is the gas phase concentration of species x and C_x^{aq} is the aqueous phase concentration of x . We transform concentration units into a molar basis with the volume

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the air parcel (V_{air}) and the volume of water inside the air parcel ($V_{\text{H}_2\text{O}}$).

$$\frac{dn_{x+h\nu}^{\text{gas}}}{dt} = V_{\text{air}} \cdot \frac{dC_{x+h\nu}^{\text{gas}}}{dt} \quad (\text{A3})$$

$$\frac{dn_{x+h\nu}^{\text{aq}}}{dt} = V_{\text{H}_2\text{O}} \cdot \frac{dC_{x+h\nu}^{\text{aq}}}{dt} \quad (\text{A4})$$

We define Z as the ratio between these two molar rates:

$$Z \equiv \frac{\frac{dn_{x+h\nu}^{\text{gas}}}{dt}}{\frac{dn_{x+h\nu}^{\text{aq}}}{dt}} \quad (\text{A5})$$

The ratio of air and water volumes can be simplified with a volumetric cloud liquid water content (LWC_v , volume of H_2O /volume of air parcel):

$$Z = \frac{V_{\text{air}} \cdot \frac{dC_{x+h\nu}^{\text{gas}}}{dt}}{V_{\text{H}_2\text{O}} \cdot \frac{dC_{x+h\nu}^{\text{aq}}}{dt}} = \frac{1}{\text{LWC}_v} \cdot \frac{J_{x+h\nu}^{\text{gas}} C_x^{\text{gas}}}{J_{x+h\nu}^{\text{aq}} C_x^{\text{aq}}} \quad (\text{A6})$$

where the volumetric liquid water volume is defined as:

$$\text{LWC}_v = \frac{\text{LWC}}{\rho_{\text{H}_2\text{O}}} \quad (\text{A7})$$

The density of water is represented by $\rho_{\text{H}_2\text{O}}$ and LWC is the liquid water content in units of mass of water per volume of air parcel. We now introduce the Henry's law

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



constant to describe the ratio of the aqueous phase concentration of x (C_x^{aq}) and the partial pressure of x (p_x^{gas}).

$$k_H = \frac{C_x^{\text{aq}}}{p_x^{\text{gas}}} \quad (\text{A8})$$

5 Converting pressure to concentration units with the ideal gas constant and temperature yields the equilibrium ratio of gas and aqueous phase concentration of x :

$$\frac{C_x^{\text{gas}}}{C_x^{\text{aq}}} = \frac{1}{k_H RT} \quad (\text{A9})$$

Our final working equation to describe the ratio of the rate of gas phase photolysis to the rate of aqueous phase photolysis is:

$$Z = \frac{1}{\text{LWC}_v \cdot k_H \cdot R \cdot T} \cdot \left(\frac{J_{x+h\nu}^{\text{gas}}}{J_{x+h\nu}^{\text{aq}}} \right) \quad (\text{A10})$$

10 A2 Derivation of W

We start with the rate laws for oxidation of species x in the gas and aqueous phase by OH:

$$\frac{dC_{x+\text{OH}}^{\text{gas}}}{dt} = -k_{x+\text{OH}}^{\text{gas}} C_x^{\text{gas}} C_{\text{OH}}^{\text{gas}} \quad (\text{A11})$$

$$15 \frac{dC_{x+\text{OH}}^{\text{aq}}}{dt} = -k_{x+\text{OH}}^{\text{aq}} C_x^{\text{aq}} C_{\text{OH}}^{\text{aq}} \quad (\text{A12})$$

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Converting to molar units:

$$\frac{dn_{x+OH}^{\text{gas}}}{dt} = V_{\text{air}} \cdot \frac{dC_{x+OH}^{\text{gas}}}{dt} \quad (\text{A13})$$

$$\frac{dn_{x+OH}^{\text{aq}}}{dt} = V_{\text{H}_2\text{O}} \cdot \frac{dC_{x+OH}^{\text{aq}}}{dt} \quad (\text{A14})$$

5 We define W as the ratio between the loss rate due to gas phase oxidation and the loss rate due to aqueous oxidation:

$$W \equiv \frac{\frac{dn_{x+OH}^{\text{gas}}}{dt}}{\frac{dn_{x+OH}^{\text{aq}}}{dt}} \quad (\text{A15})$$

We eliminate the dependence of air and water volume with the volumetric liquid water content:

$$10 \quad W = \frac{V_{\text{air}} \cdot \frac{dC_{x+OH}^{\text{gas}}}{dt}}{V_{\text{H}_2\text{O}} \cdot \frac{dC_{x+OH}^{\text{aq}}}{dt}} = \frac{1}{\text{LWC}_v} \cdot \frac{k_{x+OH}^{\text{gas}} C_x^{\text{gas}} C_{\text{OH}}^{\text{gas}}}{k_{x+OH}^{\text{aq}} C_x^{\text{aq}} C_{\text{OH}}^{\text{aq}}} \quad (\text{A16})$$

Inserting the Henry's law constant to describe the ratio of gas and aqueous phase concentrations yields the final working equation:

$$W = \frac{1}{\text{LWC}_v \cdot k_H \cdot R \cdot T} \cdot \frac{k_{x+OH}^{\text{gas}} C_{\text{OH}}^{\text{gas}}}{k_{x+OH}^{\text{aq}} C_{\text{OH}}^{\text{aq}}} \quad (\text{A17})$$

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A3 Derivation of Y

We start with rate laws describing gas phase oxidation by OH and gas phase photolysis:

$$\frac{dC_{x+OH}^{\text{gas}}}{dt} = -k_{x+OH}^{\text{gas}} C_x^{\text{gas}} C_{OH}^{\text{gas}} \quad (\text{A18})$$

5

$$\frac{dC_{x+h\nu}^{\text{gas}}}{dt} = -J_{x+h\nu}^{\text{gas}} C_x^{\text{gas}} \quad (\text{A19})$$

We define the ratio of these rates in molar units as Y:

$$Y \equiv \frac{\frac{dn_{x+OH}^{\text{gas}}}{dt}}{\frac{dn_{x+h\nu}^{\text{gas}}}{dt}} \quad (\text{A20})$$

Since both rates describe the gas phase, the ratio of molar rates is equal to the ratio of concentration-based rates:

10

$$\frac{\frac{dn_{x+OH}^{\text{gas}}}{dt}}{\frac{dn_{x+h\nu}^{\text{gas}}}{dt}} = \frac{\frac{dC_{x+OH}^{\text{gas}}}{dt}}{\frac{dC_{x+h\nu}^{\text{gas}}}{dt}} \quad (\text{A21})$$

Inserting Eqs. (A18) and (A19) into Eq. (A21) yields our final working equation for Y:

$$Y = \frac{k_{x+OH}^{\text{gas}} C_{OH}^{\text{gas}}}{J_{x+h\nu}^{\text{gas}}} \quad (\text{A22})$$

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Appendix B

Chemical sink analysis at 273 K

Z as a function of temperature can be calculated from Z at 298 K. Both Henry's Law constants and for carbonyls, hydration equilibrium constants are sensitive to temperature. When temperature dependent Henry's Law data are available, they are typically fit to the function (Sander et al., 2011):

$$\ln k_{\text{Hx}} = A + B \cdot T^{-1} + C \cdot \ln(T) \quad (\text{B1})$$

where A , B , and C are fitting parameters. Z calculated at T_1 can be adjusted to temperature T_2 with the unitless correction factor, CF_{Hlaw} :

$$\text{CF}_{\text{Hlaw}}^{T_1 \rightarrow T_2} = e^{B \cdot \left(\frac{1}{T_1} - \frac{1}{T_2}\right) + C \cdot \ln\left(\frac{T_1}{T_2}\right)} \quad (\text{B2})$$

Hydration equilibrium constants are also a function of temperature. For a di-carbonyl, Z calculated at T_1 can be adjusted to temperature T_2 with the unitless correction factor, CF_{Khyd} , which accounts for this temperature dependence:

$$\text{CF}_{\text{Khyd}}^{T_1 \rightarrow T_2} = \frac{e^{\frac{\Delta S_{\text{hyd}}}{R} - \frac{\Delta H_{\text{hyd}}}{R \cdot T_2}} + 1}{e^{\frac{\Delta S_{\text{hyd}}}{R} - \frac{\Delta H_{\text{hyd}}}{R \cdot T_1}} + 1} \quad (\text{B3})$$

where ΔS_{hyd} is the change in entropy upon hydration and ΔH_{hyd} is the change in enthalpy upon hydration. Z at T_2 is the product of the correction factors and Z at T_1 :

$$Z(T_2) = \text{CF}_{\text{Hlaw}}^{T_1 \rightarrow T_2} \cdot \text{CF}_{\text{Khyd}}^{T_1 \rightarrow T_2} \cdot Z(T_1) \quad (\text{B4})$$

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The factor W can also be corrected for other temperatures by accounting for differences in both the Henry's Law parameter and the ratio of the gaseous and aqueous rate constants. W calculated at T_1 can be adjusted to its value at T_2 :

$$W(T_2) = CF_{\text{Hlaw}}^{T_1 \rightarrow T_2} \cdot CF_k^{T_1 \rightarrow T_2} \cdot W(T_1) \quad (\text{B5})$$

5 where CF_k accounts for the temperature dependence on the rate constants:

$$CF_k^{T_1 \rightarrow T_2} = \exp \left[\left(\frac{1}{T_2} - \frac{1}{T_1} \right) \left(\frac{E_A^{\text{aq.}}}{R} - \frac{E_A^{\text{gas}}}{R} \right) \right] \quad (\text{B6})$$

where $E_A^{\text{aq.}}$ is the activation energy of the OH oxidation reaction in the aqueous phase and E_A^{gas} is the activation energy of the gas phase reaction.

Using Eqs. (B4) and (B5), we can predict Z and W factors at 273 K from the values calculated at 298 K. The correction factor used to account for the changing gas solubility parameters at different temperatures, CF_{Hlaw} , is dependent on the fitting parameters used to correlate Henry's Law parameters with temperatures (see Eq. B2). For most compounds, these fitting parameters are available in the literature (Sander et al., 2011; Sander, 1999). Typically $C = 0$. For the compounds investigated, B ranges from 4700 K to 7700 K, yielding correction factors of 0.04 to 0.004 going from temperatures of 298 K to 273 K. This correction factor is significant and could change the magnitude of Z by a few orders of magnitude while changes in the percent hydrated have a relatively small effect. Therefore, with the lack of temperature dependant hydration equilibrium data for many relevant compounds and the relatively small effect on Z , we only address changes due to increased Henry's Law parameters at depressed temperatures.

W can also be corrected by CF_k , which captures the temperature dependence in the gas and aqueous OH rate constants. However, the availability of temperature dependent rate data for the compounds investigated is limited. Fortunately, CF_k tends to be on the order of 1 for reasonable temperature corrections. For example, when the activation energy of the aqueous and gaseous reactions differ by 2500 K, (an upper estimate

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from a selection of published OH rate constants for several water soluble compounds) $CF_k = 0.5$. Overestimating or underestimating our value of W by not correcting by this factor will not significantly change the results of our analysis. The lack of published activation energies and the limited influence temperature changes have on CF_k motivate the assumption that CF_k is unity. Y may also depend on temperature. Absorption cross section and quantum yield data are extremely limited, but for the four species for which we determined Y as a function of SZA and altitude, Y was not a strong function of temperature. Therefore, we assume that between 298 K and 273 K the magnitude of Y will not change as to affect the results of our analysis.

Figure A1 shows the magnitude of each chemical sink for the compounds with published Henry's Law temperature dependences at 273 K. In general, aqueous processes play a more significant role due to their enhanced solubility, but for the species investigated, aqueous photolysis still remains a minor sink.

Comparison of the chemical sinksS. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Appendix C

Nomenclature and units

(Variable, typical units, and definition are presented in their order of appearance in the text)

$J[\text{s}^{-1}]$	photolysis rate constant
$F_A[\text{cm}^{-2}\text{s}^{-1}\text{nm}^{-1}]$	actinic flux
$\sigma[\text{cm}^2]$	absorption cross section
Φ [unitless]	quantum yield of photolysis
$\lambda[\text{nm}]$	wavelength of light
$\text{LWC}\left[\frac{\text{kg}_{\text{H}_2\text{O}}}{\text{m}^3_{\text{air}}}\right]$	Liquid water content in terms of the mass of water per volume of air
Z [unitless]	ratio of the rate of gas phase photolysis to the rate of aqueous phase photolysis
n_x [mol]	moles of species x
$R\left[\frac{\text{L atm}}{\text{mol K}}\right]$ or $R\left[\frac{\text{J}}{\text{mol K}}\right]$	ideal gas constant
T [K]	temperature
$\text{LWC}_v\left[\frac{\text{m}^3_{\text{H}_2\text{O}}}{\text{m}^3_{\text{air}}}\right]$	volume based liquid water content (water volume per volume of air)
$k_{\text{Hx}}\left[\frac{\text{mol}_x}{\text{L}_{\text{H}_2\text{O}}\text{atm}}\right]$	Henry's Law constant of species x
$J_{x+h\nu}^{\text{gas}}\left[\text{s}^{-1}\right]$	photolysis rate constant of species x in the gas phase
$J_{x+h\nu}^{\text{aq}}\left[\text{s}^{-1}\right]$	photolysis rate constant of species x in the aqueous phase
t [s]	time
K_{hyd} [unitless]	hydration equilibrium constant
α_N [unitless]	mole fraction of form N in the aqueous phase upon hydration
α_{free} [unitless]	mole fraction of the free form (unhydrated) in solution at equilibrium
α_{hyd} [unitless]	mole fraction of the hydrated form in solution at equilibrium
β_N [unitless]	normalization constant for the photolysis rate constant of form N
β_{free} [unitless]	normalization constant for the photolysis rate constant of the free form
β_{hyd} [unitless]	normalization constant for the photolysis rate constant of the hydrated form
$\beta_{\text{part,hyd}}$ [unitless]	normalization constant for the photolysis rate constant of the partially hydrated form
Y_x [unitless]	constant used to penalize the aqueous photolysis rate of species x for the presence of non-photolabile gem-diol groups
W [unitless]	ratio of the rate of gas phase reaction with OH to the rate of aqueous phase reaction with OH
$k_{x+\text{OH}}^{\text{gas}}\left[\text{m}^3_{\text{air}}\text{mol}^{-1}\text{s}^{-1}\right]$	gas phase rate constant for the reaction of x with OH

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$k_{x+\text{OH}}^{\text{aq}} \left[\text{m}^3_{\text{H}_2\text{O}} \cdot \text{mol}^{-1} \text{s}^{-1} \right]$	aqueous phase rate constant for the reaction of x with OH
$C_{\text{OH}}^{\text{gas}} \left[\frac{\text{mol}_{\text{OH}}}{\text{m}^3_{\text{gas}}} \right]$	concentration of OH in the gas phase
$C_{\text{OH}}^{\text{aq}} \left[\frac{\text{mol}_{\text{OH}}}{\text{m}^3_{\text{H}_2\text{O}}} \right]$	concentration of OH in the aqueous phase
$C_x^{\text{gas}} \left[\frac{\text{mol}_x}{\text{m}^3_{\text{gas}}} \right]$	concentration of species x in the gas phase
$C_x^{\text{aq}} \left[\frac{\text{mol}_x}{\text{m}^3_{\text{H}_2\text{O}}} \right]$	concentration of species x in the aqueous
f_{A^-} [unitless]	mole fraction of the acid that is in the anion form at equilibrium
f_{HA} [unitless]	mole fraction of the acid that is in the acid form at equilibrium
Y [unitless]	ratio of the rate of gas phase reaction with OH to the rate of gas phase photolysis
SZA [deg latitude]	solar zenith angle
z [km]	altitude
Γ [K km^{-1}]	lapse rate
DN [unitless]	parameter (defined in this work) that indicates the ratio of the rate of gas phase reaction with OH relative to the rate of aqueous phase reaction with OH
$CF_{\text{Hlaw}}^{\tau_1 \rightarrow \tau_2}$ [unitless]	correction factor used to adjust Z and W due the temperature dependance of Henry's Law constants
$CF_{\text{Khyd}}^{\tau_1 \rightarrow \tau_2}$ [unitless]	correction factor used to adjust Z and W due the temperature dependance of hydration equilibrium
$\Delta S_{\text{hyd}} \left[\text{J mol}^{-1} \text{K}^{-1} \right]$	change in entropy upon hydration
$\Delta H_{\text{hyd}} \left[\text{J mol}^{-1} \right]$	change in enthalpy upon hydration
$CF_k^{\tau_1 \rightarrow \tau_2}$ [unitless]	correction factor used to adjust W due the temperature dependance of OH rate constants
$E_A^{\text{aq}} \left[\text{J mol}^{-1} \right]$	activation energy of species x in the aqueous phase
$E_A^{\text{gas}} \left[\text{J mol}^{-1} \right]$	activation energy of species x in the gas phase
$\rho_{\text{H}_2\text{O}} \left[\frac{\text{kg}_{\text{H}_2\text{O}}}{\text{m}^3_{\text{H}_2\text{O}}} \right]$	the density of water
$V_{\text{air}} \left[\text{m}^3_{\text{air}} \right]$	air volume
$V_{\text{H}_2\text{O}} \left[\text{m}^3_{\text{H}_2\text{O}} \right]$	water volume

Supplementary material related to this article is available online at:
[http://www.atmos-chem-phys-discuss.net/12/10015/2012/
acpd-12-10015-2012-supplement.pdf](http://www.atmos-chem-phys-discuss.net/12/10015/2012/acpd-12-10015-2012-supplement.pdf).

Acknowledgements. The authors would like to acknowledge the National Science Foundation grants ATM-0831518 and CHE-0909227 for funding this work.

References

Andreae, M. O. and Crutzen, P. J.: Atmospheric Aerosols: Biogeochemical Sources and Role in Atmospheric Chemistry, *Science*, 276, 1052–1058, doi:10.1126/science.276.5315.1052, 1997.

Atkinson, R.: A structure-activity relationship for the estimation of rate constants for the gas-phase reactions of OH radicals with organic compounds, *Int. J. Chem. Kinet.*, 19, 799–828, doi:10.1002/kin.550190903, 1987.

Atkinson, R.: Estimation of gas-phase hydroxyl radical rate constants for organic chemicals, *Environ. Toxicol. Chem.*, 7, 435–442, doi:10.1002/etc.5620070604, 1988.

Atkinson, R.: Gas-phase tropospheric chemistry of organic compounds: A review, *Atmos. Environ. A-Gen.*, 24, 1–41, 1990.

Atkinson, R.: Kinetics and Mechanisms of the Gas-Phase Reactions of the NO₃ Radical with Organic Compounds, *J. Phys. Chem. Ref. Data*, 20, 459–507, 1991.

Atkinson, R. and Arey, J.: Atmospheric Degradation of Volatile Organic Compounds, *Chem. Rev.*, 103, 4605–4638, doi:10.1021/cr0206420, 2003.

Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and IUPAC Subcommittee: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II – gas phase reactions of organic species, *Atmos. Chem. Phys.*, 6, 3625–4055, doi:10.5194/acp-6-3625-2006, 2006.

Audiffren, N., Renard, M., Buisson, E., and Chaumerliac, N.: Deviations from the Henry's law equilibrium during cloud events: a numerical approach of the mass transfer between phases and its specific numerical effects, *Atmos. Res.*, 49, 139–161, 1998.

ACPD

12, 10015–10058, 2012

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Comparison of the
chemical sinks**S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Bailey, P. S.: The Reactions of Ozone with Organic Compounds, *Chem. Rev.*, 58, 925–1010, 1958.
- Bateman, A. P., Nizkorodov, S. A., Laskin, J., and Laskin, A.: Photolytic processing of secondary organic aerosols dissolved in cloud droplets, *Phys. Chem. Chem. Phys.*, 13, 12199–12212, 2011.
- 5 Bell, R. P. and Gold, V.: The Reversible Hydration of Carbonyl Compounds, in: *Advances in Physical Organic Chemistry*, Academic Press, 1–29, 1966.
- Bell, R. P. and McDougall, A. O.: Hydration equilibria of some aldehydes and ketones, *T. Faraday Soc.*, 56, 1281–1285, 1960.
- 10 Buxton, G. V., Greenstock, C. L., Helman, W. P., and Ross, A. B.: Critical Review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals in aqueous Solution, *J. Phys. Chem. Ref. Data*, 17, 513–886, 1988.
- Chebbi, A. and Carlier, P.: Carboxylic acids in the troposphere, occurrence, sources, and sinks: A review, *Atmos. Environ.*, 30, 4233–4249, 1996.
- 15 Chu, L. and Anastasio, C.: Formation of Hydroxyl Radical from the Photolysis of Frozen Hydrogen Peroxide, *J. Phys. Chem. A*, 109, 6264–6271, doi:10.1021/jp051415f, 2005.
- De Bruyn, W. J., Swartz, E., Hu, J. H., Shorter, J. A., Davidovits, P., Worsnop, D. R., Zahniser, M. S., and Kolb, C. E.: Henry's law solubilities and Setchenow coefficients for biogenic reduced sulfur species obtained from gas-liquid uptake measurements, *J. Geophys. Res.*, 100, 7245–7251, doi:10.1029/95jd00217, 1995.
- 20 Donahue, N. M., Robinson, A. L., and Pandis, S. N.: Atmospheric organic particulate matter: from smoke to secondary organic aerosol, *Atmos. Environ.*, 43, 94–106, 2009.
- Dong, S. and Dasgupta, P. K.: Solubility of gaseous formaldehyde in liquid water and generation of trace standard gaseous formaldehyde, *Environ. Sci. Technol.*, 20, 637–640, doi:10.1021/es00148a016, 1986.
- 25 Epstein, S. A., Shemesh, D., Tran, V. T., Nizkorodov, S. A., and Gerber, R. B.: Absorption spectra and photolysis of methyl peroxide in liquid and frozen water, *J. Phys. Chem. A*, doi:10.1021/jp211304v, 2012.
- Ervens, B. and Volkamer, R.: Glyoxal processing by aerosol multiphase chemistry: towards a kinetic modeling framework of secondary organic aerosol formation in aqueous particles, *Atmos. Chem. Phys.*, 10, 8219–8244, doi:10.5194/acp-10-8219-2010, 2010.
- 30

- Ervens, B. and Volkamer, R.: Glyoxal processing by aerosol multiphase chemistry: towards a kinetic modeling framework of secondary organic aerosol formation in aqueous particles, *Atmos. Chem. Phys.*, 10, 8219–8244, doi:10.5194/acp-10-8219-2010, 2010.
- 5 Ervens, B., George, C., Williams, J. E., Buxton, G. V., Salmon, G. A., Bydder, M., Wilkinson, F., Dentener, F., Mirabel, P., Wolke, R., and Herrmann, H.: CAPRAM 2.4 (MODAC mechanism): An extended and condensed tropospheric aqueous phase mechanism and its application, *J. Geophys. Res.*, 108, 4426, doi:10.1029/2002jd002202, 2003a.
- Ervens, B., Gligorovski, S., and Herrmann, H.: Temperature-dependent rate constants for hydroxyl radical reactions with organic compounds in aqueous solutions, *Phys. Chem. Chem. Phys.*, 5, 1811–1824, 2003b.
- 10 Finlayson-Pitts, B. J. and Pitts, J. N.: *Chemistry of the Upper and Lower Atmosphere*, Academic Press, San Diego, 198 pp., 2000.
- Gelencsér and Varga: Evaluation of the atmospheric significance of multiphase reactions in atmospheric secondary organic aerosol formation, *Atmos. Chem. Phys.*, 5, 2823–2831, doi:10.5194/acp-5-2823-2005, 2005.
- 15 Goldstein, A. H. and Galbally, I. E.: Known and Unexplored Organic Constituents in the Earth's Atmosphere, *Environ. Sci. Technol.*, 41, 1514–1521, doi:10.1021/es072476p, 2007.
- Graedel, T. E. and Goldberg, K. I.: Kinetic Studies of Raindrop Chemistry 1. Inorganic and Organic Processes, *J. Geophys. Res.*, 88, 10865–10882, doi:10.1029/JC088iC15p10865, 1983.
- 20 Graedel, T. E. and Weschler, C. J.: Chemistry within aqueous atmospheric aerosols and raindrops, *Rev. Geophys.*, 19, 505–539, doi:10.1029/RG019i004p00505, 1981.
- Haag, W. R. and Yao, C. C. D.: Rate constants for reaction of hydroxyl radicals with several drinking water contaminants, *Environ. Sci. Technol.*, 26, 1005–1013, doi:10.1021/es00029a021, 1992.
- 25 Hummel, J. R. and Reck, R. A.: A Global Surface Albedo Model, *J. Appl. Meteorol.*, 18, 239–253, doi:10.1175/1520-0450(1979)18[239:agsam]2.0.co;2, 1979.
- Kolb, C. E., Jayne, J. T., Worsnop, D. R., and Davidovits, P.: Solubility data requirements and new experimental methods in atmospheric aerosol research, *Pure and applied chemistry*, Research Triangle Park, NC, ETATS-UNIS, 10 pp., 1997.
- 30 Kwok, E. S. C. and Atkinson, R.: Estimation of hydroxyl radical reaction rate constants for gas-phase organic compounds using a structure-reactivity relationship: An update, *Atmos. Environ.*, 29, 1685–1695, 1995.

Comparison of the chemical sinksS. A. Epstein and
S. A. Nizkorodov

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Comparison of the
chemical sinks**S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Le Botlan, D. J., Mechin, B. G., and Martin, G. J.: Proton and carbon-13 nuclear magnetic resonance spectrometry of formaldehyde in water, *Anal. Chem.*, 55, 587–591, doi:10.1021/ac00254a041, 1983.

Lee, M., Heikes, B. G., and O'Sullivan, D. W.: Hydrogen peroxide and organic hydroperoxide in the troposphere: a review, *Atmos. Environ.*, 34, 3475–3494, 2000.

Limbeck, A. and Puxbaum, H.: Dependence of in-cloud scavenging of polar organic aerosol compounds on the water solubility, *J. Geophys. Res.*, 105, 19857–19867, doi:10.1029/2000jd900123, 2000.

Madronich, S.: Photodissociation in the Atmosphere 1. Actinic Flux and the Effects of Ground Reflections and Clouds, *J. Geophys. Res.*, 92, 9740–9752, doi:10.1029/JD092iD08p09740, 1987.

Marini, A., Munoz-Losa, A., Biancardi, A., and Mennucci, B.: What is Solvatochromism?, *J. Phys. Chem. B*, 114, 17128–17135, doi:10.1021/jp1097487, 2010.

Monod, A., Poulain, L., Grubert, S., Voisin, D., and Wortham, H.: Kinetics of OH-initiated oxidation of oxygenated organic compounds in the aqueous phase: new rate constants, structure activity relationships and atmospheric implications, *Atmos. Environ.*, 39, 7667–7688, 2005.

Moortgat, G. K.: Important photochemical processes in the atmosphere, *Pure Appl. Chem.*, 73, 487–490, 2001.

Nozriere, B., Dziedzic, P., and Cordova, A.: Products and Kinetics of the Liquid-Phase Reaction of Glyoxal Catalyzed by Ammonium Ions (NH₄⁺), *J. Phys. Chem. A*, 113, 231–237, doi:10.1021/jp8078293, 2008.

Pankow, J. F. and Asher, W. E.: SIMPOL.1: a simple group contribution method for predicting vapor pressures and enthalpies of vaporization of multifunctional organic compounds, *Atmos. Chem. Phys.*, 8, 2773–2796, doi:10.5194/acp-8-2773-2008, 2008.

Rohrer, F. and Berresheim, H.: Strong correlation between levels of tropospheric hydroxyl radicals and solar ultraviolet radiation, *Nature*, 442, 184–187, 2006.

Ruggaber, A., Dlugi, R., Bott, A., Forkel, R., Herrmann, H., and Jacobi, H. W.: Modelling of radiation quantities and photolysis frequencies in the aqueous phase in the troposphere, *Atmos. Environ.*, 31, 3137–3150, 1997.

Saxena, P. and Hildemann, L. M.: Water-soluble organics in atmospheric particles: A critical review of the literature and application of thermodynamics to identify candidate compounds, *J. Atmos. Chem.*, 24, 57–109, doi:10.1007/bf00053823, 1996.

**Comparison of the
chemical sinks**S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Schwarzenbach, R. P., Gschwend, P. M., and Imboden, D. M.: Frontmatter, in: Environmental Organic Chemistry, John Wiley & Sons, Inc., i-xiii, 2005.

Seinfeld, J. and Pandis, S.: Atmospheric Chemistry and Physics, John Wiley and Sons, Inc., New York, USA, 1998.

5 Vaitilingom, M., Amato, P., Sancelme, M., Laj, P., Leriche, M., and Delort, A. M.: Contribution of microbial activity to carbon chemistry in clouds, *Appl. Environ. Microb.*, 76, 23, doi:10.1128/AEM.01127-09, 2010.

Vaitilingom, M., Charbouillot, T., Deguillaume, L., Maisonobe, R., Parazols, M., Amato, P., Sancelme, M., and Delort, A. M.: Atmospheric chemistry of carboxylic acids: microbial im-
10 plication versus photochemistry, *Atmos. Chem. Phys.*, 11, 8721–8733, doi:10.5194/acp-11-8721-2011, 2011.

Vione, D., Maurino, V., Minero, C., Pelizzetti, E., Harrison, M. A. J., Olariu, R.-I., and Arsene, C.: Photochemical reactions in the tropospheric aqueous phase and on particulate matter, *Chem. Soc. Rev.*, 35, 441–453, 2006.

15 Wallace, J. M. and Hobbs, P. V.: Atmospheric Science: An Introductory Survey, International Geophysics Series, edited by: Dmowska, R., Hartmann, D., and Rossby, H. T., Academic Press, 2006.

Winiwarter, W., Fierlinger, H., Puxbaum, H., Facchini, M. C., Arends, B. G., Fuzzi, S., Schell, D., Kaminski, U., Pahl, S., Schneider, T., Berner, A., Solly, I., and Krusz, C.: Henry's law and the behavior of weak acids and bases in fog and cloud, *J. Atmos. Chem.*, 19, 173–188, doi:10.1007/bf00696588, 1994.

20 Xu, H., Wentworth, P. J., Howell, N. W., and Joens, J. A.: Temperature dependent near-UV molar absorptivities of aliphatic aldehydes and ketones in aqueous solution, *Spectrochim. Acta A-M*, 49, 1171–1178, 1993.

25 Zellner, R., Exner, M., and Herrmann, H.: Absolute OH quantum yields in the laser photolysis of nitrate, nitrite and dissolved H₂O₂ at 308 and 351 nm in the temperature range 278–353 K, *J. Atmos. Chem.*, 10, 411–425, doi:10.1007/bf00115783, 1990.

30 Zepp, R. G., Baughman, G. L., and Schlotzhauer, P. F.: Comparison of photochemical behavior of various humic substances in water: I. Sunlight induced reactions of aquatic pollutants photosensitized by humic substances, *Chemosphere*, 10, 109–117, 1981.

Comparison of the chemical sinks


 S. A. Epstein and
 S. A. Nizkorodov


Fig. 1. Schematic diagram detailing the four chemical sinks being investigated in this manuscript. Dotted \dot{n} represents the rate of change in moles per time per volume of air parcel due to each of the four processes: gaseous oxidation by OH (\dot{n}_{gas}^{OHox}), aqueous oxidation by OH ($\dot{n}_{aq.}^{OHox}$), gaseous photolysis ($\dot{n}_{gas}^{photolysis}$), and aqueous photolysis ($\dot{n}_{aq.}^{photolysis}$). The ratios of the removal rates W , Z , and Y are defined in the text; they are independent of the air parcel volume. From these three ratios, the relative rates of any of these four processes can be calculated; an example is shown with a broken arrow.

[Title Page](#)
[Abstract](#) | [Introduction](#)
[Conclusions](#) | [References](#)
[Tables](#) | [Figures](#)
[◀](#) | [▶](#)
[◀](#) | [▶](#)
[Back](#) | [Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 2. Best estimate for the maximum ratio of gas phase photolysis to aqueous phase photolysis, Z , for a series of peroxides and methyl nitrate in an air parcel with a LWC = 3 g m^{-3} at 298 K. Species are labeled in order of increasing Z : (1) hydrogen peroxide, (2) ethyl peroxide, (3) methyl peroxide, and (4) methyl nitrate. The two inward arrows adjacent to “1” show the range of Z for hydrogen peroxide across all solar zenith angles with experimentally measured gaseous and aqueous rate constants from (Sander et al., 2011; Chu and Anastasio, 2005).

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Best estimate for the maximum ratio of gas phase photolysis to aqueous phase photolysis, Z , for a series of carbonyls in an air parcel with a LWC = 3 g m^{-3} at 298 K. The “>” label next to species 13 indicates that it lies at $Z \gg 10^4$ where $J_{\text{aqueous}} = 0$. Species are labeled in order of increasing Z : (1) glyceraldehyde, (2) pyruvic acid, (3) gluteraldehyde, (4) methyl glyoxal, (5) glyoxal, (6) glycolaldehyde, (7) glyoxylic acid, (8) acetone, (9) butanone, (10) butanal, (11) propanal, (12) acetaldehyde, and (13) formaldehyde.

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov




Fig. 4. The ratio of the rate of gas phase OH oxidation to the rate of aqueous phase oxidation for a series of peroxides and ethers in an air parcel with a liquid water content of 3 g m^{-3} at 298 K. Labeled horizontal lines are bound by differences in $C_{\text{OH}}^{\text{gas}}/C_{\text{OH}}^{\text{aqueous}}$ (defined as the moles of OH per volume of air/the moles of OH per volume of liquid) during different times of the day. The line demarcating OH at Henry's law equilibrium is shown on the left side of the plot. Species are labeled in order of increasing W : (1) hydrogen peroxide, (2) ethyl peroxide, (3) methyl peroxide, (4) methyl tert-butyl ether, (5) dimethyl ether, and (6) ethyl ether.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov




Fig. 5. The ratio of the rate of gas phase OH oxidation to the rate of aqueous phase oxidation for a series of carbonyls in an air parcel with a liquid water content of 3 g m^{-3} at 298 K. Labeled horizontal lines are bound by differences in $C_{\text{OH}}^{\text{gas}}/C_{\text{OH}}^{\text{aqueous}}$ (defined as the moles of OH per volume of air / the moles of OH per volume of liquid) during different times of the day. The line demarcating OH at Henry's law equilibrium is shown on the left side of the plot. Species are labeled in order of increasing W : (1) glyoxal, (2) methyl glyoxal, (3) formaldehyde, (4) acetone, (5) 2-butanone, (6) acetaldehyde, (7) butanal, and (8) propanal.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. The ratio of the rate of gas phase OH oxidation to the rate of aqueous phase oxidation for a series of acids in an air parcel with a liquid water content of 3 g m^{-3} at 298 K. Labeled rectangles are bound by differences in $C_{\text{OH}}^{\text{gas}}/C_{\text{OH}}^{\text{aqueous}}$ (defined as the moles of OH per volume of air / the moles of OH per volume of liquid) during different times of the day and the pH of the cloud or fog droplet ranging between 2 and 6. The line demarcating OH at Henry's law equilibrium is shown on the left side of the plot. Species are labeled in order of increasing W at $\text{pH} = 6$: (1) tartaric acid (R,R), (2) malic acid, (3) malonic acid, (4) lactic acid, (5) succinic acid, (6) pyruvic acid, (7) formic acid, (8) propanoic acid, (9) glyoxylic acid, (10) 2-methyl propanoic acid, (11) pentanoic acid, (12) hexanoic acid, (13) acetic acid, (14) 3-methyl butanoic acid, and (15) 2,2-dimethyl 1-propanoic acid.

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 7. The ratio of the rate of gas phase OH oxidation to the rate of aqueous phase oxidation, W as a function of the dimensionless number, $DN = (LWC_V \cdot R \cdot T \cdot k_H)^{-1}$ for the series of compounds investigated under daytime OH concentrations in an air parcel with a $LWC = 3 \text{ g m}^{-3}$ at 298 K. The magenta dashed line indicates the transition between gaseous reactions being more important, $DN \gg 10$ and aqueous reactions being more important, $DN \ll 10$. Similar plots with nighttime OH concentrations and $LWC = .001 \text{ g m}^{-3}$ are shown in the Supplement and yield an identical DN cutoff.

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov




Fig. 8. The ratio of the rate of gas phase OH oxidation to the rate of gaseous photolysis, Y , as a function of solar zenith angle, SZA, for compounds where the absorption cross sections and quantum yields are published, at 298 K. The width of the red swath is due to the uncertainty in the empirical function relating OH concentration to the rate of O_3 photolysis. The dashed line indicates where the rate of gas phase OH oxidation is equal to the rate of gaseous photolysis, $Y = 1$.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Comparison of the chemical sinksS. A. Epstein and
S. A. Nizkorodov

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 9. The ratio of the rate of gas phase OH oxidation to the rate of gaseous photolysis, Y , as a function of solar zenith angle and altitude for compounds where the absorption cross sections and quantum yields are published as a function of temperature and pressure. Upper and lower limits of these plots, due to the uncertainty in the empirical function relating OH concentration to the rate of photolysis of O^1D , are presented in the Supplement.

Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov




Fig. 10. Estimated magnitudes of each of the 4 chemical sinks investigated for compounds with sufficient published data. All acids are assumed to be in cloud or fog droplets at a pH = 6. Calculations were performed with a solar zenith angle of 20° . The right panel details selected species in a parcel with a LWC = 3 g m^{-3} at 298 K and represents the upper limit for the contribution of aqueous processes in all situations at 298 K. The left panel shows the contribution of each of the 4 sinks in an air parcel with a more realistic LWC = 0.05 g m^{-3} at 298 K. The contributions of aqueous processes in this panel are also an upper estimate for the selected LWC.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Comparison of the chemical sinks

S. A. Epstein and
S. A. Nizkorodov




Fig. A1. Estimated magnitudes of each of the 4 chemical sinks investigated for compounds with sufficient published data. All acids are assumed to be in cloud or fog droplets at a pH = 6. Calculations were performed with a solar zenith angle of 20°. The right panel details selected species in a parcel with a LWC = 3 g m⁻³ at 273 K and represents the upper limit for the contribution of aqueous processes in all situations at 273 K. The left panel shows the contribution of each of the 4 sinks in an air parcel with a more realistic LWC = 0.05 g m⁻³ at 273 K. The contributions of aqueous processes in this panel are also an upper estimate for the selected LWC.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
