

This discussion paper is/has been under review for the journal *Climate of the Past* (CP).
Please refer to the corresponding final paper in CP if available.

A collection of sub-daily pressure and temperature observations for the early instrumental period with a focus on the “year without a summer” 1816

Y. Brugnara^{1,2}, R. Auchmann^{1,2}, S. Brönnimann^{1,2}, R. J. Allan³, I. Auer⁴, M. Barriandos⁵, H. Bergström⁶, J. Bhend⁷, R. Brázdil^{8,9}, G. P. Compo^{10,11}, R. C. Cornes¹², F. Dominguez-Castro^{13,14}, A. F. V. van Engelen¹⁵, J. Filipiak¹⁶, J. Holopainen¹⁷, S. Jourdain¹⁸, M. Kunz¹⁹, J. Luterbacher²⁰, M. Maugeri²¹, L. Mercalli²², A. Moberg^{23,24}, C. J. Mock²⁵, G. Pichard²⁶, L. Řezníčková^{8,9}, G. van der Schrier¹⁵, V. Slonosky²⁷, Z. Ustrnul²⁸, M. A. Valente²⁹, A. Wypych²⁸, and X. Yin³⁰

¹Oeschger Centre for Climate Change Research, Bern, Switzerland

²Institute of Geography, University of Bern, Bern, Switzerland

³Hadley Centre, Met Office, Devon, UK

⁴ZAMG-Central Institute for Meteorology and Geodynamics, Vienna, Austria

⁵Department of Modern History, University of Barcelona, Barcelona, Spain

⁶Department of Earth Sciences, Uppsala University, Uppsala, Sweden

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A collection of sub-daily observations

Y. Brugnara et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



⁷Federal Office of Meteorology and Climatology, MeteoSwiss, Zurich, Switzerland

⁸Institute of Geography, Masaryk University, Brno, Czech Republic

⁹Global Change Research Centre, Academy of Sciences of the Czech Republic, Brno, Czech Republic

¹⁰Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA

¹¹Physical Sciences Division, Earth System Research Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO, USA

¹²CRU-Climatic Research Unit, School of Environmental Sciences, University of East Anglia, Norwich, UK

¹³Department of Physics, Universidad de Extremadura, Badajoz, Spain

¹⁴Departamento de Ingeniería Civil y Ambiental, Escuela Politécnica Nacional, Quito, Ecuador

¹⁵KNMI-Royal Netherlands Meteorological Institute, De Bilt, the Netherlands

¹⁶Institute of Geography, University of Gdańsk, Gdańsk, Poland

¹⁷Department of Geosciences and Geography, University of Helsinki, Helsinki, Finland

¹⁸Météo-France, Direction de la Climatologie, Toulouse, France

¹⁹Institute for Meteorology and Climate Research (IMK), Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

²⁰Department of Geography, Climatology, Climate Dynamics and Climate Change, Justus Liebig University of Giessen, Giessen, Germany

²¹Università degli Studi di Milano, Department of Physics, Milan, Italy

²²SMI-Società Meteorologica Italiana, Turin, Italy

²³Department of Physical Geography, Stockholm University, Stockholm, Sweden

²⁴Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

²⁵Department of Geography, University of South Carolina, Columbia, SC, USA

²⁶Department of History, Université Aix-Marseille, Aix-en-Provence, France

²⁷McGill University, Centre for Interdisciplinary Studies on Montreal, Montreal, Canada

²⁸Jagiellonian University, Department of Climatology, Krakow, Poland

²⁹Instituto Dom Luiz, Faculdade de Ciências da Universidade de Lisboa, Portugal

³⁰ERT, Inc., Asheville, NC, USA

Received: 7 April 2015 – Accepted: 26 April 2015 – Published: 13 May 2015

Correspondence to: Y. Brugnara (yuri.brugnara@giub.unibe.ch)

Published by Copernicus Publications on behalf of the European Geosciences Union.

CPD

11, 1741–1794, 2015

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The eruption of Mount Tambora (Indonesia) in April 1815 is the largest documented volcanic eruption in history. It caused a large global cooling during the following year, felt particularly in parts of Europe and North America, where the year 1816 became known as the “year without a summer”. This paper describes an effort made to collect surface meteorological observations from the early instrumental period, with a focus on the years immediately following the eruption (1815–1817). Although the collection aimed in particular at pressure observations, correspondent temperature observations were also recovered. Some of the series had already been described in the literature, but a large part of the data, recently digitised from original weather diaries and contemporary magazines and newspapers, is presented here for the first time. The collection puts together more than fifty sub-daily series from land observatories in Europe and North America, and from ships in the tropics. The pressure observations have been corrected for temperature and gravity and reduced to mean sea level, moreover an additional statistical correction was applied to take into account common error sources in mercury barometers. To assess the reliability of the corrected dataset, the variance of the pressure observations is compared with modern climatologies and single observations are used for synoptic analyses of three case studies in Europe. All raw observations will be made available to the scientific community in the International Surface Pressure Databank.

1 Introduction

The measurement of atmospheric pressure has a long history, which begins with the famous experiment of Evangelista Torricelli in 1643. It was not long until, in 1657, a first European network of meteorological observatories, all equipped with a barometer, was set up by the Accademia del Cimento (Middleton, 1972). Similar short-lived attempts of organized networks would follow in the 18th century (e.g., Kington, 1974; Moberg,

CPD

11, 1741–1794, 2015

**A collection of
sub-daily
observations**

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1998; Brázdil et al., 2008). Eventually the barometer, as well as the thermometer, became a commercial product and an object of desire for anybody interested in the natural sciences, including not only scientists but also educated individuals from the middle and high classes, like physicians or clergymen. Some of them used to keep meteorological diaries, in the same way the scientists in the astronomical observatories and in some university had began to do. This phenomenon led to the recording of millions of pressure and temperature observations, at the beginning only in Europe, but gradually also in the various ocean basins, on board of intercontinental ships, and finally in the colonies. The French revolution and the Napoleonic wars caused a temporary decline in the quantity of meteorological observations in some European countries between the end of the 18th century and the beginning of the 19th century, accompanied by the dissolution of existing meteorological networks, but in the meantime the quality of the instruments continued to progress. Finally, in the 1850s a new era for meteorology began with the creation of the first national weather services. These two centuries of development of the basic instruments for the atmospheric sciences are usually referred as the “early instrumental period”.

Between the 1990s and the 2000s, three European Union-funded projects, ADVICE, IMPROVE and EMULATE (Jones et al., 1999; Camuffo and Jones, 2002; Ansell et al., 2006), triggered a large effort to digitise historical observations of temperature and pressure, particularly those of long and continuous series, some longer than 250 years, which were in some cases corrected and homogenised. These projects marked an important development from earlier manual efforts, which also sought to use historic barometric pressure observations to analyse changes in the atmospheric circulation but which were limited by an inability to automate the calculations (Cornes, 2014). A few years ago, most of the existing digitised pressure observations were collected and successfully assimilated into a global reanalysis that reconstructed four-dimensional meteorological fields back to 1870 (Compo et al., 2006, 2011), recently extended further back to 1850.

**A collection of
sub-daily
observations**

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The collection described in this paper represents a first step towards a reanalysis of the first half of the 19th century. Although some of the series cover longer periods, the focus is on the years 1815–1817, the period most influenced by the eruption of Mount Tambora in Indonesia.

Located on the island of Sunbawa, about 300 km east of Bali, Tambora erupted between 10 and 11 April 1815 (Stothers, 1984; Oppenheimer, 2003). The consequences were a significant global cooling, estimated between 0.5 and 1 K (e.g., Wagner and Zorita, 2005; Kandlbauer et al., 2013), as well as more delayed changes in the atmospheric circulation that deeply affected the climate of the mid-latitudes in the Northern Hemisphere (e.g., Fischer et al., 2007; Wegmann et al., 2014). The highlight of this period is the infamous “year without a summer” (Stommel and Stommel, 1979), 1816, a year characterised by strong and persistent negative temperature anomalies in Central and Western Europe (e.g., Luterbacher and Pfister, 2015, and references therein) as well as in eastern North America (e.g., Chenoweth, 1996; Briffa et al., 1998) during the growing season, with major socio-economical impacts due to widespread crop failures (e.g., Pfister, 1999). Comparable impacts might also have been triggered by Tambora through an exceptional winter drought in most of the Iberian Peninsula (Trigo et al., 2009; Domínguez-Castro et al., 2012). The global crisis caused by the 1816 climate anomaly has been described as “the last great subsistence crisis in the Western World” (Post, 1977).

Despite the large amount of meteorological observations available for the early instrumental period, only a small part has been exploited in modern climate research (Brönnimann et al., 2006). The huge amount of documents, spread over thousands of libraries and archives, and the significant financial and human investments needed for recovery and digitisation, explain why the majority of the data have never been analysed so far. Another difficulty arises from data quality, in particular for temperature: the homogenisation with modern data is usually not an easy task (e.g., Camuffo, 2002a, b; Böhm et al., 2010). Pressure is to some extent less problematic, when accompanied by detailed metadata, because the barometer does not have to be outdoors. Obser-

CPD

11, 1741–1794, 2015

A collection of sub-daily observations

Y. Brugnara et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**A collection of
sub-daily
observations**

Y. Brugnara et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 vations made with mercury barometers need however several corrections based on
the characteristics of the barometer, on the variations of temperature and on the lat-
titude (e.g., Moberg et al., 2002; Camuffo et al., 2006). Unfortunately, in most cases
the observations were registered without any correction and it is usually very difficult,
if not impossible, to find any information about the barometer. The temperature of the
barometer, fundamental for the correction, was also often not reported. This means
that assumptions have to be made, which increase the uncertainty of the original ob-
servations. Despite this, we will show that most of the data can be retained for scientific
use.

10 Even though a recognised official standard for outside temperature measurement
did not exist in the early 19th century, some common rules had been long agreed
inside the scientific community, mainly inspired by the recommendations of French
physicist Réaumur (Réaumur, 1732). Thermometers were usually placed on north-
facing walls or windows to minimize the effect of direct and indirect sunlight. In some
cases, an iron screen was used to shield the instrument from solar radiation (e.g.,
15 Camuffo, 2002c). We do not correct temperature observations in this work and we
make a limited use of them in the analysis. However, we use outside temperature to
reduce pressure observations to sea level and sometimes also to correct the thermal
expansion of the mercury in the barometer, when the temperature of the barometer
is not available. Böhm et al. (2010) calculated that at the Kremsmünster observatory
20 (Austria), when the sunlight hits the historical thermometer location (northeast-facing
window) in summer, the average overestimation in the observed temperature is of about
2 K, although in the most extreme cases it can reach even 5 K. Errors of this magnitude
have a negligible effect on the reduction of pressure observations to sea level at low
25 elevation.

internet (Google Books and similar) have grown considerably over the last years and gave an important contribution to the collection. In particular, contemporary scientific magazines have proven to be a prolific source.

Table 1 summarises the main characteristics of each land record. Almost half of the series unfortunately do not have the temperature of the barometer, nor were the pressure observations corrected for temperature. In fact, one can distinguish between two categories of observatories: the scientific observatories and the “amateurs”. The former category includes astronomical observatories, universities and other scientific organizations. It offers in general a higher scientific level, since the observations were carried out by professional scientists, usually astronomers or physicists. Moreover, metadata are more abundant and detailed. These observatories are printed in bold in Table 1. The amateurs were sometimes scientists who kept a personal weather diary, but in most cases they were learned and wealthy individuals (physicians, aristocrats, clergymen, etc.) with a strong interest in the natural sciences. Their measurements may be in general less accurate and information about corrections or the temperature of the barometer are rarely given. Metadata are sometimes completely absent or very difficult to find. A few stations belonging to this category can actually be considered at the borderline, in the sense that their activity was supervised by a scientific institution, which often provided the instruments, following the model of the *Societas Meteorologica Palatina* in the 18th century (see Kington, 1974). This is the case for most of the observation sites in Sweden (Moberg, 1998) and for Hohenpeissenberg (Germany), where the monks of a monastery kept a meteorological register for the Bavarian Academy of Sciences (Winkler, 2006).

The series of Paris is split into two parts because we had different sources: University of Barcelona provided one uncorrected pressure observation per day in the period 1811–1820, digitised from the original registers of the Paris astronomical observatory (Cornes et al., 2012), while four observations per day in the period 1816–1817, corrected for temperature, were digitised at the University of Bern from the *Annales de chimie et de physique*, where the corrected observations of the astronomical observa-

CPD

11, 1741–1794, 2015

**A collection of
sub-daily
observations**

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tory were published starting in 1816. There is no overlap between the two series (we removed the 1816–1817 data in the first series).

Ships' log-books also contain pressure and air temperature observations, and sometimes sea surface temperature (which was also digitised). The four records in the collection are from British vessels, they are briefly described in Table 2.

2.2 Pressure measurement in the early 19th century

At the beginning of the 19th century many different models of mercury barometers were employed for meteorological observations. They can be divided into three main categories: the fixed-cistern barometer, the Fortin barometer and the siphon barometer. Probably a fourth category should be reserved to marine barometers, which needed a special construction to be employed on moving ships.

The fixed-cistern barometer is an adaptation of the original experiment of Torricelli and was the most commonly used barometer in the early 19th century: it is composed of a cistern, where the mercury is exposed to the air pressure, and a vertical thin glass tube, closed on its upper end (where a vacuum is created) and equipped with a scale (either engraved directly on the tube or fixed externally, sometimes together with a vernier to increase the resolution), while its open end is immersed in the mercury of the cistern. The mercury is in hydrostatic equilibrium with the air, a change of the air pressure causes a change of the level of the mercury in the tube and a (smaller) change of the level in the cistern. A correction, calculated from the dimensions of the cistern and of the tube, must be applied on the readings made on the tube to take into account the change of the level of the mercury in the cistern.

The correction is avoided in the case of the Fortin barometer, which is provided with a variable displacement cistern, where the level of the mercury has to be set to the zero (marked by the tip of an ivory pin) through a screw before the pressure value is read on the column (Fig. 2). The Fortin barometer is named after its inventor, the French instrument-maker Jean Nicolas Fortin. Techniques to keep the level in the cistern constant (or to measure it) already existed in the 18th century (e.g., overflowing

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cisterns, leather bags, flowing gauges, etc.), but none of them had the success of Fortin's model, which was introduced at the beginning of the 19th century. At the time of the Tambora eruption, the Fortin barometer was a relatively new invention and only a very limited number of observatories had one.

5 Siphon barometers do not have a cistern, instead the tube is u-shaped at the bottom and the end of the shorter leg is exposed to air; the level of the mercury in both legs of the tube is needed to obtain the pressure value. The siphon barometer was often criticised at the time inside the scientific community, because of the additional reading required, the lack of transportability and the exposition of the mercury to dust, humidity and oxidation, which could affect the reliability of the measurements. Nevertheless, it maintained numerous advocates among scientists in Central Europe (Middleton, 1964).
10 In 1816 Joseph Louis Gay-Lussac eventually developed a transportable siphon barometer which temporarily increased the popularity of this kind of barometer in the following years.

15 Independently from the barometer's model, further corrections due to the thermal expansion of mercury and the change of gravity with latitude are necessary. In some cases, the capillarity inside the tube and the construction of the scale are also sources of significant errors and drifts (e.g., Camuffo et al., 2006), as is a lack of maintenance and many other factors. The reader is referred to Middleton (1964) for a detailed
20 overview on errors affecting observations made with mercury barometers.

From metadata we know what type of barometer was employed in 1815–1817 for only twelve observatories in the collection. Seven of them (Cambridge, Haarlem, Hohenpeissenberg, London, Stockholm, Vienna and Zwanenburg) employed fixed-cistern barometers, three (Aarau, Düsseldorf and Padua) had siphon barometers, two (Milan and Bologna) had Fortin-like barometers (provided with a floating gauge instead of the
25 ivory pin).

**A collection of
sub-daily
observations**

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2.1 Cistern's level correction for fixed-cistern barometers

The level l read on the scale of a fixed-cistern barometer is underestimated for high values ($l > l_0$, where l_0 is the zero-level, i.e. the level where no correction is needed) and overestimated for low values ($l < l_0$), due to the change of level in the cistern.

Therefore the following correction formula must be applied to the raw observations:

$$L = l + \frac{d^2}{D^2 - d^2}(l - l_0) \quad (1)$$

where L is the corrected level, d is the inside diameter of the tube and D that of the cistern (assuming a circular section).

For the large majority of the early instrumental records, d , D and l_0 are unknown. Even if we knew them, we cannot say for sure whether or not the correction was applied before recording the observations, or whether the correction was necessary at all. Most commercial barometers (including the ones intended for scientific use) were actually sold without the indication of l_0 (Middleton, 1964). In our metadata only in one case, for Cambridge (Harvard College), the observer clearly stated that “the barometer is provided with a floating gauge and scale of correction”.

We can try to quantify the maximum error that can arise from uncorrected observations. One case where the cistern's level correction could be applied in the literature is for the series of Stockholm: Moberg et al. (2002) estimated a correction of 1 % to $(l - l_0)$, this means that even for extreme high or low pressure values the error is lower than 0.5 hPa. We can have another example using the metadata for the observatory in London (Cornes, 2008): for the barometer in use there at the time the correction would be even lower, since the cistern/tube ratio was slightly larger than in Stockholm. A similar ratio is found for the barometer in Zwanenburg (Geurts and van Engelen, 1992). We can expect smaller cisterns by some amateur observers, however the errors introduced by the missing corrections are unlikely to be larger than 1 hPa.

CPD

11, 1741–1794, 2015

**A collection of
sub-daily
observations**

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2.2 Capillarity and drifts

In all mercury barometers, but in particular in fixed-cistern and Fortin barometers, a too thin tube can lead to underestimations in the readings due to capillarity. This error becomes larger than 1 hPa for $d < 8$ mm (Camuffo et al., 2006). The barometers in Stockholm and London had a tube with an internal diameter of only 3 and 6 mm, respectively, therefore they were probably affected by a substantial error. Capillarity was indeed the largest source of error in barometers and could be fully bypassed only in the second half of the 19th century with the adoption of reference primary barometers (Middleton, 1964). Nevertheless, correction tables had been around since at least 1776 (Cavendish, 1776), although their use is never mentioned in the metadata in our possession. The error introduced by capillarity can be assumed constant over a period of few years, with the exception of siphon barometers, in which the tube is exposed to air (and thus to humidity and dust).

The scale was often prone to physical changes, like mechanical drifts or periodical changes due to thermal expansion or to the humidity's effect on the wood of the support. The latter was estimated in Moberg et al. (2002) as negligible, however it depends on the individual instrument. Other significant errors and drifts can arise from the quality of the mercury or from bubbles of air that enters the tube. In general most barometers had probably a drift of some kind and were less reliable after a few decades of use.

2.3 Data processing

In this section we describe the procedure that was necessary to transform the raw data to a common consistent format that we could use for the analysis. After the conversion of all variables to metric units and of the observation times to the standard UTC, we corrected the pressure observations for temperature and local gravity, and we reduced them to mean sea level. We followed, when appropriate, the directives of the World Meteorological Organization (WMO, 2008). At the end of the procedure we interpolated the observations to regular 6 hourly time steps, in order to have simultaneous values.

CPD

11, 1741–1794, 2015

**A collection of
sub-daily
observations**

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3.1 Unit conversion

In 1815 only France had officially adopted the metric system, elsewhere metric units were rarely used. The English inch (= 25.40 mm) was the standard length unit in the English-speaking world. In the rest of the world, the most common unit for barometer scales was the Paris inch (= 27.07 mm). We encountered three other non-metric units, which were used only in specific countries: the Swedish inch (= 29.69 mm) in Sweden, the Vienna inch (= 26.34 mm) in Austria and the Rijnland inch (= 26.15 mm) in the Netherlands. The English and the Swedish inch had decimal sub-units (the resolution was usually one hundredth of an inch), the others were divided in 12 “lines”, which were in turn divided in 4 to 16 “points”.

The temperature was measured using either the Fahrenheit or the Réaumur scale. The only exceptions were in France and in Sweden, where the Celsius scale had already been adopted. We converted all temperature observations to °C.

2.3.2 Observation times

Observation times are available in various formats in the original records. Usually the observations were fixed at specific hours, but for some series they were indicated only qualitatively (e.g., “morning”) and in some others one of the observations was made at sunrise or sunset, whose time varies during the year. In 1815 all the countries of the observatories in the collection had already adopted the Gregorian calendar.

We assumed all times to refer to local solar time, since official standardised times did not exist. For qualitative observation times, we applied the following fixed conversions when we did not have any information from the available metadata: morning = 8 a.m.; noon = 12 p.m.; afternoon = 4 p.m.; evening = 8 p.m. However, when quantitative observation times are indicated only at the beginning of a manuscript (e.g., only on the first page of a meteorological register), we assume that they hold for the whole manuscript or in the whole series of manuscripts (e.g., if there is one volume per year and quantitative observation times are indicated only for the first year).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In cases for which observation times are noted as “sunrise” and “sunset”, the local sunrise and sunset is computed based on the date and latitude of the station using the following equation:

$$H_{\text{sun}} = \arccos(-\tan \phi \cdot \tan \delta) \cdot \frac{24}{2\pi} \quad (2)$$

where H_{sun} is the half-day length in hours, ϕ the latitude of the station and δ the declination of the sun, computed applying the algorithms described in Meeus (1999). The local sunrise (sunset) time is $0.5H_{\text{sun}}$ before (after) local noon.

If observation times for single observations are missing, but observations have been taken at regular intervals, we replaced the missing observation times with the most frequent observation time for this interval (e.g., 9 p.m. for evening observations if 9 p.m. is the most frequent known time for evening observations at one specific observatory). If the observation schedule was not regular, we assumed a local observation time of 2 p.m.

We finally translated local observation dates and times to UTC. For this we used a simple equation based on the longitude of the station:

$$t_{\text{UTC}} = t_{\text{loc}} - \lambda \cdot \frac{24}{360} \quad (3)$$

where λ is the longitude of the station in degrees east, t_{loc} is the local time and t_{UTC} is the UTC time.

2.3.3 Reduction to 0 °C

About half of the observatories in our data set used to record the temperature of the barometer. It was in fact common to have a mercury thermometer fixed on the same support of the barometer. Since the mercury expands and shrinks depending on the temperature, observations made with a mercury barometer must be corrected accordingly:

$$L_0 = (1 - \gamma T)L_{\text{mm}} \quad (4)$$

A collection of sub-daily observations

Y. Brugnara et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



temperature for the pressure reduction). We obtained similar results using observations from a certain part of the day (e.g., only morning or afternoon observations); in particular, the average errors in summer always remain within ± 1 hPa (not shown). Climatologies from the reanalysis introduce errors similar to those introduced by outside temperatures, slightly larger when the barometer is not in a heated room. In Sect. 3.3 we try to correct these errors using a statistical method.

Temperature has in general a vertical gradient along the barometer, meaning that the observed temperature of the barometer is actually the temperature of only one part of it (depending on where the thermometer is attached). Therefore the correction can introduce errors of the order of some tenths of hPa even when the temperature of the barometer is available.

The variance is more strongly affected when using climatologies (Fig. 3c and d). Using outside temperature introduces a random error in the variance that does not depend on the season and is usually smaller than 5 % for all stations but one: in Natchez (Mississippi) there is a systematic overestimation of the variance of about 10 %, which could depend on the sub-tropical climate of this station (i.e., a smaller pressure variance than any other station in the collection) or simply on the quality of the temperature observations (e.g., unshielded thermometer). Climatologies introduce a seasonal cycle in the variance error for some stations, with an underestimation (overestimation) of the variance in winter (summer).

We did not apply corrections for the thermal expansion of other parts of the barometer (cistern, tube, scale), which are usually one order of magnitude smaller than the correction for mercury and depend on the material used to build the barometer.

We also used Eq. (4) to rebase to 0°C pressure observations that had been reduced to some other temperature at the time of the readings. This results in a small inconsistency, because the correction tables in use at the time were purely empirical, being γ not known with sufficient precision. Therefore the original corrections do not correspond exactly to those resulting from Eq. (4).

CPD

11, 1741–1794, 2015

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The series of Milan, Salem, Stockholm and Turin had already been reduced to 0 °C in precedent works by data contributors (see the respective references for more details). In Exeter, the observer started to register the temperature at the barometer only in 1817; outside air temperature was used before that year (absolute differences between temperature at the barometer and outside temperature were on average smaller than 2.5 K during 1817).

2.3.4 Conversion to pressure units and correction for local gravity

The conversion of pressure readings from mm to hPa follows from the hydrostatic equation:

$$P_n = \rho g_{\varphi,h} L_0 \times 10^{-5} \quad (5)$$

where P_n is the absolute pressure in hPa reduced to normal gravity, $\rho = 1.35951 \times 10^4 \text{ kg m}^{-3}$ is the density of mercury at 0 °C, $g_{\varphi,h}$ is the local gravity (see below) and L_0 is the barometric reading in mm (corrected for temperature). This is equivalent to the usual procedure of first converting pressure readings from mm to hPa by using normal gravity acceleration in Eq. (5) and then correcting for local gravity by using:

$$P_n = \frac{g_{\varphi,h}}{g_n} P_0 \quad (6)$$

where P_0 is the absolute pressure not reduced to normal gravity and $g_n = 9.80665 \text{ m s}^{-2}$ is the normal gravity acceleration.

We estimated the local gravity $g_{\varphi,h}$ from the latitude φ and elevation h (in m a.s.l.), assuming flat terrain around the station (see WMO, 2008):

$$g_{\varphi,h} = [9.80620 \cdot (1 - 0.0026442 \cdot \cos 2\varphi - 0.0000058 \cdot \cos^2 2\varphi) - 0.000003086 \cdot h] \text{ ms}^{-2}. \quad (7)$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3.5 Reduction to mean sea level

To use the pressure observations for synoptic analysis, we reduced P to sea level:

$$P_0 = P \cdot \exp\left(\frac{\frac{g_{\varphi,h}}{R} \cdot h}{T_S + a \cdot \frac{h}{2}}\right) \quad (8)$$

where $R = 287.05 \text{ J kg}^{-1} \text{ K}^{-1}$ is the gas constant for dry air, $a = 6.5 \times 10^{-3} \text{ K m}^{-1}$ is the standard lapse rate of the fictitious air column below the station, and T_S is the outside temperature at the station in K.

We did not apply further corrections described in WMO (2008), since the uncertainty in our dataset is much higher than that required for modern barometers (i.e., $\pm 0.1 \text{ hPa}$).

Similarly to the reduction of pressure readings to 0°C , we used in-situ air temperature observations where available and resorted to climatological temperatures from the 20th Century Reanalysis (1871–1900) otherwise. We did not use temperature of the barometer to reduce pressure readings to sea level.

The series of Stockholm and Turin had already been reduced to sea level by the respective data contributors.

2.3.6 Quality control

We inspected visually each sea level pressure (SLP) series (and differences with nearby stations) to flag erroneous outliers and clear inhomogeneities in the period 1815–1817. Nearly all outliers derive from mistakes in the digitisation or in the transcriptions by the observer. When possible (i.e., when the original sources were readily available) we corrected them, otherwise we flagged them as erroneous and excluded them from the analysis.

The total number of pressure observations flagged after the quality control is 4657, corresponding to 4.1 % of the 1815–1817 dataset. Most of the flagged observations correspond to long periods in a few series where we detected large inhomogeneities:

Madrid (whole year 1815 flagged), New Haven (most of autumn/winter 1815/16), Valencia (all summer observations), Växjö (whole 1817) and Ylitornio (11 months in 1817). The number of flagged observations for each series is indicated in Table 1.

2.3.7 Interpolation on regular time steps

Another requirement for a synoptic analysis is that observations must be simultaneous. To achieve this, we linearly interpolated all pressure observations to four daily equally-spaced time steps: 00:00, 06:00, 12:00 and 18:00 UTC. If no observations of a certain station were available within ± 6 h from a certain time step, then we did not interpolate and considered the station to have no data for that specific time step. In Europe (where our analysis will focus), most observations were made very close to 06:00, 12:00 and 18:00 UTC; interpolated values for 00:00 UTC are in general less reliable and will not be analysed.

We did not interpolate outside temperature observations, because of their strong daily cycle.

3 Analysis

3.1 The post-Tambora period in monthly datasets

We start the analysis with a brief overview of the circulation and temperature anomalies that characterized the period from 1815 to 1817 in Europe. For this, we exploit seasonal gridded SLP fields statistically reconstructed by Küttel et al. (2010) using station pressure series and ships' log-book information from the northern North Atlantic. We also use the monthly temperature series for Central Europe from Dobrovolný et al. (2010), based on 11 homogeneous temperature series of stations located in Southern Germany, Bohemia, Austria and Switzerland in AD 1760–2007 and on documentary index series in AD 1500–1759.

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A collection of
sub-daily
observations**

Y. Brugnara et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Figure 4 shows the monthly temperature anomalies in Central Europe with respect to the period 1801–1830. From June 1815 to December 1816, 17 months out of 19 had negative anomalies. However, the largest negative anomaly was registered in April 1817, the coldest April of the entire series (i.e., in more than 500 years). The summer (June to August) of 1816 was the coldest in the instrumental part of the series and the second coldest since AD 1500.

Figure 5 shows the maps of seasonal SLP anomalies in Europe, calculated from the reconstruction using the same reference period (1801–1830).

Winters following large tropical volcanic eruptions are often stormier and warmer than the average over Northern Europe, as well as drier over the Iberian Peninsula (Dawson et al., 1997; Fischer et al., 2007). This is caused by the increased meridional temperature gradient in the stratosphere produced by volcanic aerosols, which supports a more positive North Atlantic Oscillation (NAO) in the troposphere (e.g., Kirchner et al., 1999). The winter of 1815/16 did not follow this rule and was colder than usual in Central and Northern Europe, despite a mild period in January (Fig. 4; see also Trigo et al., 2009). SLP anomalies (Fig. 5) resemble in fact a weak negative NAO and are very similar to those reconstructed for the other seasons of 1816. On the contrary, the winter of 1816/17 had a strong positive NAO and brought substantial warm anomalies in Central Europe (Fig. 4).

The spring of 1817 was again much colder than climatology, but the SLP pattern was different than that of 1816. In Sect. 3.3 we describe more in detail this pattern and its effects on Central and Southern Europe.

3.2 Storminess

One of the advantages of daily pressure observations with respect to monthly data is the possibility to study variability on the time scales of the typical large-scale weather phenomena. In particular, the variance of bandpass filtered daily pressure observations (hereafter “storminess”) is related to the frequency of stormy weather caused by extra-tropical cyclones and is commonly used for storm track analysis (e.g., Blackmon et al.,

1977). In this section, we apply a 2–6 days bandpass Lanczos filter (Duchon, 1979) with a 31 day convolution vector, to analyse winter and summer storminess in 1815–1817 in Europe and north-eastern North America.

We use only interpolated SLP observations at 12:00 UTC, because this is the only time step available for every series. Furthermore we require at least 90% of the 12:00 UTC values to be available in a certain season to calculate the variance for that season. To analyse winters we apply the filter to the 120 day period from 15 November to 14 March (13 March in leap years), for summers to the period from 18 May to 14 September.

The storminess for the winters of 1815/16, 1816/17 and 1817/18 is shown as SD in the last three panels of Fig. 6, where instead of absolute values we plotted the anomalies from the 1981–2010 climatology of the closest grid point in the Twentieth Century Reanalysis (Fig. 6, top panel). This analysis also constitutes a useful tool to verify the quality of the data. It is particularly evident from the map of 1816/17 that one station in Spain (Valencia) is not reliable, having a too high variability, and probably also one in North America (New Haven), which seems to have a too low variability when compared to the neighbouring stations. The observations in these two stations were corrected for temperature using, respectively, in-situ outside temperature and climatologies from the reanalysis, but it is unlikely, as demonstrated by Fig. 3, that the correction is the main responsible for errors of this magnitude. For Valencia, however, a systematic error similar to that described in Sect. 2.3.3 for Natchez is a possible contributor to the overestimation of the variance.

The difference between the winters of 1815/16 and 1816/17, which is very clear when looking at mean SLP fields (Fig. 5), disappears for the variance. The storminess anomalies suggest an eastward shift of the storm track in both winters, since the variance in all stations in North America is reduced by about 20%, while it is increased by approximately the same amount in Scandinavia. The few stations available in 1817/18 are enough to see a very different situation in terms of storminess, with a reduction in Northern Europe and positive anomalies in Southern Europe and New England.

A collection of sub-daily observations

Y. Brugnara et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



**A collection of
sub-daily
observations**

Y. Brugnara et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

SLP has climatologically a much lower variability in summer (Fig. 4, top panel) and it is difficult to interpret the results in terms of storm track, since baroclinic instability is much reduced. The summers of 1815 and 1816 (Fig. 4) show a quite similar pattern of variability in Europe, in particular a reduced storminess in Northern Europe. Perhaps surprisingly, the summer with the highest variability in Europe is that of 1817. In New England, the storminess of 1816 is similar to that of 1817. Besides, the maps confirm the suspect of a too low variability in the series of New Haven.

3.3 Synoptic analysis for three case studies in Europe

3.3.1 Statistical correction

Even though the results of the previous section demonstrate a good consistency among most of the series for what concerns variability, the lack of metadata for many of them causes large systematic errors in the mean values. A statistical approach is the only viable option to obtain absolute SLP values accurate enough for a synoptic analysis; namely, we use the reconstruction by Küttel et al. (2010) as reference to correct the land series in Europe.

It is important to mention that the reconstruction is not independent, in fact the monthly means of sixteen series in our collection were used as input for the reconstruction. However they were all homogenised by Küttel et al. (2010), therefore we are confident that the reconstruction offers the best possible estimation of mean SLP and that the application of the corrections guarantees a better reliability of synoptic weather maps.

Using the original SLP observations (not the interpolated values), we calculated seasonal means for each series in the period 1815–1817 and then applied a constant offset necessary to match the 1815–1817 seasonal means of the nearest grid point in the reconstruction. This was possible only if enough data were available: we calculated the offset using only the years with at least 90 % of the days in the target season having at least one observation available. When a series does not have enough data in any year

**A collection of
sub-daily
observations**

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for a certain season, we applied the average of the offsets from the available seasons. If the data is insufficient in every season, the series is not used in this section. This was the case for Derby, which has only one month of data. Moreover, we excluded the series of Valencia, which showed low reliability in the previous section. We did not correct the already homogenised series of Milan and Stockholm.

Since the reconstruction is based on monthly means, in turn calculated from daily means, we must apply a further correction to the offsets to take into account that our data are instantaneous observations rather than daily means. For this, we estimated the mean daily cycle of SLP for each season from the 1981–2010 climatology of the closest grid point of the MERRA reanalysis (Rienecker et al., 2011). MERRA offers the advantage of an hourly resolution, as well as a higher spatial resolution ($\frac{1}{2}^\circ$ latitude \times $\frac{2}{3}^\circ$ longitude) than the Twentieth Century Reanalysis. For stations with variable observation times, we used for the calculation the observation times (rounded to the hour) adopted most frequently at the target station in the target season. The resulting corrections are very small for all series, smaller than 1 hPa even for stations with only one observation per day.

On the other hand, the total statistical corrections are in some cases larger than 10 hPa (Fig. 8), while their root mean square is 4.9 hPa. The average correction (thick line in Fig. 8) has a seasonal cycle with a peak-to-peak amplitude of 1.9 hPa, indicating overestimated values in winter relatively to summer, nothing more than what we expected because of wrong temperature corrections for barometers in heated rooms (Fig. 3a and b).

Another important source of systematic errors is the uncertainty of the barometer elevation: considering a standard atmosphere near the sea level, an uncertainty of 20 m (which applies to most stations) results in an uncertainty in SLP of about 2.5 hPa. Moreover, the statistical correction can also take into account capillarity (see Sect. 2.2.2), which is probably the reason why the majority (about two thirds) of the applied offsets are positive (capillarity always causes an underestimation in mercury barometers).

the ground. According to Howard, this was the lowest value measured in London since 1797.

According to our dataset, temperatures were particularly low in Sweden, reaching -38.5°C in Umeå and -37°C in Härnösand, while the absolute minimum was measured in Ylitornio (Finland) with -40°C (not shown). The homogenised series of Stockholm allows a comparison with modern data: in the period 1981–2010 there has been only one cold spell (in January 1987) that was more severe than that of February 1816. However, cold spells of this magnitude were probably not so rare in the early instrumental period. In fact, Moberg et al. (2002) and Bergström and Moberg (2002) found a particularly high frequency of very cold winter days in Stockholm and Uppsala in the late 18th and early 19th century, although they warned that the data might still be affected by inhomogeneities. Daily temperatures lower than those of February 1816 were registered for example in January 1814 and again in December 1817. Similar results can be obtained by analysing the temperature series of St. Petersburg in north-western Russia (Jones and Lister, 2002).

3.3.3 Summer 1816

We analyse here one case at the beginning of July, one of the coldest periods of summer 1816 in Central Europe (e.g., about 7°C colder than usual in Geneva, see Auchmann et al., 2012). Abundant rainfall fell in the Alps where, in the night between the 3th and the 4th, a huge landslide, about 300 m wide, killed at least 14 people near the town of Uznach in eastern Switzerland (Erdrutsch in der Au (Goldingertal): Situationsplan, State Archives St. Gallen, Ref. KPG 1/65.1, 7/1816).

A shallow low-pressure system crosses the Alps between 1 and 2 July (Fig. 10), which is consistent with cold and rainy weather, and then probably moves to south-eastern Europe on the 3rd, when easterly winds are observed in the stations in eastern Europe (not shown). Afterwards, the Alps remained under the influence of unstable air coming from the Atlantic for several days. The weather diary kept for Aarau, in northern

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Switzerland, reports precipitation every day until the 19 July, always accompanied by westerly winds except for one day.

An area of high pressure is present over Scandinavia during the whole period shown in Fig. 10, suggesting fair weather there (confirmed by temperatures). In particular, on the north-eastern corner the maps show the temperature registered in Ylitornio at 2 p.m. LT, which are remarkably high for that latitude (the maximum temperature is reached on the 5 July at 31 °C, not shown). The quality of the measurements is questionable (see Klingbjer and Moberg, 2003), however it is interesting to point out that the average 2 p.m. temperature measured in Ylitornio in summer 1816 was 19 °C and that the absolute maximum in the entire series (1800–1825) is 34 °C. Therefore our data suggest the occurrence of an heatwave in north-eastern Europe in conjunction with the cold period in Central Europe. Again, the daily temperature series of St. Petersburg supports our conclusion, indicating 13 consecutive days (6–18 July) with mean daily temperature > 20 °C, the longest such series in the years 1815–1817. The month of July as a whole had nevertheless slightly negative temperature anomalies in that region (e.g., Luterbacher and Pfister, 2015), showing how even relatively long-lasting events can be overlooked by monthly means.

3.3.4 April 1817

After a relatively mild winter, the spring of 1817 struck a big final blow to Europe. In particular, as already mentioned, the month of April was extremely cold (see Fig. 4).

To gather more information on the most important weather events that distinguished this month, we did some research in contemporary newspapers and other historical sources. The most badly affected area was probably the northern slope of the Alps. Exceptional snowfalls and avalanches were often reported in that month especially in Austria: in Innsbruck (574 m a.s.l.), for instance, the snow fell in 18 days out of 30 (Fliri, 1998), while over two metres of snow were reported in Annaberg (976 m), near Vienna, after 16 consecutive days with snowfall (Lemberger Zeitung, 9 May 1817). At Buchlovice (234 m, south-east Moravia) a priest, Šimon Hausner, recorded snowfall on

**A collection of
sub-daily
observations**

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A collection of
sub-daily
observations**

Y. Brugnara et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the approach of the gale; so that there was scarcely time to take in the sails. In a few minutes we lost sight of the Augusta, which hitherto had kept at a small distance from us. A thick fog enveloped our ship; a cold rain, mixed with hailstones, which the storm furiously drove before it, covered the deck with pieces of ice of considerable size, and almost froze the crew. The ship was tossed violently; the yards and tackle were torn and broken: the waves rushed through the window into the forecastle, partly filled the hold with water; and at last, when the storm was at its height, the bowsprit broke short off. The hurricane raged with the utmost fury till noon, when the sea grew calmer, and the bleak Bora being succeeded by a mild east wind, we cast anchor in the middle of the sea, about three miles to the west of Rovigno.”

As suggested by the two German scientists, who demonstrate a remarkable knowledge of climatology, the storm was related to a severe bora wind event (Yoshino, 1976). The event was also felt in most of the Po valley (Northern Italy), where four of the stations in our dataset are located. The observatories of Padua and Bologna, which are close to the Adriatic coast, reported on that day thunderstorms, very strong wind from the north-east and snow flakes. Newspapers reported heavy snowfall in the eastern Alps during the same event, in particular about 50 cm of snow were measured in northern Slovenia and 10 cm in the city of Klagenfurt (Gazzetta di Milano, 8 May 1817).

In Fig. 12 we show the synoptic maps for 10 and 11 April. The position of the Austrian frigates in the morning of the 11th is marked by a star in the third map. Again, we are dealing with a blocking pattern, characterized by high pressure over north-western Europe and low pressure over Scandinavia. This configuration represents the negative phase of the so-called Scandinavian Pattern (e.g., Rogers, 1990), which is normally completed by a third pole (in this case, an area of high pressure) over central Siberia. This pattern stayed in place for most of the month of April 1817, continuously pumping Arctic cold air towards Central and Southern Europe. Enzi et al. (2014) singled out this

configuration as the most common responsible for widespread exceptional snowfall in Italy.

A cold air outbreak is the condition necessary for a severe Adriatic bora storm, and the synoptic pattern of 11 April (third panel in Fig. 12) is in fact a typical pattern for severe bora events (Jurčec, 1989). The maps show clearly the build-up of a pressure gradient between the northern and the southern slope of the Alps caused by the interaction of the cold air with the orographic barrier. It is likely that an orographically induced cyclone formed in the Mediterranean as a typical “cut-off” and that it stationed there for several days (e.g., Tibaldi and Buzzi, 1983). Spix and Martius (1824) wrote that when the *Austria* was near the coast of Southern Italy on the 22th, they could see the promontory of the Gargano, which reaches a maximum elevation of 1065 m, “covered with snow very low down”. They also repeatedly reported of stormy weather in the following days, culminating on the 27th in another violent storm that forced the frigate to seek shelter in Malta. They wrote:

“On the following morning we were already forty-two leagues to the west of Malta, when the wind suddenly settled in the N.N.W. It soon increased, and the waves ran so high, that it was impossible to keep the course to the south-west. The frigate rolled so violently, that in a short time the tackling was materially damaged; every thing movable was thrown backwards and forwards; and it seemed dangerous longer to expose the ship to the fury of the waves.”

The direction of the wind suggests that a cyclone was centred close to eastern Sicily. Note that the position reported in the official translation is probably affected by a mistake in the unit conversion: the original German version reports “vierzig Seemeilen”, literally 40 nautical miles (74 km). The word Seemeile was however commonly used also to indicate the “league” (i.e., 3 nautical miles), although it is not clear why the translator converted it to 42 leagues instead of 40. We think anyhow that the literal translation gives a more realistic position for the ship, which would have been only a few kilometres from the coast of Africa otherwise.

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A collection of
sub-daily
observations**

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



was necessary to produce reliable absolute sea level pressure values, because meta-
data are usually insufficient for this purpose. An analysis of the data in the period 1815–
1817 revealed a realistic and spatially consistent behaviour of the corrected pressure
observations, both concerning their variability and their absolute values. We found only
a small fraction of the 49 land series to have evident problems in terms of data quality
in at least part of the period.

Direct applications of this dataset are limited by its low spatial coverage. For instance,
in Europe the lack of data in key regions like the North Atlantic and the Mediterranean
prevents a complete analysis of synoptic patterns, although we could still give several
interesting insights on some of the most important large-scale meteorological events of
1815–1817. We also showed that documentary sources can give valuable assistance.

On the other hand the data can be a useful boundary condition for numerical weather
model simulations in order to produce complete four-dimensional reanalysis. In this
view the dataset will be made available in its raw (uncorrected) format in the Interna-
tional Surface Pressure Databank (Yin et al., 2008) and will be assimilated in a reanal-
ysis using the model of the Twentieth Century Reanalysis (Compo et al., 2011).

This paper gives also a picture of the quantity of meteorological observations avail-
able in the early instrumental period, the majority of which has not been digitised yet,
and of their potential for climate research. In particular, future initiatives aimed at the
recovery of historical records in the North Atlantic and the Mediterranean would give an
important contribution to our understanding of the climatic changes occurred in Europe
during the 18th and the 19th century.

**The Supplement related to this article is available online at
doi:10.5194/cpd-11-1741-2015-supplement.**

Acknowledgements. This work was supported by the Atmospheric Circulation Reconstructions
over the Earth (ACRE) initiative (www.met-acre.org), the Swiss National Science Foundation
(SNF) Sinergia project FUPSOL-II (Grant CRSII2-147659) and the EU Horizon 2020 EUSTACE

project (Grant Agreement no 640171). Renate Auchmann was supported by the SNF project TWIST (200021_146599 / 1), Rob Allan by the EU FP7 ERA-CLIM2 (European Reanalysis of Global Climate Observations 2) project and the Met Office Hadley Centre Climate Program (HCCP), Rudolf Brázdil by the Grant Agency of the Czech Republic for the project no. P209/11/0956 and Ladislava Řezníčková by the Ministry of Education, Youth and Sports of the CR within the National Sustainability Program I (NPU I), grant no. L01415. Fernando Domínguez-Castro was partially supported by the Prometeo Project, Secretariat of Higher Learning, Science, Technology and Innovation. Support for the Twentieth Century Reanalysis Project dataset is provided by the U.S. Department of Energy, Office of Science Innovative and Novel Computational Impact on Theory and Experiment (DOE INCITE) program, and Office of Biological and Environmental Research (BER), and by the National Oceanic and Atmospheric Administration Climate Program Office. We are grateful to the many who helped in the collection of manuscripts and metadata, including the personnel of the archives and libraries that we visited or contacted. Very appreciated were in particular the contributions of James P. Bowen, William Brown, Michele Brunetti, Dario Camuffo, Martín Jacques-Coper, A. José Leonardo, Matthias Röthlisberger, Arturo Sanchez-Lorenzo, Alexander Stickler and Clive Wilkinson. 20th Century Reanalysis data were provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA, from their Web site at <http://www.esrl.noaa.gov/psd/>. MERRA data were provided by the NASA Global Modeling and Assimilation Office (GMAO) and the NASA GES DISC.

References

Abel, C.: Narrative of a Journey in the Interior of China, Longman and Co., London, 1818. 1781
Ansell, T. J., Jones, P. D., Allan, R. J., Lister, D., Parker, D. E., Brunet, M., Moberg, A., Jacobeit, J., Brohan, P., Rayner, N. A., Aguilar, E., Alexandersson, H., Barriendos, M., Brandsma, T., Cox, N. J., Della-Marta, P. M., Drebs, A., Founda, D., Gerstengarbe, F., Hickey, K., Jónsson, T., Luterbacher, J., Nordli, O., Oesterle, H., Petrakis, M., Philipp, A., Rodwell, M. J., Saladie, O., Sigro, J., Slonosky, V., Srnc, L., Swail, V García-Suárez, A. M., Tuomenvirta, H., Wang, X., Wanner, H., Werner, P., Wheeler, D., and Xoplaki, E.: Daily mean sea level pressure reconstructions for the European-North Atlantic region for the period 1850–2003, *J. Climate*, 19, 2717–2742, 2006. 1745

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Auchmann, R., Brönnimann, S., Breda, L., Bühler, M., Spadin, R., and Stickler, A.: Extreme climate, not extreme weather: the summer of 1816 in Geneva, Switzerland, *Clim. Past*, 8, 325–335, doi:10.5194/cp-8-325-2012, 2012. 1766
- Bergström, H. and Moberg, A.: Daily air temperature and pressure series for Uppsala (1722–1998), *Climatic Change*, 53, 213–252, 2002. 1766
- Blackmon, M. L., Wallace, J. M., Lau, N.-C., and Mullen, S. L.: An observational study of the Northern Hemisphere wintertime circulation, *J. Atmos. Sci.*, 34, 1040–1053, 1977. 1761
- Böhm, R., Jones, P. D., Hiebl, J., Frank, D., Brunetti, M., and Maugeri, M.: The early instrumental warm-bias: a solution for long central European temperature series 1760–2007, *Climatic Change*, 101, 41–67, 2010. 1746, 1747
- Brázdil, R., Kiss, A., Luterbacher, J., and Valášek, H.: Weather patterns in eastern Slovakia 1717–1730, based on records from the Breslau meteorological network, *Int. J. Climatol.*, 28, 1639–1651, 2008. 1745
- Briffa, K. R., Jones, P. D., Schweingruber, F. H., and Osborn, T. J.: Influence of volcanic eruptions on Northern Hemisphere summer temperature over the past 600 years, *Nature*, 393, 450–455, 1998. 1746
- Brönnimann, S., Annis, J., Dann, W., Ewen, T., Grant, A. N., Griesser, T., Krähenmann, S., Mohr, C., Scherer, M., and Vogler, C.: A guide for digitising manuscript climate data, *Clim. Past*, 2, 137–144, doi:10.5194/cp-2-137-2006, 2006. 1746
- Camuffo, D.: Calibration and instrumental errors in early measurements of air temperature, *Climatic Change*, 53, 297–329, 2002a. 1746
- Camuffo, D.: Errors in early temperature series arising from changes in style of measuring time, sampling schedule and number of observations, *Climatic Change*, 53, 331–352, 2002b. 1746
- Camuffo, D.: History of the long series of daily air temperature in Padova (1725–1998), *Climatic Change*, 53, 7–75, 2002c. 1747
- Camuffo, D. and Jones, P. (Eds.): *Improved Understanding of Past Climatic Variability from Early Daily European Instrumental Sources*, Springer, Dordrecht, 2002. 1745
- Camuffo, D., Cocheo, C., and Sturaro, G.: Corrections of systematic errors, data homogenisation and climatic analysis of the Padova pressure series (1725–1999), *Climatic Change*, 78, 493–514, 2006. 1747, 1751, 1753
- Cavendish, H.: An account of the meteorological instruments used at the Royal Society's House, *Phil. Trans.*, 66, 375–401, 1776. 1753

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Chenoweth, M.: Ships' logbooks and "The Year Without a Summer", *B. Am. Meteorol. Soc.*, 77, 2077–2094, 1996. 1746, 1748
- Compo, G. P., Whitaker, J. S., and Sardeshmukh, P. D.: Feasibility of a 100-year reanalysis using only surface pressure data, *B. Am. Meteorol. Soc.*, 87, 175–190, 2006. 1745
- 5 Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Matsui, N., Allan, R. J., Yin, X., Gleason, B. E., Vose, R. S., Rutledge, G., Bessemoulin, P., Brönnimann, S., Brunet, M., Crouthamel, R. I., Grant, A. N., Groisman, P. Y., Jones, P. D., Kruk, M. C., Kruger, A. C., Marshall, G. J., Maugeri, M., Mok, H. Y., Nordli, Ø., Ross, T. F., Trigo, R. M., Wang, X. L., Woodruff, S. D., and Worley, S. J.: The twentieth century reanalysis project, *Q. J. Roy. Meteor. Soc.*, 137, 1–28, 2011. 1745, 1756, 1772
- 10 Cornes, R.: The barometer measurements of the Royal Society of London: 1774–1842, *Weather*, 63, 230–235, 2008. 1752
- Cornes, R.: Historic storms of the northeast Atlantic since circa 1700: a brief review of recent research, *Weather*, 69, 121–125, 2014. 1745
- 15 Cornes, R. C., Jones, P. D., Briffa, K. R., and Osborn, T. J.: A daily series of mean sea-level pressure for Paris, 1670–2007, *Int. J. Climatol.*, 32, 1135–1150, 2012. 1749
- Davy, J.: Observations on the temperature of the ocean and atmosphere, and on the density of sea-water, made during a voyage to Ceylon, *Philos. T. R. Soc. Lond.*, 107, 275–292, 1817. 1781
- 20 Dawson, A. G., Hickey, K., McKenna, J., and Foster, D.: A 200-year record of gale frequency, Edinburgh, Scotland: possible link with high-magnitude volcanic eruptions, *Holocene*, 7, 337–341, 1997. 1761
- Dobrovolný, P., Moberg, A., Brázdil, R., Pfister, C., Glaser, R., Wilson, R., van Engelen, A., Limanówka, D., Kiss, A., Halíčková, M., Macková, J., Riemann, D., Luterbacher, J., and Böhm, R.: Monthly, seasonal and annual temperature reconstructions for Central Europe derived from documentary evidence and instrumental records since AD 1500, *Climatic Change*, 101, 69–107, 2010. 1760, 1785
- 25 Domínguez-Castro, F., Ribera, P., García-Herrera, R., Vaquero, J. M., Barriendos, M., Cuadrat, J. M., and Moreno, J. M.: Assessing extreme droughts in Spain during 1750–1850 from rogation ceremonies, *Clim. Past*, 8, 705–722, doi:10.5194/cp-8-705-2012, 2012. 1746
- 30 Duchon, C. E.: Lanczos filtering in one and two dimensions, *J. Appl. Meteorol.*, 18, 1016–1022, 1979. 1762

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Enzi, S., Bertolin, C., and Diodato, N.: Snowfall time-series reconstruction in Italy over the last 300 years, *Holocene*, 24, 346–356, 2014. 1769
- Fischer, E., Luterbacher, J., Zorita, E., Tett, S., Casty, C., and Wanner, H.: European climate response to tropical volcanic eruptions over the last half millennium, *Geophys. Res. Lett.*, 34, L05707, doi:10.1029/2006GL027992, 2007. 1746, 1761
- Fliri, F.: *Naturchronik von Tirol*, Wagner, Innsbruck, 1998. 1767, 1771
- Geurts, H. A. M. and van Engelen, A. F. V.: *Beschrijving antieke meetreeksen*, Koninklijk Nederlands Meteorologisch Instituut 165-V, KNMI, De Bilt, 1992. 1752
- Hall, B.: *Account of a voyage of discovery to the west coast of Corea, and the great Loo-Choo island*, Murray, London, 1818. 1781
- Jelinek, C.: *Anleitung zur Anstellung meteorologischer Beobachtungen und Sammlung von Hilfstafeln*, K.k. Hof- und Staatsdruckerei, Vienna, 1869. 1783
- Jones, P. and Lister, D.: The daily temperature record for St. Petersburg (1743–1996), *Climatic Change*, 53, 253–267, 2002. 1766
- Jones, P. D., Davies, T. D., Lister, D. H., Slonosky, V., Jónsson, T., Bärring, L., Jónsson, P., Maheras, P., Kolyva-Machera, F., Barriendos, M., Martin-Vide, J., Rodriguez, R., Alcoforado, M. J., Wanner, H., Pfister, C., Luterbacher, J., Rickli, R., Schuepbach, E., Kaas, E., Schmith, T., Jacobeit, J., and Beck, C.: Monthly mean pressure reconstructions for Europe for the 1780–1995 period, *Int. J. Climatol.*, 19, 347–364, 1999. 1745
- Jurčec, V.: Severe Adriatic bora storms in relation to synoptic developments, *Hrvatski meteorološki časopis*, 24, 11–20, 1989. 1770
- Kandlbauer, J., Hopcroft, P. O., Valdes, P., and Sparks, R.: Climate and carbon cycle response to the 1815 Tambora volcanic eruption, *J. Geophys. Res.*, 118, 12–497, 2013. 1746
- Kington, J.: The Societas Meteorologica Palatina: an eighteenth-century meteorological society, *Weather*, 29, 416–426, 1974. 1744, 1749
- Kirchner, I., Stenchikov, G. L., Graf, H.-F., Robock, A., and Antuña, J. C.: Climate model simulation of winter warming and summer cooling following the 1991 Mount Pinatubo volcanic eruption, *J. Geophys. Res.*, 104, 19039–19055, 1999. 1761
- Klingbjör, P. and Moberg, A.: A composite monthly temperature record from Tornedalen in northern Sweden, 1802–2002, *Int. J. Climatol.*, 23, 1465–1494, 2003. 1767
- Küttel, M., Xoplaki, E., Gallego, D., Luterbacher, J., Garcia-Herrera, R., Allan, R., Barriendos, M., Jones, P., Wheeler, D., and Wanner, H.: The importance of ship log data: recon-

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- structuring North Atlantic, European and Mediterranean sea level pressure fields back to 1750, *Clim. Dynam.*, 34, 1115–1128, 2010. 1760, 1763, 1786
- Luterbacher, J. and Pfister, C.: The year without a summer, *Nat. Geosci.*, 8, 246–248, 2015. 1746, 1767
- 5 Mangianti, F. and Beltrano, M. C.: La neve a Roma dal 1741 al 1991, MIPAAF-UCEA, Rome, 1993. 1771
- Meeus, J. H.: *Astronomical Algorithms*, 2nd edn., Willmann-Bell, Richmond, VA, 1999. 1755
- Middleton, W. E. K.: *The History of the Barometer*, Johns Hopkins, Baltimore, MD, 1964. 1751, 1752, 1753
- 10 Middleton, W. E. K.: *The Experimenters: A Study of the Accademia del Cimento*, Johns Hopkins, Baltimore, MD, 1972. 1744
- Moberg, A.: *Meteorological observations in Sweden made before AD 1860*, *Paläoklimaforschung/PalaeoClim. Res.*, 23, 99–119, 1998. 1744, 1749
- Moberg, A., Bergström, H., Krigsman, J. R., and Svanered, O.: Daily air temperature and pressure series for Stockholm (1756–1998), *Climatic Change*, 53, 171–212, 2002. 1747, 1752, 1753, 1766
- 15 Oppenheimer, C.: Climatic, environmental and human consequences of the largest known historic eruption: Tambora volcano (Indonesia) 1815, *Prog. Phys. Geog.*, 27, 230–259, 2003. 1746
- 20 Pfister, C.: *Wetternachhersage*, Haupt, Bern, 1999. 1746, 1771
- Post, J. D.: *The Last Great Subsistence Crisis in the Western World*, Johns Hopkins University Press, Baltimore, MD, 1977. 1746
- Réaumur, R. A. F.: Règles pour construire des thermomètres dont les degrés sont comparables, in: *Mémoires de l'Academie Royale Des Sciences pour 1730*, 452–507, Durand, Paris, 1732. 25 1747
- Rex, D. F.: Blocking action in the middle troposphere and its effect upon regional climate, *Tellus*, 2, 275–301, 1950. 1765
- Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, Siegfried D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's modern-era retrospective analysis for research and applications, *J. Climate*, 24, 3624–3648, 2011. 1764
- 30

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Rogers, J. C.: Patterns of low-frequency monthly sea level pressure variability (1899–1986) and associated wave cyclone frequencies, *J. Climate*, 3, 1364–1379, 1990. 1769
- Spix, J. B. and Martius, C. F. P.: Travels in Brazil, in the Years 1817–1820, Vol. 1, Longman and Co., London, 1824. 1768, 1770
- 5 Stommel, H. and Stommel, E.: The year without a summer, *Sci. Am.*, 240, 176–186, 1979. 1746
- Stothers, R. B.: The great Tambora eruption in 1815 and its aftermath, *Science*, 224, 1191–1198, 1984. 1746
- Tibaldi, S. and Buzzi, A.: Effects of orography on Mediterranean lee cyclogenesis and its relationship to European blocking, *Tellus A*, 35, 269–286, 1983. 1770
- 10 Trigo, R. M., Vaquero, J. M., Alcoforado, M.-J., Barriendos, M., Tabora, J., García-Herrera, R., and Luterbacher, J.: Iberia in 1816, the year without a summer, *Int. J. Climatol.*, 29, 99–115, 2009. 1746, 1761
- Wagner, S. and Zorita, E.: The influence of volcanic, solar and CO₂ forcing on the temperatures in the Dalton Minimum (1790–1830): a model study, *Clim. Dynam.*, 25, 205–218, 2005. 1746
- 15 Wegmann, M., Brönnimann, S., Bhend, J., Franke, J., Folini, D., Wild, M., and Luterbacher, J.: Volcanic influence on European summer precipitation through monsoons: possible cause for “Years without Summer”, *J. Climate*, 27, 3683–3691, 2014. 1746
- Wetter, O., Pfister, C., Weingartner, R., Luterbacher, J., Reist, T., and Trösch, J.: The largest floods in the High Rhine basin since 1268 assessed from documentary and instrumental evidence, *Hydrolog. Sci. J.*, 56, 733–758, 2011. 1771
- 20 Winkler, P.: Hohenpeißenberg 1781–2006 – das älteste Bergobservatorium der Welt, Deutscher Wetterdienst, Offenbach am Main, 2006. 1749
- WMO: Guide to meteorological instruments and methods of observation, WMO-No. 8, World Meteorological Organization, Geneva, 2008. 1753, 1758, 1759
- 25 Yin, X., Gleason, B., Compo, G., Matsui, N., and Vose, R.: The International Surface Pressure Databank (ISPD) land component version 2.2, Tech. rep., National Climatic Data Center, Asheville, NC, available at: ftp://ftp.ncdc.noaa.gov/pub/data/ispd/doc/ISPD2_2.pdf (last access: 11 May 2015), 2008. 1772
- 30 Yoshino, M.: Local Wind Bora, University of Tokyo Press, Tokyo, 1976. 1769

A collection of sub-daily observations

Y. Brugnara et al.

Table 1. List of land stations included in the collection, in alphabetical order. Observatories managed directly by scientific organizations are written in bold. Metadata refer to the period 1815–1817. Legend of abbreviations: Lon = longitude in degrees East, Lat = latitude in degrees North, Elev = elevation of the barometer in m a.s.l. (rounded to the unit), Obs = typical number of pressure observations per day, Loc = exact location (within 100 m) from metadata, TCorr = data used for temperature correction, Tot = number of pressure observations in 1815–1817, Flag = number of flagged observations after quality control, Y = available, N = not available, TB = temperature of the barometer, TA = outside air temperature, CL = outside temperature climatology, CO = observations already corrected for temperature. A question mark indicates estimated elevations.

Name	Lon	Lat	Elev	Obs	Loc	TCorr	Years	Tot	Flag
Aarau	8.04	47.39	380?	2	N	CO	1815–1816	1431	1
Albany	−73.75	42.65	12?	3	Y	TA	1815	543	0
Althorp	−1.00	52.28	105?	2	Y	TA	1816–1817	1400	0
Armagh	−6.65	54.35	64	3	Y	TB	1796–1965	3286	0
Avignon	4.80	43.95	22	4	N	TB	1816	982	0
Barcelona	2.17	41.38	20?	3	Y	TA	1811–1820	3288	12
Barnnton	−3.29	55.96	50?	1	N	TA	1815–1817	968	5
Bologna	11.35	44.50	74	1	Y	TA	1815–1817	1088	0
Boston	−0.03	52.98	10?	1	N	TA	1816–1817	713	0
Brunswick	−69.96	43.91	25?	3	Y	CL	1815–1817	3112	0
Cádiz	−6.30	36.53	15?	3	N	TA	1816–1820	1461	0
Cambridge	−71.12	42.37	9	3	N	CO	1815–1816	818	0
Coimbra	−8.42	40.21	95?	4	Y	TB	1815–1817	3665	1
Cracow	19.96	50.06	212	3	Y	TB	1816	1098	19
Derby	−1.48	52.93	50?	2	N	TA	1817	64	0
Düsseldorf	6.77	51.23	35?	3	N	TA	1816–1817	1187	2
Edinburgh	−3.18	55.96	110?	2	Y	CO	1817	340	0
Exeter	−3.53	50.72	47?	3	Y	TB	1813–1817	3058	1
Gdańsk	18.65	54.35	14	3	Y	TB	1815–1817	3278	4
Geneva	6.15	46.23	405?	2	Y	CO	1796–1863	2129	0
Göteborg	11.97	57.71	15?	3	N	TB	1815–1817	3288	0
Haarlem	4.65	52.38	2	3	Y	TA	1801–1841	3288	6
Härnösand	17.94	62.63	15?	3	N	TA	1815–1816	2027	0
Hohenpeissenberg	11.02	47.80	995	3	Y	CO	1781–2009	3288	3
Karlsruhe	8.40	49.01	121	3	Y	TB	1815–1817	3288	3
London	−0.12	51.52	24	2	Y	TB	1815–1817	2192	76
Lviv	24.03	49.84	295?	3	Y	CO	1815–1817	2576	0
Madrid	−3.71	40.41	650?	3	N	TA	1814–1817	1488	1096

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



CPD

11, 1741–1794, 2015

A collection of
sub-daily
observations

Y. Brugnara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 1. Position of the land observatories (red dots) and routes of the ships. For the latter, filled symbols denote locations for which pressure data are available, colours indicate time from December 1815 (magenta) to March 1817 (blue).




Figure 2. Drawing of the cisterns of a Fortin barometer (left) and of a fixed-cistern barometer (right). In the Fortin barometer a screw (indicated by the letter “S”) allows to adjust the level of the mercury in the cistern. From Jelinek (1869).




Figure 3. (a) Monthly averages of the differences between temperature of the barometer and outside temperature for the stations where both are available. (b) Monthly averages of the differences between temperature of the barometer and temperature climatologies from the Twentieth Century Reanalysis for the stations where the temperature of the barometer is available. In both panels the corresponding error in the pressure reduction to 0 °C is shown on the right axis, calculated considering an uncorrected barometer reading of 760 mm. (c) Monthly ratios between the variance of pressure observations corrected using outside temperature and the same observations corrected using the temperature of the barometer, for the same stations as in (a). (d) Monthly ratios between the variance of pressure observations corrected using climatologies and the same observations corrected using the temperature of the barometer, for the same stations as in (b). All plots are based on the period 1815–1817, climatologies are calculated from the period 1871–1900.




Figure 4. Monthly temperature anomalies in Central Europe in the period 1815–1817 with respect to 1801–1830 (Dobrovolný et al., 2010).

A collection of sub-daily observations

Y. Brugnara et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A collection of
sub-daily
observations**

Y. Brugnara et al.




Figure 5. Seasonal SLP anomalies (in hPa) in Europe for winter (DJF), spring (MAM), summer (JJA) and autumn (SON) for the years 1815–1817 (reference period 1801–1830), reconstructed by Küttel et al. (2010).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**A collection of
sub-daily
observations**

Y. Brugnara et al.




Figure 6. SD of daily (12:00 UTC) bandpass filtered SLP (in hPa) in winter (as defined in the text). The top panel shows the 1981–2010 climatology in the Twentieth Century Reanalysis. The other panels show land observations for 1815/16, 1816/17 and 1817/18, in terms of anomalies from the top panel. Empty squares indicate stations with insufficient data (see text).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A collection of
sub-daily
observations**

Y. Brugnara et al.




Figure 7. Similar to Fig. 6, but for summer (as defined in the text).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Figure 8. Seasonal corrections applied for the case studies. Each color represents a different station in Europe, the thick black line is the average of all corrections.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 9. Synoptic maps for the two main cold spells in Central Europe during winter 1815/16. Coloured points represent SLP observations (in hPa). To facilitate interpretation, isobars at intervals of 5 hPa are drawn using inverse distance weights and the approximate position of pressure minima and maxima are indicated by the letters L and H, respectively. White numbers represent temperatures (in °C) observed within ± 1 h.

A collection of sub-daily observations

Y. Brugnara et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



A collection of sub-daily observations

Y. Brugnara et al.




Figure 10. Similar to Fig. 9, but showing maps for the first four days of July 1816 at 12:00 UTC. Note that the colour scale has changed and isobars are drawn at intervals of 2 hPa.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A collection of sub-daily observations

Y. Brugnara et al.




Figure 12. Similar to Fig. 9, but showing maps for 10–11 April 1817 at 12h interval. A star marks the position of the frigates *Austria* and *Augusta* in the morning of the 11th.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A collection of sub-daily observations

Y. Brugnara et al.




Figure 13. Similar to Fig. 9, but showing maps for 25–27 April 1817 at 12h interval. A star marks the position of the frigate *Austria* in the morning of the 27th.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

