



**A multi-resolution
assessment (CMAQ)**

K. W. Appel et al.

A multi-resolution assessment of the Community Multiscale Air Quality (CMAQ) Model v4.7 wet deposition estimates for 2002–2006

K. W. Appel¹, K. M. Foley¹, J. O. Bash¹, R. W. Pinder¹, R. L. Dennis¹, D. J. Allen²,
and K. Pickering³

¹Atmospheric Modelling and Analysis Division, National Exposure Research Laboratory,
Office of Research and Development, US Environmental Protection Agency, RTP, NC, USA
²Dept. of Atmospheric and Oceanic Science, University of Maryland, College Park, MD, USA
³Atmospheric Chemistry and Dynamics Branch, NASA-Goddard, Greenbelt, MD, USA

Received: 15 October 2010 – Accepted: 12 November 2010 – Published: 9 December 2010

Correspondence to: K. W. Appel (appel.wyat@epa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[I◀](#)

[▶I](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

This paper examines the operational performance of the Community Multiscale Air Quality (CMAQ) model simulations for 2002–2006 using both 36-km and 12-km horizontal grid spacing with a primary focus on the performance of the CMAQ model in predicting wet deposition of sulfate (SO_4^-), ammonium (NH_4^+) and nitrate (NO_3^-). Performance of the wet deposition species is determined by comparing CMAQ predicted concentrations to concentrations measured by the National Acid Deposition Program (NADP), specifically the National Trends Network (NTN). For SO_4^- wet deposition, the CMAQ model estimates were generally comparable between the 36-km and 12-km simulations for the eastern US, with the 12-km simulation giving slightly higher estimates of SO_4^- wet deposition than the 36-km simulation on average. The normalized mean bias (NMB) was slightly higher for the 12-km simulation, however, both simulations had annual biases that were less than $\pm 15\%$ for each of the five years. The model estimated SO_4^- wet deposition values improved when they were adjusted to account for biases in the model estimated precipitation. The CMAQ model underestimates NH_4^+ wet deposition over the eastern US using both the 36-km and 12-km horizontal grid spacing, with a slightly larger underestimation in the 36-km simulation. The largest underestimations occur during the winter and spring periods, while the summer and fall have slightly smaller underestimations of NH_4^+ wet deposition. Annually, the NMB generally ranges between -10% and -16% for the 12-km simulation and -12% to -18% for the 36-km simulation over the five-year period for the eastern US. The underestimation in NH_4^+ wet deposition is likely due, in part, to the poor temporal and spatial representation of ammonia (NH_3) emissions, particularly those emissions associated with fertilizer applications and NH_3 bi-directional exchange. The model performance for estimates of NO_3^- wet deposition are mixed throughout the year, with the model largely underestimating NO_3^- wet deposition in the spring and summer in the eastern US, while the model has a relatively small bias in the fall and winter. Model estimates of NO_3^- wet deposition tend to be slightly lower for the 36-km simulation as compared

GMDD

3, 2315–2360, 2010

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the 12-km simulation, particularly in the spring. Annually for the eastern US, the NMB ranges from roughly -12% to -20% for the 12-km simulation and -18% to -26% for the 36-km simulation. The underestimation of NO_3^- wet deposition in the spring and summer is due, in part, to a lack of lightning generated NO emissions in the upper troposphere, which can be a large source of NO in the spring and summer when lightning activity is the high. CMAQ model simulations that include the production of NO from lightning show a significant improvement in the NO_3^- wet deposition estimates in the eastern US in the summer. Model performance for the western US was generally not as good as that for the eastern US for all three wet deposition species.

1 Introduction

Atmospheric deposition of sulfur and nitrogen cause deleterious impacts on terrestrial and aquatic ecosystems due to acidification and excess nutrients (Lovett and Tear, 2008; Driscoll et al., 2001, 2003; Fenn et al., 2003). Sulfur deposition from SO_2 and SO_4^{2-} emissions contributes to acidification and nitrogen deposition from nitrogen oxide (NO_x) and ammonia (NH_3) emissions contribute to acidification and excess nitrogen nutrients. Estimates of wet and dry deposition of nitrogen and sulfur are needed for sensitive ecosystems, as total deposition estimates are used to assess whether current or projected pollutant levels exceed a point where significant harmful effects on sensitive elements of the environment are likely to occur (Geiser et al., 2010). Monitoring of wet deposition is relatively sparse and monitoring of dry deposition is extremely sparse, contributing to significant interpolation errors when these data are used to estimate deposition in unmonitored areas. Thus, a regional air quality model, like the Community Multiscale Air Quality (CMAQ; Byun and Schere, 2006) model, can be used to provide a more spatially complete estimate of total deposition to the sensitive ecosystems. However, the model estimates must first be evaluated to establish the credibility of the model in replicating the observed wet deposition.

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A multi-resolution assessment (CMAQ)

K. W. Appel et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Evaluating the ability of the air quality model to replicate observed net (wet + dry) deposition is difficult. The National Atmospheric Deposition Program (NADP) monitoring sites provide the most complete spatial coverage of observed wet deposition across the US on a temporal scale suitable for air quality model evaluations. Evaluation of dry deposition is even more challenging because monitoring network (e.g. Clean Air Status and Trends Network) dry deposition levels are based on modelled values of deposition velocity and, hence, are not a true measure of dry deposition. Therefore, this work focuses on wet deposition to provide a test of the ability of the model to mix, transport, transform and scavenge the pollutant emissions at the regional scale. Many sensitive ecosystems are in complex terrain where orographic effects influence the precipitation patterns and consequently wet deposition. Thus, quantifying precipitation biases as part of the wet deposition evaluation is critical.

This paper examines the performance of the CMAQ model sulfate (SO_4^-), nitrate (NO_3^-) and ammonium (NH_4^+) wet deposition estimates for the 2002–2006 period over the continental United States (CONUS) using two model grid-spacing options, namely 12-km and 36-km grid spacing. The performance of the CMAQ model estimates is examined temporally using various averaging periods (i.e., monthly, seasonal, annual and multi-annual) and spatially across different regions, as the model performance can vary significantly in space. In cases where deficiencies in model performance are identified, model improvements, such as the production of NO_x from lightning and the inclusion of bi-directional flux of NH_3 , are tested and their impacts on model performance assessed. Together, these analyses provide insight into the strengths and weaknesses of the CMAQ model in estimating wet deposition of sulfur and nitrogen to sensitive ecosystems.

2 Input data and model configuration

2.1 Meteorology

The CMAQ model requires gridded meteorological data to provide estimates of various meteorological parameters such as temperature, wind speed and direction, relative humidity and planetary boundary layer (PBL) height. The 5th generation Mesoscale Model (MM5; Grell et al., 1994) is a Eulerian meteorological model that provides estimates of the meteorological parameters required by the CMAQ model and has been used and tested extensively with the CMAQ model over the past 15 years. For this work, the MM5 version 3.7.4 was used for both the 36-km and 12-km simulations. The 36-km MM5 domain consists of 165 by 129 grid cells covering the entire CONU, and includes portions of Canada and Mexico. The 12-km domain consists of 290 by 251 grid cells covering the eastern two-thirds of the US, southern Canada and northern Mexico.

Boundary conditions for the 2002–2005 36-km and 12-km MM5 simulations were provided by the 40-km Eta Data Assimilation System (EDAS) data; while the 12-km North American Model (NAM) data were used as boundary conditions for the 2006 36-km and 12-km MM5 simulations, with any missing data filled in using the 32-km North American Regional Reanalysis data. The MM5 simulations utilized the Kain-Fritsch 2 (KF2) cumulus parameterization (Kain, 2004); the asymmetric convective model version 2 (ACM2) PBL scheme (Pleim, 2007a, b); the Reisner 2 explicit microphysics scheme (Reisner et al., 1998); the Dudhia shortwave radiation scheme (Dudhia, 1989); the RRTM longwave radiation scheme (Mlawer et al., 1997); and the Pleim-Xiu land surface model (LSM; PX; Xiu and Pleim, 2001; Pleim and Xiu, 1995). Both the 36-km and 12-km MM5 simulations utilized 34 vertical layers, with the surface layer set at approximately 36 metres. The meteorological outputs from both sets of MM5 simulations were processed to create model-ready inputs for CMAQ using the Meteorology-Chemistry Interface Processor (MCIP; Otte et al., 2005) version 3.4.

GMDD

3, 2315–2360, 2010

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[I◀](#)

[▶I](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



2.2 Emissions

The 2002 National Emissions Inventory (NEI) version 3 was used as the primary basis for the 2002–2006 emissions inputs. Version 3 of the 2002 NEI is documented at <http://www.epa.gov/ttn/chief/net/2002inventory.html#documentation>. For the major point sources, namely electric generating units (EGUs), year specific continuous emission monitoring systems (CEMS) data were used. Year specific updates to mobile emissions were done using the MOBILE6 model and daily estimates of fire emissions based on satellite detection of fires were included as well. NH₃ emissions from agricultural cropping practices in CMAQ are provided by a separate model based on the Carnegie Mellon University (CMU) ammonia emission model (Goebes et al., 2003), which are then combined with the NEI. Monthly NH₃ emissions from livestock were adjusted according to the inverse-modelling recommendations of Gilliland et al. (2006). For inventories outside of the US, which include Canada, Mexico and offshore emissions, the latest available base year inventories were used. The CMAQ model-ready emissions were created using the Sparse Matrix Operator Kernel Emissions (SMOKE) modelling system (Houyoux et al., 2000).

2.3 CMAQ model configuration

The CMAQ simulations were performed at the 36-km horizontal grid spacing for the CONUS, while for the eastern two-thirds of the US a CMAQ simulation using 12-km horizontal grid spacing was performed. Chemical boundary conditions for the 12-km simulation were provided by the 36-km simulation, while boundary conditions for the 36-km CMAQ simulation were provided by a non-year specific GEOS-Chem (Bey et al., 2001) simulation. The boundary data for the 36-km CMAQ simulation were created by taking the median value of a 2.0 degree by 2.5 degree (latitude-longitude) 24-vertical layer 2002 GEOS-Chem simulation and averaging the three-hourly data to monthly values. These monthly averages were then used as boundary conditions for all five years of the 36-km CMAQ model simulations.

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The air quality simulations utilized CMAQv4.7 (Foley et al., 2010), the latest version of the model available at that time. The simulations included a 10-day spin-up period for the 36-km simulations, while a 3-day spin-up period was used for the 12-km simulations. The CMAQ simulations were performed using the same horizontal dimensions as their respective meteorology simulation except that the horizontal dimensions were reduced by five grid cells on each of the four lateral boundaries to avoid artifacts that can appear along the domain boundaries in the meteorological simulations. However, unlike the meteorological simulations which utilized 34-vertical layers, the CMAQ simulations used 24-vertical layers. The CMAQ model simulations used the AERO5 aerosol module (Carlton et al., 2010), the Carbon-Bond 05 (CB05) chemical mechanism with chlorine chemistry extensions (Yarwood et al., 2005) and the ACM2 PBL scheme (Pleim, 2007a, b).

2.4 Assessing model performance

The assessment of the CMAQ model's wet deposition estimates is accomplished by comparing the simulated wet deposition estimates to observed wet deposition values available from the National Acid Deposition Program's (NADP; <http://nadp.sws.uiuc.edu>) National Trends Network (NTN). The NTN measures total weekly wet deposition of several atmospheric pollutants, including SO_4^- , NH_4^+ and NO_3^- . Since all of the SO_2 in rainwater is oxidized to SO_4^- by the time the samples are analysed for the NTN (high prevalence of oxidants), the CMAQ estimates of SO_4^- wet deposition include 150% of the model estimated SO_2 wet deposition to account for the SO_2 captured in the observations. Because in solution the favoured phase of NH_3 is NH_4^+ at the pH of rainwater, the CMAQ estimates of NH_4^+ wet deposition include 106% of the model estimated NH_3 wet deposition to account for the reduced nitrogen (both NH_4^+ and NH_3) captured in the NTN observations. Likewise, because in solution HNO_3 reacts with water and dissociates to NO_3^- as the favoured phase, the CMAQ estimates of NO_3^- wet deposition include 98.4% of the model estimated nitric acid wet deposition to account for NO_3^- captured as nitric acid and converted to NO_3^- in the NTN measurements.

The NTN consists of approximately 185 sites in the eastern US (east of 110° W longitude) and 38 sites in the western US (west of 110° W longitude). Only observations that were flagged as valid in the NTN data file were used in the performance analysis. Observations and model estimates are paired in time and space using the EPA's Site Compare programme, which is available for download as a tool from the Community Modelling and Analysis System (CMAS) website (<http://www.cmascenter.org>). Visualization of observations and model estimates, and computation of model performance statistics accomplished through the use of the Atmospheric Model Evaluation Tool (AMET; Appel et al., 2010), are available for download through the CMAS website.

2.5 Precipitation bias adjustment

At least some portion of the error present in the CMAQ estimated wet deposition is due to errors in the precipitation estimates from the meteorological model. Since both the NTN observed and MM5 estimated precipitation data are available for each NTN site, the modelled wet deposition can be adjusted to account for the error present in the model estimated precipitation. This adjustment is accomplished here by linearly adjusting the CMAQ estimated wet deposition by the ratio of the observed to estimated precipitation (see Eq. 1). For example, in the case where the observed precipitation is greater than the model estimated precipitation, the ratio is greater than one and, therefore, the model estimated wet deposition is increased.

$$\frac{\sum_{\text{Seasonal/Annual}} RT_{\text{Observed}}}{\sum_{\text{Seasonal/Annual}} RT_{\text{Modelled}}} \times \sum_{\text{Seasonal/Annual}} WD_{\text{Modelled}} = \text{Bias Adjusted } WD_{\text{Modelled}} \quad (1)$$

In Eq. (1), “RT” represents the seasonal/annual total accumulated precipitation (either observed or modelled), “WD” represents the seasonal/annual accumulated raw wet deposition estimate from the model and the “Bias Adjusted WD” is the precipitation bias adjusted seasonal/annual wet deposition estimate from the model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A multi-resolution assessment (CMAQ)

K. W. Appel et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The precipitation adjustment technique assumes that the observed to modelled precipitation ratio is well correlated with the observed to modelled deposition ratio. In other words, it is not assumed that the wet deposition scales linearly with precipitation, but only that the relationship between the errors in the model precipitation estimates and the error in the CMAQ deposition estimates is linear. Since the bias adjustment was applied over the aggregated seasonal and annual totals, there were no instances in which the observed precipitation was greater than zero while the model estimated precipitation was zero. However, in instances where there is observed precipitation but no model predicted precipitation, the current method of bias adjustment would keep the model estimated wet deposition zero for all species. The impact of the precipitation bias adjustment on model performance will be presented for each of the wet deposition species.

3 Assessment of CMAQ wet deposition performance

In order to provide a comprehensive assessment of the CMAQ wet deposition estimates, several different types of analyses will be presented. The performance of the model estimates are assessed on several time scales, including monthly, seasonally, annually and finally a multi-annual assessment of model performance. The performance for the 36-km and 12-km CMAQ simulations will be compared to examine how similar or dissimilar the model estimates are for a given time period. Since the 12-km CMAQ domain only covers the eastern two-thirds of the US, comparison to the 36-km results will be limited to the same geographic region (herein referred to as 36-km East). Results for the western one-third of the US will be limited to estimates from the 36-km CMAQ simulation (herein referred to as 36-km West) only, since no 12-km model data are available for the western US for the current analysis. The model estimates will also be examined spatially to identify regional biases.

3.1 Summary of precipitation performance

Simulated precipitation is a critical driver in the performance of the CMAQ-simulated wet deposition estimates, especially since large biases in model estimated precipitation can translate into biases in the CMAQ model estimates. Tables 1 and 2 present seasonal and annual normalized mean bias (NMB) and root mean square error (RMSE) for precipitation for the 12-km, 36-km East and 36-km West domains for the five years simulated. For the eastern US, the precipitation bias and error are lowest in the winter (December, January and February) and spring (March, April and May) seasons, when the majority of the precipitation is on the synoptic scale (i.e. large-scale frontal systems) and can generally be well resolved by the model. In the summer (June, July and August) and early fall (September, October and November), a large amount of the precipitation is sub-grid scale convective rain, which meteorological models tend to have difficulty representing accurately through the various parameterizations, which results in higher precipitation biases in those seasons. See Fig. S1 in the supplemental data for spatial plots of the NTN observed and MM5 estimated annual precipitation (12-km simulation only).

While the precipitation estimates for the 12-km and 36-km East simulations have similar patterns in their bias, the precipitation estimates for the 12-km simulation are consistently higher than those of the 36-km East simulation (indicated by systematically larger NMB values), which results in slightly larger biases in the winter, spring and summer for the 12-km simulation, but a smaller bias in the fall when precipitation is underestimated in both simulations. The bias and error in precipitation tend to be larger for the western US (based on the 36-km West simulation) than for the eastern US. The large bias is especially evident in the summer, when precipitation is grossly overestimated in the 36-km West simulation (summer average NMB = 54.5% for the five-year period). Overall, the annual NMB for the 12-km simulation was typically less than 5%, the exception being 2002 when the model precipitation estimates were biased significantly higher (NMB = 12.9%). The NMB for the 36-km East and 36-km West

GMDD

3, 2315–2360, 2010

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



simulations was typically slightly larger than the 12-km East simulation, with annual NMB generally ranging between $\pm 11\%$ for the five year period.

3.2 SO_4^- wet deposition

Model estimates from both the 12-km and 36-km simulations capture the seasonal trends in the observed monthly accumulated (accumulated over all sites) SO_4^- wet deposition for the 2002–2006 period, with the estimates from the 12-km CMAQ simulation consistently higher than those from the 36-km East simulation (Fig. 1). The CMAQ model generally overestimates SO_4^- wet deposition in the eastern US, with the 12-km simulation overestimating SO_4^- wet deposition for 50 of the 60 months, while the 36-km East simulation overestimates SO_4^- wet deposition for 33 of the 60 months. However, 88% of the estimates from the 36-km East simulation and 80% of the estimates from the 12-km simulation have a NMB of less than $\pm 15\%$ (Fig. 2). The largest overestimations of SO_4^- wet deposition occur in the late fall and winter, generally between October and March.

Overall, the bias in SO_4^- wet deposition estimates for the eastern US was relatively small for both the 12-km and 36-km East simulations (Table 3). The bias for the 12-km (36-km East) CMAQ simulation is highest in the winter, with the annual NMB ranging from 8.1% (–0.8%) to 30.7% (23.1%) and a five-year average NMB of 17.2% (9.0%). However, SO_4^- wet deposition is relatively small in the winter compared to the other seasons, so RMSE values in the winter are lower than the other seasons (Table 4). The NMB is smallest in the summer, ranging from 1.7% to 14.5% for the 12-km simulation (five-year average NMB = 5.2%) and 0.0% to 9.3% for the 36-km East simulation (five-year average NMB = –3.5%). The RMSE is largest in the summer, with annual RMSE values ranging between 1.6–2.1 kg/ha for the two simulations. Bias in the spring and fall periods generally falls between the performance for the summer and winter. The average annual NMB (RMSE) for the five-year period was 7.9% (3.56 kg/ha) for the

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



12-km simulation and 0.8% (3.10 kg/ha) for the 36-km East simulation, indicating SO_4^- wet deposition is generally overestimated, although only very slightly in the 36-km East simulation.

The SO_4^- wet deposition performance for the western US is considerably worse than for the eastern US, with the NMB exceeding 40% in 18 of the 60 (30%) months (Fig. 2). This result is not surprising given the challenging meteorological (recall the large precipitation biases in the western US) and air quality conditions that exist in the western US due to its complex topography. Also note that SO_4^- wet deposition in the western US is an order of magnitude less than that in the eastern US (Fig. 1), which may also contribute to the larger normalized bias. As was the case for the eastern US, the poorest model performance for the western US was in the winter, which had an average NMB of 31.6% (RMSE = 0.28 kg/ha) for the five-year period, while the summer had the best model performance, with a five-year average NMB of just 1.9% (RMSE = 0.25 kg/ha). The model bias was slightly higher in the spring (24.3%) than the fall (13.9%). The average NMB for the entire five-year period was 18.9% (RMSE = 0.82 kg/ha). Given the complexity of the terrain over much of the western US, a simulation utilizing finer grid spacing (e.g. 12-km) may result in improved performance, as some of the finer details of the topography would be captured in the modelling system.

Annual SO_4^- wet deposition is highest in the eastern half of the US where the largest SO_2 emissions occur (see Fig. S2 in the supplemental data). The highest amounts of SO_4^- wet deposition occur in the Ohio Valley and Great Lakes regions and stretching into parts of the Northeast. While these spatial features are well captured by the CMAQ model for all five years, the model tends to overestimate the annual SO_4^- wet deposition in the Ohio Valley region, with some model estimates exceeding 27 kg/ha in areas where observations indicate annual SO_4^- wet deposition of 19–20 kg/ha. The model also underestimates the SO_4^- wet deposition along parts of the coast of the Gulf of Mexico, although to varying degrees throughout the five-year period. Overall, the model captures the spatial variations in annual SO_4^- wet deposition.

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The change in annual SO_4^- wet deposition model bias as a result of applying the precipitation bias adjustment described in Sect. 2.5 for the 12-km simulation is shown in Fig. 3, which indicates at least some improvement in model bias for each of the five years by applying the precipitation bias adjustment. However, the improvement varies significantly from year to year, with the largest improvement in model performance for 2002 (annual NMB decreases from 21% to 2%), while for 2003 and 2006 the NMB improves by 3% or less. Spatially, the largest precipitation bias typically occurs in the Northeast and Great Lakes regions (particularly in 2002), and those regions show the largest improvement in bias and error as a result of the adjustment for precipitation bias (see Figs. S3 and S4 in the supplemental data for regional statistics).

A bootstrap sampling technique was used to test the robustness of the precipitation bias adjustment. For each year, the NTN observations were re-sampled with replacement 1000 times. The sample size for each of the 1000 samples matched the number of observations available for that year. The base model SO_4^- wet deposition estimates and precipitation bias corrected model estimates were matched to these pseudo-sets of observations, and the Root Mean Square Error (RMSE) for each sample was computed. The bootstrap distribution of RMSE values for the base model results and precipitation bias adjusted results is shown in Fig. 4. The largest decrease in RMSE occurs in 2002, 2004 and 2005, while the decrease in RMSE is much smaller in 2003 and 2006, which confirms that the precipitation bias adjustment significantly improves the model performance in 2002, but provides only a minor improvement in 2003 and 2006. The improvement in model performance gained by applying the precipitation bias adjustment is highly dependent on the performance of meteorological model estimates of precipitation, with greater improvement in model performance when the precipitation estimates are poor (as was the case in 2002).

3.3 NH_4^+ wet deposition

The pattern of NH_4^+ wet deposition closely follows the seasonal SO_4^- wet deposition pattern, with a peak in NH_4^+ wet deposition in the eastern US in the summer and a

NH_4^+ wet deposition each year, but consistently underestimates the magnitude of NH_4^+ wet deposition.

Unlike the SO_4^- wet deposition, applying the precipitation adjustment to the CMAQ estimated NH_4^+ wet deposition generally results in an increase in bias (Fig. 7) and a slight increase in error (Fig. 8) for each of the five years. This suggests that the overestimation in model-estimated precipitation is at least partially compensating for an underestimation in NH_4^+ wet deposition. The increase in bias is largest in 2002, where the NMB increases from -3% to -19% , while for the other years the increase in bias is smaller, generally ranging from 3% to 7% (see Fig. S6 in the supplemental data). It is important to note that the NH_3 emissions used in the CMAQ model simulation are constrained using the results of inverse modelling, so some increase in NH_4^+ wet deposition bias is expected when the model estimates are adjusted for precipitation bias.

The underestimation in NH_4^+ wet deposition may be due, in large part, to the poor temporal and spatial representation of NH_3 emissions, particularly those emissions associated with fertilizer applications and bi-directional exchange of NH_3 from soil and vegetation surfaces. In order to improve the NH_3 emissions, a bi-directional NH_3 exchange mechanism was developed for the CMAQ model which was in turn coupled with an agricultural management tool and a soil nitrogen geochemical cycling model to estimate NH_3 emissions from fertilized croplands (Cooter et al., 2010). The agricultural management tool estimates fertilizer application as a function of crop nutrient demand and the soil geochemical model was used to estimate the nitrification and denitrification processes in the soil column and provided the soil water solution ammonium and hydrogen ion concentrations needed in the bi-directional NH_3 model. Agricultural land use categories and crop profiles were proven by the US Department of Agriculture's 2002 Census of Agriculture (2002 Census of Agriculture, 2004).

To evaluate the impact that bi-directional NH_3 exchange has on the CMAQ estimated NH_4^+ wet deposition, a 2002 12-km eastern US CMAQ simulation that included bi-directional exchange was performed and the results were corrected for precipitation

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



bias (Fig. 9). Including the bi-directional exchange which significantly reduces the bias in the precipitation corrected annual NH_4^+ wet deposition, with the NMB reduced more than a factor of three (from -19% to -6%). The reduction in the model bias was due to improving the temporal resolution of NH_3 emissions from a monthly profile to hourly, representing grid cell level spatial variability instead of county level and modelling the soil nitrification, de-nitrification, vegetative uptake and soil evasion of NH_3 following fertilizer application rather than using state level fertilizer sales as a surrogate for emissions. It is anticipated that a beta version of the bi-directional NH_3 exchange will be available for the next version of the CMAQ model.

3.4 NO_3^- wet deposition

The NO_3^- wet deposition performance is dominated by large underestimations in the summer (Fig. 10), which is consistent with the performance of CMAQ model estimates of aerosol fine particulate NO_3^- (Appel et al., 2008). The CMAQ model estimates of NO_3^- wet deposition for the fall and winter seasons are relatively consistent for the eastern US, with the NMB ranging between $\pm 20\%$ for both the 12-km and 36-km East CMAQ simulations (Fig. 11). In the spring, NO_3^- wet deposition is underestimated in the eastern US, with an average NMB of -14.5% (RMSE = 0.88 kg/ha) and -22.6% (RMSE = 0.95 kg/ha) for the 12-km and 36-km East CMAQ simulations, respectively (Tables 7 and 8). For the western US the NMB is unbiased in the spring. For the summer, the NO_3^- wet deposition is largely underestimated for both the eastern and western US, with a NMB greater than -40% for all three domains, while the RMSE is roughly 1.5 kg/ha for the eastern US and 0.5 kg/ha for the western US. For the entire five-year period, the model underestimates NO_3^- wet deposition with a five-year average NMB of -14.9% (RMSE = 2.54 kg/ha) and -21.4% (RMSE = 2.70 kg/ha) for the 12-km and 36-km East simulations, respectively, and a NMB of -6.9% (RMSE = 1.00 kg/ha) for the 36-km West simulation.

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A multi-resolution assessment (CMAQ)

K. W. Appel et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the flashes observed by the National Lightning Detection Network (NLDN), where the NLDN cloud-to-ground (CG) flash rates are multiplied by $Z+1$ to account for the contribution of intracloud flashes (IC) to the total flash rate. Z is the climatological IC/CG ratio from Boccippio et al. (2001). This method captures the day-to-day variability in flash rates, while retaining an accurate estimate of the monthly total (Allen et al., 2009). For each flash, it is assumed that 500 moles of NO are produced (DeCaria et al., 2005; Ott et al., 2007), which is a reasonable mid-latitude value. The NO is vertically distributed from the surface to the model layer containing the convective cloud top using climatological vertical flash rate information from the Northern Alabama Lightning Mapping Array (Koshak et al., 2004).

For the summer of 2004, a CMAQ model simulation using 36-km grid spacing was performed for the CONUS that included lightning produced NO as described above. Over the entire summer, NO produced from lightning was equal to 30% of the anthropogenic NO emissions. Because most of the NO produced from lightning is created in the upper troposphere, the impact to surface concentrations is small, as in Kaynak et al. (2008). However, over the eastern US where lightning flash counts are greatest, the impact to NO_3^- wet deposition is substantial. Figure 14 shows the bias in NO_3^- wet deposition at NADP monitoring sites for the CMAQ simulation without lightning NO, including lightning NO, and including lightning NO and the precipitation bias adjustment. For the monitoring locations east of 100 degrees W longitude, the CMAQ simulation with the lightning NO production has a low bias and captures the range of variability shown at the surface monitors. At the monitors west of 100 degrees W longitude, the impact is small and the bias persists, owing to the low lightning flash counts in this region.

4 Summary

The CMAQ modelling system was used to estimate SO_4^{2-} , NH_4^+ and NO_3^- wet deposition for the years 2002–2006 for the CONUS using a 36-km grid spacing and the eastern

**A multi-resolution
assessment (CMAQ)**

K. W. Appel et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

US using a 12-km grid spacing. The resulting wet deposition estimates from the model were compared with surface based observations of wet deposition species available across the US from the NTN for the five-year period. For SO_4^- wet deposition, the operational performance of the CMAQ model estimates were generally comparable for the 36-km and 12-km simulations for the eastern US, with the 12-km simulation on average yielding slightly higher estimates of SO_4^- wet deposition than the 36-km simulation. When compared to observations from the NTN, the NMB for the CMAQ model estimates was slightly higher for the 12-km simulation, however, both simulations had annual NMB that were less than $\pm 15\%$ each year. Bias and error in the model SO_4^- wet deposition estimates were significantly reduced for three of the five years (smaller improvements for the other two years) when the estimates were adjusted to account for biases in the model estimated precipitation.

The CMAQ modelling system underestimates NH_4^+ wet deposition in the eastern US in both the 36-km and 12-km simulations, with the underestimation tending to be slightly larger in the 36-km simulation. The largest underestimation of NH_4^+ wet deposition occurs in the winter and spring periods, while the summer and fall have slightly lower underestimations. The underestimation is likely due, in part, to the poor temporal and spatial representation of NH_3 emissions, particularly those emissions associated with fertilizer applications and bi-directional exchange of NH_3 flux from the soil and vegetation. Implementation of a bi-directional NH_3 flux mechanism in the CMAQ model, along with improvements in the temporal and spatial representation of fertilizer applications, improve the underestimation of NH_4^+ wet deposition and these changes will likely be included in the next release of the CMAQ model.

The performance for model estimates of NO_3^- wet deposition are mixed throughout the year, with the model largely underestimating NO_3^- wet deposition in the spring and summer in the eastern US, while the bias in the fall and winter is relatively small. Model estimates of NO_3^- wet deposition tend to be slightly lower for the 36-km simulation as compared to the 12-km simulation, particularly in the spring. One large source of the underestimation of NO_3^- wet deposition is from a lack of NO produced from

lightning in the upper troposphere, which can be a large source of NO, particularly in the summer in the eastern US when lightning activity is the high. CMAQ model simulations, that include the production of NO from lightning, show a substantial reduction in the NO₃⁻ wet deposition underestimation in the eastern US in the summer as compared to simulations without lightning NO. There is little impact on bias in the western US when lightning generated NO is included due to the relatively low amount of lightning activity in the western US.

Overall, the performance for the 36-km and 12-km CMAQ model simulations was similar for the eastern US, while for the western US the performance of the 36-km simulation was generally not as good as either eastern US simulation. On an annual basis, the model performance for all three wet deposition species was relatively consistent (NMB <30%), with mostly small variations in normalized bias (standard deviation <3%) over the five-year period for the eastern US. Annual variations in NMB were larger for the western US, with a standard deviation >5.5%. This suggests that the modelling system handles the year-to-year variability relatively well in meteorology and emissions that occur over longer periods of time, particularly for the eastern US. As annual air quality model simulations become more routine, it is likely that the five-year performance assessment presented here could be extended to cover a longer time-period (e.g. a decade). Additionally, expanding the 12-km simulation to include the western US may result in improved model performance over the 36-km simulation given the complexity of the terrain in the western US.

Supplementary material related to this article is available online at:
<http://www.geosci-model-dev-discuss.net/3/2315/2010/gmdd-3-2315-2010-supplement.zip>

Acknowledgements. The authors would like to thank Lara Reynolds, Nancy Hwang, Lucille Bender and other members of Computer Sciences Corporation for their help in performing the MM5 and CMAQ simulations used in this work.

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Disclaimer

The United States Environmental Protection Agency, through its Office of Research and Development, funded and managed the research described here. It has been subjected to Agency review and approved for publication.

References

Allen, D. J., Pickering, K., Pinder, R. W., and Pierce, T.: Impact of Lightning-NO emissions on Eastern US Photochemistry during the Summer of 2004 as Determined using the CMAQ Model. Presented at the 8th Annual CMAS Conference, Chapel Hill, NC, 19–21 October 2009.

Appel, K. W., Bhave, P. V., Gilliland, A. B., Sarwar, G., and Roselle, S. J.: Evaluation of the Community Multiscale Air Quality (CMAQ) model version 4.5: Sensitivities impacting model performance; Part II–particulate matter, *Atmos. Environ.*, 42, 6057–6066, 2008.

Appel, K. W., Gilliam, R. C., Davis, N., and Zubrow, A.: Overview of the Atmospheric Model Evaluation Tool (AMET) v1.1 for evaluating meteorological and air quality models, accepted for publication in *Environ. Modell. Softw.*, 2010.

Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and Schultz, M. G.: Global modelling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res.*, 106, 23073–23096, 2001.

Boccippio, D., Cummings, K., Christian, H., and Goodman, S.: Combined satellite- and surface-based estimation of the intracloud-cloud-to-ground lightning ratio over the continental United States, *Mon. Weather Rev.*, 129, 108–122, 2001.

Byun, D. W. and Schere, K. L.: Review of the governing equations, computational algorithms, and other components of the Models-3 Community Multiscale Air Quality (CMAQ) modelling system, *Appl. Mech. Rev.*, 55, 51–77, 2006.

Carlton, A. G., Bhave, P. V., Napelenok, S. L., Edney, E. O., Sarwar, G., Pinder, R. W., Pouliot, G. A., and Houyoux, M.: Model representation of secondary organic aerosol in CMAQv4.7, *Environ. Sci. Technol.*, 44, 8553–8560, 2010.

Cooter, E., Bash, J. O., Walker, J. T., Jones, M. R., and Robarge, W.: Estimation of NH₃ bi-directional flux over managed agricultural soils, *Atmos. Environ.*, 44, 2107–2115, 2010.

GMDD

3, 2315–2360, 2010

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DeCaria, A. J., Pickering, K. E., Stenchikov, G. L., and Ott, L. E.: Lightning-generated NO_x and its impact on tropospheric ozone production: A three-dimensional modelling study of a STERAO-A thunderstorm, *J. Geophys. Res.*, 110, D14303, doi:10.1029/2004JD005556, 2005.

5 Driscoll, C. T., Lawrence, G. B., Bulger, A. J., Butler, T. J., Cronan, C. S., Eagar, C., Lambert, K. F., Likens, G. E., Stoddard, J. L., and Weathers, K. C.: Acidic Deposition in the Northeastern United States: Sources and Inputs, Ecosystem Effects, and Management Strategies, *Bioscience*, 51(3), 180–198, 2001.

10 Driscoll, C. T., Whitall, D., Aber, J., Boyer, E., Castro, M., Cronan, C., Goodale, C. L., Groffman, P., Hopkinson, C., Lambert, K., Lawrence, G., and Ollinger, S.: Nitrogen Pollution in the Northeastern United States: Sources, Effects, and Management Options, *Bioscience*, 53(4), 357–374, 2003.

Dudhia, J.: Numerical study of convection observed during the winter monsoon experiment using a mesoscale two-dimensional model, *J. Atmos. Sci.*, 46, 3077–3107, 1989.

15 Fang, Y., Fiore, A. M., Horowitz, L. W., Levy II, H., Hu, Y., and Russell, A. G.: Sensitivity of the NO_y budget over the United States to anthropogenic and lightning NO_x in summer, *J. Geophys. Res.*, 115, D18312, doi:10.1029/2010JD014079, 2010.

Fenn, M. E., Baron, J. S., Allen, E. B., Rueth, H. M., Nydick, K. R., Geiser, L., Bowman, W. D., Sickman, J. O., Meixner, T., Johnson, D. W., and Neitlich, P.: Ecological Effects of Nitrogen Deposition in the Western United States, *Bioscience*, 53(4), 404–420, 2003.

20 Foley, K. M., Roselle, S. J., Appel, K. W., Bhawe, P. V., Pleim, J. E., Otte, T. L., Mathur, R., Sarwar, G., Young, J. O., Gilliam, R. C., Nolte, C. G., Kelly, J. T., Gilliland, A. B., and Bash, J. O.: Incremental testing of the Community Multiscale Air Quality (CMAQ) modelling system version 4.7, *Geosci. Model Dev.*, 3, 205–226, doi:10.5194/gmd-3-205-2010, 2010.

25 Geiser, L. H., Jovan, S. E., Glavich, D. A., and Porter, M. K.: Lichen-based critical loads for atmospheric nitrogen deposition in Western Oregon and Washington Forests, USA, *Environ. Pollut.*, 158, 2412–2442, 2010.

Gilliland, A. B., Appel, K. W., Pinder, R., and Dennis, R. L.: Seasonal NH_3 emissions for the continental United States: inverse model estimation and evaluation, *Atmos. Environ.*, 40, 4986–4998, 2006.

30 Goebes, M. D., Strader, R., and Davidson, C.: An ammonia emission inventory for fertilizer application in the US, *Atmos. Environ.*, 37, 2539–2550, 2003.

Grell, G. A., Dudhia, A. J., and Stauffer, D. R.: A description of the Fifth-Generation

**A multi-resolution
assessment (CMAQ)**

K. W. Appel et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

PennState/NCAR Mesoscale Model (MM5). NCAR Technical Note NCAR/TN-398+STR, available at <http://www.mmm.ucar.edu/mm5/doc1.html>, 1994.

Houyoux, M. R., Vukovich, J. M., Coats Jr., C. J., Wheeler, N. J. M., Kasibhatla, P.: Emission inventory development and processing for the seasonal model for regional air quality, *J. Geophys. Res.*, 105 (D7), 9079–9090, 2000.

Kain, J. S.: The Kain-Fritsch convective parameterization: An update, *J. Appl. Meteor.*, 43, 170–181, 2004.

Kaynak, B., Hu, Y., Martin, R. V., Russell, A. G., Choi, Y., and Wang, Y.: The effect of lightning NO_x production on surface ozone in the continental United States, *Atmos. Chem. Phys.*, 8, 5151–5159, doi:10.5194/acp-8-5151-2008, 2008.

Koshak, W. J., Solakiewicz, R. J., Blakeslee, R. J., Goodman, S. J., Christian, H. J., Hall, J. M., Bailey, J. C., Krider, E. P., Bateman, M. G., Boccippio, D. J., Mach, D. M., McCaul, E. W., Stewart, M. F., Buechler, D. E., Petersen, W. A., and Cecil, D. J.: North Alabama Lightning Mapping Array (LMA): VHF Source Retrieval Algorithm and Error Analyses, *J. Atmos. Ocean. Tech.*, 21, 543–558, 2004.

Lovett, G. M. and Tear, T. H.: Threat from Above: Air Pollution impacts on Ecosystems and Biological Diversity in the Eastern United States. The Nature Conservancy and the Cary institute of Ecosystem Studies (www.ecostudies.org/reprints/Threats_from_above.pdf), 2008.

Mlawer, E. J., Taubman, S. J., Brown, P. D., Iacono, M. J., and Clough, S. A.: Radiative transfer for inhomogeneous atmosphere: RRTM, a validated correlated-k model for the long-wave, *J. Geophys. Res.*, 102(D14), 16663–16682, 1997.

Ott, L. E., Pickering, K. E., Stenchikov, G. L., Huntrieser, H., and Schumann, U.: Effects of lightning NO_x production during the 21 July European Lightning Nitrogen Oxides Project storm studied with a three-dimensional cloud-scale chemical transport model, *J. Geophys. Res.*, 112, D05307, doi:10.1029/2006JD007365, 2007.

Otte, T. L., Pouliot, G., Pleim, J. E., Young, J. O., Schere, K. L., Wong, D. C., Lee, P. C. S., Tsidulko, M., McQueen, J. T., Davidson, P., Mathur, R., Chuang, H. Y., DiMego, G., and Seaman, N. L.: Linking the Eta model with the Community Multiscale Air Quality (CMAQ) modelling system to build a national air quality forecasting system, *Weather Forecast.*, 20, 367–384, 2005.

Pleim, J. E.: A combined local and nonlocal closure model for the atmospheric boundary layer. Part I: model description and testing, *J. Appl. Meteor. Clim.*, 46, 1383–1395, 2007a.

Pleim, J. E.: A combined local and nonlocal closure model for the atmospheric boundary layer.

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Part II: application and evaluation in a mesoscale meteorological model, *J. Appl. Meteor. Clim.*, 46, 1396–1409, 2007b.

Pleim, J. E. and Xiu, A.: Development and testing of a surface flux and planetary boundary layer model for application in mesoscale models, *J. Appl. Meteor.*, 34, 16–32, 1995.

5 Reisner, J., Rasmussen, R. M., and Bruintjes, R. T.: Explicit forecasting of supercooled liquid water in winter storms using the MM5 mesoscale model, *Q. J. Roy. Meteor. Soc.*, 124, 1071–1107, 1998.

Xiu, A. and Pleim, J. E.: Development of a land-surface model. Part I: application in a mesoscale meteorological model, *J. Appl. Meteor.*, 40, 192–209, 2001.

10 Yarwood, G., Roa, S., Yocke, M., and Whitten, G.: Updates to the carbon bond chemical mechanism: CBo5. Final report to the US EPA, RT-0400675, available at <http://www.camx.com>, 2005.

2002 Census of Agriculture: US Department of Agriculture, US Summary and State Data, vol. 1, Geographic Area Series Part 51, AC-02-A-51, National Agricultural Statistics Service, 2004.

15

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Seasonal and annual NMB (%) for precipitation for the 12-km and 36-km CMAQ model simulations.

	CMAQ Domain	2002	2003	2004	2005	2006	Five-year average
Winter	12-km	-0.4	-1.8	-1.4	-1.9	-1.8	-1.5
	36-km East	-2.6	-7.1	-4.8	-4.9	-10.8	-6.0
	36-km West	-10.0	0.6	-3.8	-3.6	-1.4	-3.6
Spring	12-km	20.2	0.5	9.3	4.9	12.8	9.5
	36-km East	8.9	-6.8	-1.6	-5.6	0.8	-0.9
	36-km West	9.7	-1.7	24.2	8.7	20.8	12.3
Summer	12-km	44.8	12.3	20.2	23.9	15.0	23.2
	36-km East	42.2	6.2	8.4	16.3	0.4	14.7
	36-km West	64.3	85.3	43.9	49.5	29.7	54.5
Fall	12-km	-16.9	-15.5	-16.1	-20.7	-15.4	-16.9
	36-km East	-16.6	-20.0	-18.4	-22.1	-22.2	-19.9
	36-km West	-11.6	8.2	-7.8	9.5	14.2	2.5
Annual	12-km	12.9	-0.1	4.1	2.4	2.4	4.3
	36-km East	9.0	-6.0	-3.5	-3.2	-8.4	-2.4
	36-km West	0.5	5.7	5.8	10.7	10.9	6.7

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 2. Seasonal and annual RMSE (cm) for precipitation for the 12-km and 36-km CMAQ model simulations.

	CMAQ Domain	2002	2003	2004	2005	2006	Five-year average
Winter	12-km	7.6	7.4	6.6	7.9	5.6	7.02
	36-km East	7.4	7.9	6.7	5.7	6.4	6.82
	36-km West	14.7	11.8	11.3	7.6	11.0	11.3
Spring	12-km	8.3	9.8	8.1	8.8	7.1	8.42
	36-km East	7.2	9.5	7.6	8.9	7.0	8.04
	36-km West	7.8	14.9	7.4	10.3	10.4	10.2
Summer	12-km	18.4	13.1	17.0	15.4	13.9	15.6
	36-km East	17.0	13.2	14.4	13.4	11.5	13.9
	36-km West	11.2	9.5	9.0	9.6	4.5	8.76
Fall	12-km	10.8	8.7	10.9	10.6	9.5	10.1
	36-km East	10.4	9.4	10.1	10.8	10.5	10.2
	36-km West	7.9	7.8	10.1	10.0	8.0	8.76
Annual	12-km	25.9	24.0	24.2	24.4	22.8	24.3
	36-km East	23.2	25.3	23.6	23.2	25.1	24.1
	36-km West	32.5	36.1	31.7	29.1	27.1	31.3

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Table 3. Seasonal and annual NMB (%) for SO_4^- wet deposition for the 12-km and 36-km CMAQ model simulations.

	CMAQ Domain	2002	2003	2004	2005	2006	Five-year average
Winter	12-km	8.1	12.7	26.4	30.7	8.1	17.2
	36-km East	−0.8	5.2	16.3	23.1	1.0	9.0
	36-km West	14.1	49.7	39.4	32.5	22.1	31.6
Spring	12-km	8.1	2.8	7.8	3.5	3.8	5.2
	36-km East	−0.6	−4.5	−1.3	−5.3	−5.8	−3.5
	36-km West	27.7	29.3	38.5	2.5	23.6	24.3
Summer	12-km	14.5	3.9	8.1	1.7	2.1	6.1
	36-km East	9.3	0.0	2.6	−2.4	−3.6	1.2
	36-km West	8.7	−9.8	25.8	11.5	−26.8	1.9
Fall	12-km	11.5	12.2	13.3	−1.8	7.2	8.5
	36-km East	5.9	5.9	5.1	−7.9	−1.4	1.4
	36-km West	−4.8	38.0	13.0	19.1	4.0	13.9
Annual	12-km	11.0	6.4	11.4	6.0	4.6	7.9
	36-km East	4.2	0.5	3.7	−1.5	−3.0	0.8
	36-km West	12.6	29.9	28.4	13.0	10.8	18.9

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Table 4. Seasonal and annual RMSE (kg/ha) for SO_4^- wet deposition for the 12-km and 36-km CMAQ model simulations.

	CMAQ Domain	2002	2003	2004	2005	2006	Five-year average
Winter	12-km	0.93	0.75	0.96	1.21	0.75	0.92
	36-km East	0.72	0.64	0.74	0.71	0.64	0.69
	36-km West	0.33	0.29	0.30	0.21	0.27	0.28
Spring	12-km	1.21	1.49	1.37	1.30	0.96	1.27
	36-km East	1.05	1.35	1.18	1.19	0.84	1.12
	36-km West	0.45	0.40	0.30	0.34	0.34	0.37
Summer	12-km	1.90	1.95	2.07	1.66	1.72	1.86
	36-km East	1.88	1.89	1.77	1.63	1.70	1.77
	36-km West	0.26	0.30	0.27	0.24	0.20	0.25
Fall	12-km	1.33	1.03	1.13	1.09	1.12	1.14
	36-km East	1.20	0.97	0.85	1.00	1.04	1.01
	36-km West	0.22	0.26	0.23	0.27	0.22	0.24
Annual	12-km	3.85	3.59	3.79	3.62	2.94	3.56
	36-km East	3.14	3.36	3.25	2.94	2.82	3.10
	36-km West	0.82	0.94	0.83	0.71	0.82	0.82

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


**A multi-resolution
assessment (CMAQ)**

K. W. Appel et al.

Table 5. Seasonal and annual NMB (%) for NH_4^+ wet deposition for the 12-km and 36-km CMAQ model simulations.

	CMAQ Domain	2002	2003	2004	2005	2006	Five-year average
Winter	12-km	-19.4	-18.3	-13.3	2.0	-18.9	-13.6
	36-km East	-23.5	-25.0	-18.9	1.5	-21.7	-17.5
	36-km West	-39.0	-41.5	-35.6	-42.2	-27.2	-37.1
Spring	12-km	-13.5	-28.1	-17.7	-20.0	-20.4	-19.9
	36-km East	-16.8	-30.5	-22.1	-24.5	-23.9	-23.6
	36-km West	-2.5	-19.7	0.8	-5.2	9.4	-3.4
Summer	12-km	-7.8	-8.6	-2.2	-7.8	-10.4	-7.4
	36-km East	-8.0	-8.0	-2.2	-8.3	-11.9	-7.7
	36-km West	-19.3	-43.4	10.3	0.3	-41.4	-18.7
Fall	12-km	-8.6	-3.5	-6.5	-20.5	-8.5	-9.5
	36-km East	-11.9	-6.2	-9.7	-20.6	-11.8	-12.0
	36-km West	-42.3	14.6	-9.4	23.0	-22.7	-7.4
Annual	12-km	-11.2	-16.0	-9.8	-13.2	-14.0	-12.8
	36-km East	-13.4	-17.9	-12.5	-15.5	-16.6	-15.2
	36-km West	-25.0	-23.5	-9.6	-5.4	-15.2	-15.7

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 6. Seasonal and annual RMSE (kg/ha) for NH_4^+ wet deposition for the 12-km and 36-km CMAQ model simulations.

	CMAQ Domain	2002	2003	2004	2005	2006	Five-year average
Winter	12-km	0.15	0.14	0.24	0.15	0.15	0.17
	36-km East	0.14	0.13	0.23	0.10	0.14	0.15
	36-km West	0.11	0.16	0.21	0.14	0.11	0.15
Spring	12-km	0.34	0.49	0.37	0.41	0.31	0.38
	36-km East	0.35	0.49	0.37	0.41	0.30	0.38
	36-km West	0.16	0.26	0.16	0.20	0.24	0.20
Summer	12-km	0.44	0.45	0.41	0.41	0.45	0.43
	36-km East	0.44	0.42	0.41	0.42	0.45	0.43
	36-km West	0.12	0.22	0.15	0.14	0.15	0.16
Fall	12-km	0.22	0.17	0.20	0.22	0.22	0.21
	36-km East	0.21	0.17	0.20	0.22	0.22	0.20
	36-km West	0.14	0.14	0.20	0.14	0.16	0.16
Annual	12-km	0.76	0.86	0.82	0.80	0.77	0.80
	36-km East	0.74	0.84	0.82	0.78	0.79	0.79
	36-km West	0.36	0.56	0.53	0.49	0.46	0.48

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Table 7. Seasonal and annual NMB (%) for NO_3^- wet deposition for the 12-km and 36-km CMAQ model simulations.

	CMAQ Domain	2002	2003	2004	2005	2006	Five-year average
Winter	12-km	12.3	10.1	16.9	20.6	8.8	13.7
	36-km East	3.9	0.5	7.4	12.0	1.8	5.1
	36-km West	5.8	21.6	24.9	11.2	17.2	16.1
Spring	12-km	-8.7	-13.3	-15.3	-15.6	-19.7	-14.5
	36-km East	-16.4	-20.9	-23.6	-24.2	-28.1	-22.6
	36-km West	-7.3	-2.7	-6.6	-1.3	18.1	0.0
Summer	12-km	-38.0	-39.4	-38.7	-39.9	-45.4	-40.3
	36-km East	-40.3	-41.9	-43.2	-43.4	-49.9	-43.7
	36-km West	-49.6	-62.0	-36.2	-26.4	-63.9	-47.6
Fall	12-km	3.7	2.4	11.5	-9.0	-1.1	1.5
	36-km East	-3.4	-4.5	3.0	-14.1	-9.2	-5.6
	36-km West	-29.0	16.3	-6.2	9.2	-16.7	-5.3
Annual	12-km	-12.5	-15.6	-12.8	-14.6	-19.7	-15.0
	36-km East	-18.4	-21.6	-20.1	-23.1	-26.4	-21.9
	36-km West	-18.0	-6.0	-4.7	-1.8	-7.4	-7.6

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


A multi-resolution assessment (CMAQ)

K. W. Appel et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 8. Seasonal and annual RMSE (kg/ha) for NO_3^- wet deposition for the 12-km and 36-km CMAQ model simulations.

	CMAQ Domain	2002	2003	2004	2005	2006	Five-year average
Winter	12-km	1.10	0.86	1.10	1.14	0.81	1.00
	36-km East	0.90	0.80	0.80	0.67	0.68	0.77
	36-km West	0.26	0.32	0.39	0.28	0.34	0.32
Spring	12-km	0.83	1.02	0.90	0.90	0.76	0.88
	36-km East	0.88	1.08	0.98	0.97	0.83	0.95
	36-km West	0.26	0.45	0.22	0.37	0.61	0.38
Summer	12-km	1.56	1.55	1.40	1.27	1.54	1.46
	36-km East	1.62	1.64	1.48	1.34	1.62	1.54
	36-km West	0.46	0.49	0.50	0.35	0.55	0.47
Fall	12-km	1.10	0.74	0.80	0.59	0.69	0.78
	36-km East	0.91	0.75	0.72	0.56	0.70	0.73
	36-km West	0.31	0.31	0.32	0.26	0.26	0.29
Annual	12-km	2.76	2.63	2.45	2.43	2.42	2.54
	36-km East	2.75	2.95	2.62	2.40	2.76	2.70
	36-km West	0.80	1.11	1.00	0.90	1.19	1.00




Fig. 1. Monthly accumulated (across all sites) SO_4^- wet deposition (kg/ha) for the eastern US NTN observations (black diamonds), 12-km CMAQ simulation (red squares), 36-km East CMAQ simulation (blue triangles), western US NTN observations (dashed; green diamonds) and 36-km West CMAQ (dashed; yellow triangles). The scale for the western US values is given on the right y-axis.

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A multi-resolution
assessment (CMAQ)

K. W. Appel et al.




Fig. 2. SO₄⁼ wet deposition NMB for the 12-km CMAQ simulation (red diamonds), 36-km East CMAQ simulation (blue squares) and the 36-km West CMAQ simulation (dashed; yellow triangles).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A multi-resolution
assessment (CMAQ)

K. W. Appel et al.




Fig. 3. Box plots of annual modelled – observed SO₄²⁻ wet deposition for model wet deposition estimates without any adjustment for precipitation bias (“Base Model”; blue) and for the model estimates adjusted for precipitation errors (“Precip. Adjusted”; red). The black line within the box represents the median bias, shading represents the range of the 25% to 75% quartile and the dashed lines represent the range of the 5% to 95% values.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A multi-resolution assessment (CMAQ)

K. W. Appel et al.




Fig. 4. Distribution of RMSE based on 1000 bootstrap samples of the modelled and observed SO_4^- wet deposition. Results for model estimates without any adjustment for precipitation bias (“Base Model”) are shown in blue and for model estimates adjusted for precipitation errors (“Precip. Adj.”) are red. The bold lines indicate the RMSE values from the original dataset.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 5. Monthly accumulated (across all sites) NH_4^+ wet deposition (kg/ha) for the eastern US NTN observations (black diamonds), 12-km CMAQ simulation (red squares), 36-km East CMAQ simulation (blue triangles), western US NTN observations (dashed; green diamonds) and 36-km West CMAQ (dashed; yellow triangles). The scale for the western US values is given on the right y-axis.

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. NH₄⁺ wet deposition NMB for the 12-km CMAQ simulation (red diamonds), 36-km East CMAQ simulation (blue squares) and the 36-km West CMAQ simulation (dashed; yellow triangles).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A multi-resolution
assessment (CMAQ)

K. W. Appel et al.




Fig. 8. Distribution of RMSE based on 1000 bootstrap samples of the modelled and observed NH_4^+ wet deposition. Results for model estimates without any adjustment for precipitation bias (“Base Model”) are shown in blue and for model estimates adjusted for precipitation errors (“Precip. Adj.”) are red. The bold lines indicate the RMSE values from the original dataset.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2002: Modeled – Observed Wet Deposition NH₄




Fig. 9. Box plots of modelled – observed NH₄⁺ wet deposition for the eastern US (12-km CMAQ simulation only) for 2002. Shown are the model NH₄⁺ wet deposition biases for the base CMAQ simulation (“Base Model”; light blue), the base simulation with precipitation bias adjustment (“Precip. Adjusted Base”; red), the simulation with bi-directional NH₃ flux only (“Bidi NH3”; dark blue) and the simulation with both precipitation bias adjusted NH₄⁺ wet deposition and bi-directional NH₃ flux included (“Precip. Adjusted Bidi NH3”; dark red).

A multi-resolution assessment (CMAQ)

K. W. Appel et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

⏪

⏩

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



A multi-resolution assessment (CMAQ)

K. W. Appel et al.




Fig. 10. Monthly accumulated (across all sites) NO_3^- wet deposition (kg/ha) for the eastern US NTN observations (black diamonds), 12-km CMAQ simulation (red squares), 36-km East CMAQ simulation (blue triangles), western US NTN observations (dashed; green diamonds) and 36-km West CMAQ (dashed; yellow triangles). The scale for the western US values is given on the right y-axis.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**A multi-resolution
assessment (CMAQ)**

K. W. Appel et al.




Fig. 11. NO₃⁻ wet deposition NMB for the 12-km CMAQ simulation (red diamonds), 36-km East CMAQ simulation (blue squares) and the 36-km West CMAQ simulation (dashed; yellow triangles).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A multi-resolution
assessment (CMAQ)

K. W. Appel et al.




Fig. 12. Box plots of annual modelled – observed NO₃⁻ wet deposition for model wet deposition estimates without any adjustment for precipitation bias (“Base Model”; blue) and for the model estimates adjusted for precipitation errors (“Precip. Adjusted”; red). The black line within the box represents the median bias, shading represents the range of the 25% to 75% quartile and the dashed lines represent the range of the 5% to 95% values.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

A multi-resolution
assessment (CMAQ)

K. W. Appel et al.




Fig. 13. Distribution of RMSE based on 1000 bootstrap samples of the modelled and observed NO₃⁻ wet deposition. Results for model estimates without any adjustment for precipitation bias (“Base Model”) are shown in blue and for model estimates adjusted for precipitation errors (“Precip. Adj.”) are red. The bold lines indicate the RMSE values from the original dataset.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



